

CATALOGUE OF INVERTEBRATE AND VERTEBRATE
Paleontological Type Specimens
OF THE HUNGARIAN NATURAL HISTORY MUSEUM



József PÁLFY
Alfréd DULAI
Mihály GASPARIK
Péter OZSVÁRT
Piroska PAZONYI
Ottilia SZIVES

Hungarian Natural History Museum
Budapest, 2008

Catalogue of Invertebrate and Vertebrate
Paleontological Type Specimens
of the Hungarian Natural History Museum

József PÁLFY, Alfréd DULAI, Mihály GASPARIK,
Péter OZSVÁRT, Piroska PAZONYI & Ottília SZIVES

**Catalogue of Invertebrate and Vertebrate
Paleontological Type Specimens
of the Hungarian Natural History Museum**

József PÁLFY, Alfréd DULAI, Mihály GASPARIK,
Péter OZSVÁRT, Piroska PAZONYI & Ottília SZIVES

Hungarian Natural History Museum
Budapest, 2008

Publication of this volume was supported by the National Cultural Fund



Authors' affiliations and addresses:

Department of Geology and Paleontology
Hungarian Natural History Museum
POB 137, Budapest, H-1431 Hungary
palfy@nhmus.hu, dulai@nhmus.hu, gasparik@nhmus.hu, sziveso@nhmus.hu

Research Group for Paleontology
Hungarian Academy of Sciences – Hungarian Natural History Museum
POB 137, Budapest, H-1431 Hungary
palfy@nhmus.hu, ozsi@nhmus.hu, pazonyi@nhmus.hu

Front cover: Holotype skull of *Iharkutosuchus makadii* ŐSI, CLARK & WEISHAMPEL, 2007. The recently discovered Late Cretaceous locality of Iharkút (Bakony Mts., Hungary) furnished type specimens of several new reptile species which are considered the most valuable new acquisitions to our collection after 2000.

Back cover: Holotype of *Pecten fotensis* CSEPREGHY-MEZNERICS, 1960. This Miocene bivalve species was described by a former head of the Department of Geology and Paleontology. With a long history of research, Tertiary mollusks are the best represented group within the type collection.

Additional copies may be obtained from:

The Library of the Hungarian Natural History Museum
POB 137, Budapest, H-1431 Hungary
Fax: +36 1 317 2320, e-mail: libr@nhmus.hu

Published by the Hungarian Natural History Museum
Responsible publisher: István Matskási
www.nhmus.hu

ISBN 978-963-9877-02-3

© Hungarian Natural History Museum, 2008

Printed in Hungary by *mondAt L&d;*, Budapest, www.mondat.hu

Contents

Introduction	7
History and development of the type catalogue	7
Notes on structure and use of type catalogue	9
Acknowledgements	14
Catalogue of invertebrate type specimens	17
1. Paleozoic types	17
1.1. Paleozoic Foraminiferida	17
1.2. Paleozoic Anthozoa	18
1.3. Paleozoic Bivalvia	19
1.4. Paleozoic Gastropoda	19
1.5. Paleozoic Cephalopoda	19
1.6. Paleozoic Brachiopoda	20
1.7. Paleozoic Trilobita	20
2. Triassic types	20
2.1. Triassic Radiolaria	20
2.2. Triassic Bivalvia	27
2.3. Triassic Gastropoda	30
2.4. Triassic Cephalopoda	30
2.5. Triassic Brachiopoda	31
2.6. Triassic Ostracoda	31
3. Jurassic types	31
3.1. Jurassic Foraminiferida	31
3.2. Jurassic Anthozoa	31
3.3. Jurassic Bivalvia	32
3.4. Jurassic Gastropoda	33
3.5. Jurassic Cephalopoda	34
3.6. Jurassic Brachiopoda	36
4. Cretaceous types	37
4.1. Cretaceous Anthozoa	37
4.2. Cretaceous Bivalvia	38
4.3. Cretaceous Gastropoda	38
4.4. Cretaceous Cephalopoda	39
5. Paleocene types	40
5.1. Paleocene Brachiopoda	40

6.	Eocene types	41
6.1.	Eocene Foraminiferida	41
6.2.	Eocene Anthozoa	48
6.3.	Eocene Bivalvia	49
6.4.	Eocene Scaphopoda	50
6.5.	Eocene Gastropoda	51
6.6.	Eocene Cephalopoda	60
6.7.	Eocene Brachiopoda	60
6.8.	Eocene Cirripedia	61
6.9.	Eocene Decapoda	61
6.10.	Eocene Echinoidea	66
7.	Oligocene types	67
7.1.	Oligocene Foraminiferida	67
7.2.	Oligocene Bivalvia	70
7.3.	Oligocene Scaphopoda	81
7.4.	Oligocene Gastropoda	81
7.5.	Oligocene Cephalopoda	94
7.6.	Oligocene Cirripedia	95
7.7.	Oligocene Decapoda	95
7.8.	Oligocene Ophiuroidea	96
8.	Miocene types	96
8.1.	Miocene Foraminiferida	96
8.2.	Miocene Polyplacophora	98
8.3.	Miocene Bivalvia	98
8.4.	Miocene Gastropoda	104
8.5.	Miocene Annelida	110
8.6.	Miocene Brachiopoda	111
8.7.	Miocene Cirripedia	111
8.8.	Miocene Decapoda	112
8.9.	Miocene Crinoidea	120
8.10.	Miocene Echinoidea	120
9.	Pliocene types	121
9.1.	Pliocene Gastropoda	121
10.	Pleistocene types	121
10.1.	Pleistocene Gastropoda	121
Catalogue of vertebrate type specimens	123	
11.	Pisces	123
12.	Reptilia	127
13.	Aves	134
14.	Mammalia	135
Index of species and subspecies names	171	
References	195	

Introduction

Type specimens constitute the scientifically most valuable part of any paleontological collection. The International Code of Zoological Nomenclature (ICZN) (INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE 1999) clearly outlines the responsibilities of public collections with regards to primary and secondary type specimens. Every effort should be made to properly curate and store these specimens and make them available to the wider scientific community. The Department of Geology and Paleontology of the Hungarian Natural History Museum (HNHM) houses a large number of both invertebrate and vertebrate types. Until recently, no up-to-date and publicly accessible type catalogue existed and our in-house inventory did not facilitate easy listing and retrieval of type specimens. Both the publishing of a printed catalogue and developing and maintaining a web-based, searchable database are aimed at better serving the research needs of the Hungarian and international paleontological community. This work provides a comprehensive listing of our invertebrate and vertebrate paleontological types. It includes type specimens preserved in the collection as well as those types known or believed to be lost or missing which are attributed in the original publications to our museum or its ancestral mother institution, the Hungarian National Museum (HNM).

History and development of the type catalogue

The eventful history of the paleontological collections of the HNHM from the 19th century beginnings within the HNM through the late 20th century is summarized by KECSKEMÉTI & NAGY (1987). The collections of HNHM survived the World War II with relatively minor losses but during the Hungarian uprising in October 1956, a fire destroyed a major part of the collections and resulted in massive damage. Unfortunately this also led to the destruction of a large number of type specimens. The types believed to be permanently lost at this or other events are included in this catalogue with the appropriate remark. After 1956, a new inventory of the surviving collections was started. Meanwhile the remaining part of the collections was supplemented by generous donations from other institutions and new acquisitions from a vigorous paleontological research program. Further type specimens were obtained in both ways.

The first modern effort to build a catalogue of type specimens is the work of BODA (1964) that includes types described from Hungary in the paleontological literature up to 1960, housed in either of the two main type repositories of the country, the HNHM and the Geological Institute of Hungary (GIH). However, only a few years after the destruction at HNHM, it was not yet possible to

8 PALEONTOLOGICAL TYPE CATALOGUE

locate all specimens or confirm their loss, hence the listing is replete with types of unknown whereabouts. Moreover, the catalogue of BODA (1964) contains only species described from the territory of present-day Hungary, thereby disregarding a significant number of types from either those parts of historical Hungary which now belong to neighbour states, or countries farther afield.

No update of the catalogue of BODA (1964) was published in the following 40 years. Developing a card-catalogue of type specimens in the HNHM had been carried out for internal use by former staff member I. Z. NAGY but his project has not been completed. Our recent effort used that card catalogue as a starting point and supplemented it by data culled from recent literature. For the first time, a digital catalogue of the paleontological types was compiled and made available to the scientific community through a searchable database on the world-wide web in 2004. The website can be visited at http://www.nhmus.hu/kutatas/adatbazisok/paleotipus/index_a.html, and a catalogue search is also possible directly at <http://specify.nhmus.hu>.

Subsequently, the collections were moved to temporary premises in 2005 and, shortly afterwards, to the newly renovated home of the Department of Geology and Paleontology in 2006. Packing, moving, unpacking and rearranging of the entire collection necessitated a quick and rudimentary computerization of our holdings. During this process we discovered that the web-based type catalogue was far from complete and did not include a significant number of type specimens.

Furthermore, this and earlier type listings compiled for internal use often reveal two common mistakes. First, a figured specimen of a type series was commonly uncritically interpreted as holotype, even if multiple specimens of the type series existed and the author made no unambiguous holotype designation. Article 73 of ICZN states that fixation of the holotype must be explicit, thus illustrating a specimen does not in itself constitute a holotype designation. Second, specimens of an original type series other than the figured one were often left unlabelled as types, based on the erroneous assumption that paratypes have to be explicitly designated by the author. However, the same article states that all specimens of a type series other than the holotype are to be regarded paratypes (or syntypes if no holo- or lectotype has been selected), unless the author specifically excludes them.

For the first case, we have reinterpreted some holo- and paratypes listed in BODA 1964 as syntypes. For the second case, in the lack of a comprehensive digital inventory, likely we have not been able to trace all previously unlabelled types. However, effort has been made to recognize original type series and those specimens have now been added to the list as additional syntypes or paratypes.

Prior to publication of this catalogue we took several steps towards completeness: 1) we gleaned further data from the primary paleontological literature; 2) we included all types discovered when handling every specimen during

the move of our collections; 3) we reprocessed the catalogue of BODA (1964) in search of possible types existing in our collections but unknown at that time; 4) we culled the hand-written inventory books in search of possible hints for the type status of specimens; and 5) we critically reviewed the material donated after 1956 by the GIH and the Eötvös University of Budapest, as important historical collections that contain type specimens.

In the new facility of the HNHM, the type specimens are now stored separately within the paleontological collections, ensuring a higher level of curation. A disturbing number of type specimens has not been possible to locate, even though some of them are recorded in the post-1956 inventories. Clearly, the standard of curation at HNHM in the past 50 years has left much to be desired. The missing types may be accounted for by a lack of special attention to the type specimens and also the recent moves of the collection. It is hoped that at least some of the currently missing type specimens will be rediscovered during the planned complete revision and full digital cataloguing of our holdings.

Type specimens of 1181 fossil species-group taxa are listed in the catalogue, of them 986 are invertebrates and 195 are vertebrates. A total of 1131 taxa have or would have name-bearing primary types in our collection. Discounting the specimens that are missing, lost or have been transferred to other institutions, the catalogue includes primary types of 917 taxa available in the collection of HNHM. In total 2017 inventory lots are found to contain type specimens and recorded in the catalogue.

The paleontological literature, in which the invertebrate and vertebrate taxa represented by types originally or subsequently deposited in the HNHM collection were introduced, consists of more than 250 items listed in the References. The oldest work dates back to 1841 and the latest publication appeared in February 2008.

Historically, paleobotanical specimens, including types, form part of the collection of Department of Botany at HNHM. A comprehensive catalogue of more than 600 type specimens of fossil plant taxa described in the 19th century was published recently (HABLY et al. 2001). Some but not all of the paleobotanical types from 20th century works are treated in a previously published catalogue (HABLY & SZAKÁLY 1989).

Notes on structure and use of type catalogue

This catalogue includes primary, name-bearing types (i.e. holotypes, syntypes, lectotypes, neotypes) and secondary types (i.e. paratypes and paralectotypes) of fossil invertebrate and vertebrate species-group taxa currently known to exist in the collection of HNHM. Types explicitly stated in the original publication as deposited in the HNHM (or HNM) but not found in the present collection are also included, with the appropriate annotation. At present we have not attempted to list the tertiary types (i.e. all other figured and/or measured speci-

mens from published redescriptions of taxa, informally sometimes also called hypotypes or plesiotypes). Interpretation of type specimens follow the rules of the ICZN. Types for invalid names (i.e. junior homonyms) are not included.

The full catalogue will be also available in digital form as a searchable, web-based database that includes more information than listed in this print version. The data structure is presented in Table 1.

The printed catalogue is subdivided into separate parts for invertebrate and vertebrate types. The basic information of taxa and their type specimens is listed in slightly different formats for invertebrates and vertebrates. An individual catalogue entry typically corresponds to an inventory lot that contains the type specimen(s) of a taxon, identified by an inventory number. If a taxon is represented by more than one type specimen in our collection, separate entries are listed where differences exist in either the inventory number, locality, or illustration of specimen. Types under more than one inventory number are merged into a single entry if all data fields other than the inventory number are identical. Complexities often led to slight deviations from the strict application of these rules.

The format for invertebrate entries is as follows:

- Genus species* AUTHOR, year
- [or *Genus (Subgenus) species subspecies* AUTHOR, year]
- Status of type, inventory number
- [Period and/or epoch], stage, formation
- Locality, town, country
- Reference, page, text-figure, plate, figure
- Revised name, reference to revising work
- Remarks

The format for pre-Pliocene vertebrate entries is as follows:

- Genus species* AUTHOR, year
- [or *Genus (Subgenus) species subspecies* AUTHOR, year]
- Status of type, inventory number
- Part of fossil
- Stage, epoch; formation
- Locality, town, country
- Reference, page, text-figure, plate, figure
- Revised name, reference to revising work
- Remarks

The format for Pliocene and Quaternary vertebrate entries is as follows:

- Genus species* AUTHOR, year
- [or *Genus (Subgenus) species subspecies* AUTHOR, year]
- Status of type, inventory number
- Part of fossil
- Epoch subdivided (i.e. Early/Middle/Late Pliocene or Pleistocene)
- Locality, town, country
- Reference, page, text-figure, plate, figure
- Revised name, reference to revising work
- Remarks

Data category	Data fields in printed and digital catalogue
Name & author	GENUS SUBGENUS SPECIES SPECIES AUTHOR¹ SUBSPECIES SUBSPECIES AUTHOR YEAR
Specimen information	STATUS OF TYPE INVENTORY NUMBER PART OF FOSSIL² Collector Year of collection
Taxonomic position	PHYLUM CLASS Order Family
Age & formation	PERIOD EPOCH STAGE, STANDARD³ STAGE, REGIONAL³ Biozone FORMATION Bed
Locality	LOCALITY TOWN COUNTRY⁴
Reference	REFERENCE⁵ PAGE TEXT-FIGURE PLATE FIGURE
Revision	REVISED NAME Reviser REFERENCE OF REVISION⁵
Remarks	REMARKS <i>Data enterer</i>
Storage	<i>New cabinet/drawer</i> <i>Old cabinet</i> <i>Old drawer</i>

Table 1 – Data structure of the type specimen database used to produce listings for this catalogue

Uppercase, bold letters: data fields listed in printed catalogue; uppercase, normal letters: data fields used for grouping of entries in printed catalogue; lowercase, normal letters: data fields available in the web-based, searchable database; italics: data for internal use only.

¹ Species author not listed if type is for subspecies.

² Only appears in printed catalogue for vertebrate types.

³ Either standard or regional stages used for different geological periods and epochs.

⁴ Country is listed only if other than Hungary.

⁵ See code numbers in References section.

The invertebrate catalogue entries are arranged primarily by geologic age, secondarily by major taxonomic groups, then the originally proposed bi- or trinomens are listed alphabetically within each subdivision.

Both the ages and the taxonomic groups that define the subdivisions were selected for practical reasons rather than strictly using certain levels in their hierarchical structure. The following age subdivisions, ranging in rank from era to epoch, are used: Paleozoic, Triassic, Jurassic, Cretaceous, Paleocene, Eocene, Oligocene, Miocene, Pliocene and Pleistocene.

Secondary, taxonomic subdivisions use the following, well-known groups, arbitrarily taken from higher taxa at the phylum, class, or order level: Radiolaria, Foraminiferida, Anthozoa, Polyplacophora, Bivalvia, Scaphopoda, Gastropoda, Cephalopoda, Annelida, Brachiopoda, Trilobita, Ostracoda, Cirripedia, Decapoda, Crinoidea, Ophiuroidea, Echinoidea. Classification of fossil taxa follows that used in the Paleobiology Database (www.paleodb.org) as of April 2008. The sequence of higher taxa reflects that used in the Tree of Life Web Project (www.tolweb.org). The catalogue of vertebrate types is arranged by higher taxa, containing subdivisions for Pisces, Reptilia, Aves, and Mammalia. The obsolete category of Pisces is used here for historical reasons to include taxa from Chondrichthyes and Actinopterygii.

Taxon names for the types are given as bi- or trinomens as they were introduced by the author in the original description. The original spelling is retained, even though current rules of ICZN (see Article 32) require to avoid the diacritical marks of non-standard Latin characters (commonly occurring in taxon names derived from Hungarian person or place names). However, preserving the original spelling in the catalogue may help tracing a taxon in the literature. Hyphenated species epithets are also retained here. Necessary corrections and emendations of originally incorrect spelling of species names are reflected in the index of species and subspecies names. Capitalization of species names, common in the old literature for names derived from proper names, has been removed throughout.

If types of a taxon require more than one entry, the holotype or lectotype is listed first. For types of equal status (e.g. syntypes, paratypes, paralectotypes), the entries are arranged in ascending order of their inventory number. In merged entries with several specimens of identical attributes, the inventory numbers are given in a list separated by commas.

If possible, an age assignment for invertebrate specimens is given at the stage level. Standard stages are used (GRADSTEIN et al. 2004) except for the Oligocene and Miocene, where the regional stages of the central Paratethys time scale are applied (PILLER et al. 2007). Although some stages are known to straddle standard epoch boundaries, for sake of a simplified hierarchy of age units, these boundary problems are sidestepped. Thus the Egerian is included in the Oligocene, and the Pannonian is taken to form part of the Miocene. Original age

assignments are revised if necessary, where up-to-date stratigraphic information is available from the type locality.

The ages of Pliocene and Pleistocene vertebrates is given using a threefold (i.e. Early–Middle–Late) subdivision of the epochs, derived by simplifying the biochronological framework of KRETZOI (1969) and JÁNOSSY (1986), with the necessary modifications to account for different concepts in placing the Pliocene/Pleistocene boundary. Age assignment of pre-Pleistocene vertebrates follows the concepts applied to invertebrates.

The lithostratigraphic unit from which a type specimen was collected is given on the basis of CSÁSZÁR et al. (1997) for Hungary and FILIPESCU (2001) for the Transylvanian part of Romania. As modern formation names were not in use prior to the 1980's, most of the formation assignments are made here using the available information on lithology, age and locality. Pliocene and younger vertebrates are not assigned to formations as they are most commonly found in fissure fills and cave deposits, not forming conventional lithostratigraphic units.

In the locality field, geographic names are given in their usual Hungarian form, for ease of finding them on local maps. The English translation of some common vernacular terms in geographic names (often hyphenated to form the latter part of a proper name) is listed as follows (Hungarian words in italics): *árok* – gully; *barlang* – cave; *dombok* – hills; *fennsík* – plateau; *hegy* – mountain; *kő* – ~cliff; *major* – farm; *mező* – meadow; *patak* – creek; *puszta* – farm; *szakadék* – gorge; *szőlők* – vineyard; *szurdok* – gorge; *tér (tere)* – square; *tető* – summit; *tó* – lake; *út* – road; *utca* – street; *völgy* – valley. Geological terms for localities, such as outcrop ("feltárás"), quarry ("köfejtő"), brickyard ("téglagyár"), are given in English. Name of town refers to the municipality to which the locality belongs, in some cases this is not the nearest settlement.

As the majority of the types were collected at localities within the present-day borders of Hungary, no country name is listed for them. Some specimens are from sites in historical Hungary that are now in the territory of Austria, Croatia, Romania, Serbia or Slovakia. Their town is listed using the current official name, with the Hungarian name given in parentheses. A lesser number of types are from other countries, including Albania, Algeria, Argentina, Bulgaria, China, Czech Republic, France, Germany, Italy, Peru, Russia, Serbia, Turkey, Ukraine, and the USA.

Reference is given to the original description, using a five or six character code, found in the reference list at the end of each item. The reference code is composed of the first three letters of the first author's name (or an otherwise meaningful three-letter abbreviation of the name), the last two digits of the year of publication, and, if needed, an extra letter to distinguish works from the same year. If the original description appeared in Hungarian as well as in a foreign language, both page numbers are given, the foreign one first and the Hungarian one in parentheses.

Illustrations (text-figures and/or figures on plates) accompanying the original description are quoted at the appropriate entry, if the specimen is identified as the figured one. Unmatched illustrations from the original description of a taxon are listed in parentheses in the first catalogue entry of that taxon.

If we became aware of a published revision, either nomenclatural, or changing the species-level identification or the generic assignment of the taxon concerned, we include the revised name and provide the reference code of the revising work. In some cases both the first occurrence of a revised combination and the most recent redescription of the revised taxon are quoted. However, it was beyond the scope of this project to research the validity or nomenclatural status of each taxa.

Any additional information may appear as remarks in the last line of an item. Remarks may include reference to a different inventory number in the original publication, notes on the type designation, or the availability of the specimen. Types known with certainty to be vanished are classified as lost. A special remark applies to many specimens known to have been destroyed in 1956. The probability of loss may also be indicated. Types may be reported missing either with or without a current inventory number. In the first case the specimen was definitely in the collection after 1956 and has a higher chance of rediscovery.

To facilitate finding information on taxa, the catalogue is followed by an index of all invertebrate and vertebrate species and subspecies names. Here an original name (using emended spelling if needed) is followed by the originally published bi- or trinomen, the author's name, and the page number on which the first catalogue entry of this taxon appears.

Although significant care was taken to compile a comprehensive catalogue that contains accurate information, errors and omissions are almost inevitable in an endeavour of this scale. Therefore users of this catalogue are encouraged to report any suggested correction and/or addition to the authors or the Department of Geology and Paleontology.

Acknowledgements

Special thanks to Zsuzsanna MOLNÁR for the tireless search for types in the collections, inventory books and literature, for her dedicated work in data entry, editing and layout, and for a relentless strive for accuracy. Our fellow curators, István FÓZY, Tibor KECSKEMÉTI, János SZABÓ and Attila VÖRÖS provided indispensable help in the collection under their care. Mariann BOSNAKOFF assisted in assigning type horizons to lithostratigraphic formations and in library searches. Attila ŐSI is thanked for his support and valuable expertise in vertebrate anatomy. The unpublished card catalogue of our retired colleague István Zoltán NAGY was used as a starting point in our compilation. Eszter HANKÓ

and Katalin BABÁK contributed to the data entry in the early stage of the project. László KORDOS (GIH) answered many queries about specimens donated to the HNHM by the GIH. We are grateful to many colleagues who generously gave advise, helped with curatorial matters or provided access to literature: Stanislav ČERMÁK (Prague), Rodica CIOBANU (Sibiu), Zoltán CSIKI (Bucharest), Sorin FILIPESCU (Cluj), Łucja FOSTOWICZ-FRELIK (Warsaw), János HÍR (Pásztó), Wolfgang KIESSLING (Berlin), László KOCSIS (Lausanne), Jürgen KRIWET (Berlin), Éva LENGYEL (HNHM), Imre MAGYAR (Budapest), Oleg MANDIC (Vienna), David MAYHEW (Rotterdam), Pál MÜLLER (GIH), Christian NEUMANN (Berlin), Olga PIROS (GIH), István SZENTE (Eötvös University of Budapest), György TOPÁL (HNHM), István VÖRÖS (HNM), Jan WAGNER (Prague) and Ferenc WANEK (Cluj). The virtual community of the PaleoNet mailing list contributed to interpretation of some complex issues. András GALÁCZ critically read an earlier version of the invertebrate catalogue. Photos on the front and back cover are by Attila ŐSI and István FÓZY, respectively. Development of a web-based type catalogue was supported through a grant from the Ministry of National Cultural Heritage in 2004. Publication of the print catalogue was made possible through a grant from the National Cultural Fund.

Catalogue of invertebrate type specimens

1. Paleozoic types

1.1. Paleozoic Foraminiferida

Fusulinella lóczyi LÖRENTHEY, 1897
syntypes
Middle Carboniferous, "fusulinid limestone"
Teng-tian-ching, Kansu Province, China
Lör97, p. 228, text-figs. 22–23
Translated description in German edition,
LÖRENTHEY 1899: p. 269. Types probably lost

Lingulina nankingensis LÖRENTHEY, 1897
syntypes
Carboniferous, "limestone"
Nanking, China
Lör97, p. 239, text-fig. 35
Translated description in German edition,
LÖRENTHEY 1899: p. 282. Types probably lost

Lingulina nankingensis LÖRENTHEY, 1897
syntypes
Carboniferous or Permian, "limestone"
pass between Kien-chuen-chou and Niu-ke,
Yunman Province, China
Lör97, p. 239, text-fig. 35
Types probably lost

Lingulina széchenyi LÖRENTHEY, 1897
syntypes
Carboniferous, "marly limestone"
Nanking, China
Lör97, p. 237, text-figs. 33–34
Translated description in German edition,
LÖRENTHEY 1899: p. 280. Types probably lost

Lingulina széchenyi LÖRENTHEY, 1897
syntypes
Carboniferous or Permian, "fusulinid
limestone"
Santa-sien, Kansu Province, China
Lör97, p. 237, text-figs. 33–34
Types probably lost

Lingulina széchenyi LÖRENTHEY, 1897
syntypes
Carboniferous or Permian, "limestone"
pass between Kien-chuen-chou and Niu-ke,
Yunnan Province, China
Lör97, p. 237, text-figs. 33–34
Types probably lost

Lingulina széchenyi LÖRENTHEY, 1897
syntypes
Carboniferous, "limestone"
Sining-fu, Kansu Province, China
Lör97, p. 237, text-figs. 33–34
Types probably lost

Lingulina széchenyi LÖRENTHEY, 1897
syntypes
Carboniferous, "crinoidal limestone"
Sa-men-kwan saddle, Ta-tia-san Mountains,
Kansu Province, China
Lör97, p. 237, text-figs. 33–34
Types probably lost

Nodosinella simplex LÖRENTHEY, 1897
holotype
Middle Carboniferous, "limestone"
Teng-tian-ching, Kansu Province, China
Lör97, p. 236, text-fig. 32
Translated description in German edition,
LÖRENTHEY 1899: p. 279. Type probably lost

Spirillina chinensis LÖRENTHEY, 1897
syntypes
Carboniferous or Permian, "limestone"
Yung-chang-fu, Yunnan Province, China
Lör97, p. 234, text-figs. 26–29
Translated description in German edition,
LÖRENTHEY 1899: p. 276. Types probably lost

Spirillina plana patella LÖRENTHEY, 1897
syntypes
Carboniferous, "crinoidal limestone"
Sa-men-kwan saddle, Ta-tia-san Mountains,
Kansu Province, China
Lör97, p. 231, text-figs. 24–25
Translated description in German edition,
LÖRENTHEY 1899: p. 272. Types probably lost

Spirillina plana patella LÖRENTHEY, 1897
 syntype
 Carboniferous or Permian, "limestone"
 Yung-chang-fu, Yunnan Province, China
 Lör97, p. 231, text-figs. 24–25
 Type probably lost

Lonsdaleoides bükkicense KOLOSVÁRY, 1951
 syntype, M.57.2902
 Carboniferous or Permian, Mályinka
 Formation
 Mályinka
 Kol51a, p. 182 (29), pl. 4, figs. 3–5

1.2. Paleozoic Anthozoa

Cyathophyllum lóczyi FRECH, 1897
 holotype
 Middle Devonian, "grey limestone"
 Hoaling-pu, Sichuan Province, China
 Fre97, p. 195, pl. 9, figs. 1–4
 Translated description in German edition,
 FRECH 1899: p. 231. Type probably lost

Lonsdaleoides bükkicense KOLOSVÁRY, 1951
 syntype, M.57.2925
 Carboniferous or Permian, Mályinka
 Formation
 Dezső-völgy, Dédes
 Kol51a, p. 182 (29), pl. 4, figs. 3–5

Dibunophyllum kissi KOLOSVÁRY, 1951
 syntype, M.57.3000
 Visean, Early Carboniferous, Szabadbattyán
 Limestone Formation
 Szár-hegy, Szabadbattyán
 Kol51b, p. 282 (280), pl. 9, figs. 14–18, pl. 11,
 figs. 28–29, pl. 12, figs. 31, 34–36

Phineus confluentiseptatus KOLOSVÁRY, 1951
 syntype, M.57.2901
 Carboniferous or Permian, Mályinka
 Formation
 Felsőszólóköve, Mályinka
 Kol51a, p. 185 (172), pl. 16, figs. 7–12

Favosites asteriscus FRECH, 1897
 syntypes
 Middle Devonian, "marly limestone"
 Hoaling-pu, Sichuan Province, China
 Fre97, p. 196, pl. 8, figs. 3, 3a–c
 Translated description in German edition,
 FRECH 1899: p. 232. Types probably lost

Phineus confluentiseptatus KOLOSVÁRY, 1951
 syntype, M.57.2933
 Carboniferous or Permian, Mályinka
 Formation
 near Szelecsikő, Dédes
 Kol51a, p. 185 (172), pl. 16, figs. 7–12

Favosites goldfussi major FRECH, 1897
 holotype
 Middle Devonian
 Hoaling-pu, Sichuan Province, China
 Fre97, p. 196, pl. 8, figs. 1, 1a–e
 Translated description in German edition,
 FRECH 1899: p. 232. Type probably lost

Pleramplexus vadászi KOLOSVÁRY, 1951
 holotype, M.57.2992
 Late Permian, Nagyvisnyó Limestone
 Formation
 Felsőszólóköve, Mályinka
 Kol51a, p. 185 (50), pl. 14, figs. 13–14

Haplothecia? chinensis FRECH, 1897
 holotype
 Middle Devonian
 Hoaling-pu, Sichuan Province, China
 Fre97, p. 195, pl. 9, figs. 5, 5a
 Translated description in German edition,
 FRECH 1899: p. 231. Type probably lost

Plerophyllum (Ufimia) baloghi KOLOSVÁRY, 1951
 holotype, M.57.2938
 Late Carboniferous, Mályinka Formation
 railway cut No. 1, Nagyvisnyó
 Kol51a, p. 184 (48), pl. 13, figs. 9–10

Plerophyllum (Ufimia) longicontraseptatum
 KOLOSVÁRY, 1951
 syntypes, M.57.2937
 Late Carboniferous, Mályinka Formation
 railway cut No. 1, Nagyvisnyó
 Kol51a, p. 184 (49), pl. 13, fig. 8

Plerophyllum (Ufimia) rakuszi KOLOSVÁRY,

1951

holotype

Late Carboniferous, Mályinka Formation

railway cut No. 1, Nagyvisnyó

Kol51a, p. 185 (49), pl. 14, fig. 8

Type probably destroyed in 1956

Polycoelia malyinkae KOLOSVÁRY, 1951

holotype

Carboniferous or Permian, Mályinka

Formation

Felsőszőlőköve, Mályinka

Kol51a, p. 40, pl. 11, figs. 4–6

Type probably destroyed in 1956

Polycoelia profundiformis KOLOSVÁRY, 1951

holotype, M.57.2900

Carboniferous or Permian, Mályinka

Formation

Felsőszőlőköve, Mályinka

Kol51a, p. 41, pl. 11, figs. 7–8

Prosmilia helenae KOLOSVÁRY, 1951

holotype, M.57.2998

Late Permian, Nagyvisnyó Limestone

Formation

Felsőszőlőköve, Mályinka

Kol51a, p. 183 (44), pl. 18, fig. 6, pl. 19,

figs. 1–2

Sinophyllum gracile KOLOSVÁRY, 1951

syntypes

Late Carboniferous, Mályinka Formation

railway cut No. 1, Nagyvisnyó

Kol51a, p. 173, pl. 16, figs. 5–6

Types probably destroyed in 1956

1.3. Paleozoic Bivalvia

Actinopteria? densiradiata LÓCZY, 1897

holotype

Devonian, "dark, marly dolomitic limestone"

Hoaling-pu, Sichuan Province, China

Lóc97, p. 23, pl. 7, fig. 1

Translated description in German edition,

LÓCZY 1899: p. 27. Type probably lost

1.4. Paleozoic Gastropoda

Bellerophon (Bucania?) incerta LÓCZY, 1897

syntypes

Carboniferous, "grey, bituminous marly limestone"

Teng-tian-ching, Kansu Province, China

Lóc97, p. 40, pl. 1, figs. 12, 13

Translated description in German edition, LÓCZY 1899: p. 47. Types probably lost

Loxonema széchenyi LÓCZY, 1897

holotype

Carboniferous, "grey, bituminous marly limestone"

Teng-tian-ching, Kansu Province, China

Lóc97, p. 43, pl. 1, figs. 8, 9

Translated description in German edition, LÓCZY 1899: p. 51. Type probably lost

Macrochilina kreitneri LÓCZY, 1897

holotype

Carboniferous, "grey, bituminous marly limestone"

Teng-tian-ching, Kansu Province, China

Lóc97, p. 45, pl. 1, fig. 10

Translated description in German edition, LÓCZY 1899: p. 52. Type probably lost

1.5. Paleozoic Cephalopoda

?*Nautilus kayseri* LÓCZY, 1897

holotype

Carboniferous, "grey, bituminous marly limestone"

Teng-tian-ching, Kansu Province, China

Lóc97, p. 38, pl. 1, figs. 7, 7a

Translated description in German edition, LÓCZY 1899: p. 44. Type probably lost

Nautilus (Temnocheilus) waageni LÓCZY, 1897

holotype

Carboniferous, "grey, bituminous marly limestone"

Teng-tian-ching, Kansu Province, China

Lóc97, p. 38, pl. 1, figs. 6, 6a–b

Translated description in German edition, LÓCZY 1899: p. 45. Type probably lost

1.6. Paleozoic Brachiopoda

Camarophoria szü-tschan-ensis LÓCZY, 1897
holotype
Devonian, "dark, marly dolomitic limestone"
Hoaling-pu, Sichuan Province, China
Lóc97, p. 31, pl. 7, fig. 21
Translated description in German edition,
LÓCZY 1899: p. 36. Type probably lost

Chonetella dubia LÓCZY, 1897
holotype
Carboniferous, "black clayey marl"
Teng-tian-ching, Kansu Province, China
Lóc97, p. 69, pl. 3, fig. 15
Translated description in German edition,
LÓCZY 1899: p. 81. Type probably lost

Chonetes flemmingi gobica LÓCZY, 1897
syntypes
Carboniferous, "black clayey marl"
Teng-tian-ching, Kansu Province, China
Lóc97, p. 65, pl. 3, fig. 14
Translated description in German edition,
LÓCZY 1899: p. 77. Types probably lost

Chonetes orientalis LÓCZY, 1897
syntypes
Devonian, "dark, marly dolomitic limestone"
Hoaling-pu, Sichuan Province, China
Lóc97, p. 24, pl. 7, figs. 12–17
Translated description in German edition,
LÓCZY 1899: p. 28. Types probably lost

Productus (Marginifera) desgodinsi LÓCZY, 1897
holotype
Carboniferous or Permian, "marly limestone
and black shale"
Yarkalo, Lan-tsang-kiang Valley, China
Lóc97, p. 90, pl. 4, fig. 6
Translated description in German edition,
LÓCZY 1899: p. 106. Type probably lost

Productus yunnanensis LÓCZY, 1897
syntypes
Carboniferous or Permian, "bituminous
limestone"
Vo-sih-vo cave, Yung-chang-fu, Yunnan
Province, China
Lóc97, p. 106, pl. 6, figs. 1, 2
Translated description in German edition,
LÓCZY 1899: p. 125. Types probably lost

Reticularia waageni LÓCZY, 1897
syntypes
Carboniferous or Permian, "marly limestone
and black shale"
Yarkalo, Lan-tsang-kiang Valley, China
Lóc97, p. 93, pl. 4, figs. 1, 2
Translated description in German edition,
LÓCZY 1899: p. 110. Types probably lost

1.7. Paleozoic Trilobita

Phillipsia kansuensis LÓCZY, 1897
syntypes
Carboniferous, "grey, bituminous marly
limestone"
Teng-tian-ching, Kansu Province, China
Lóc97, p. 36, pl. 1, figs. 1–5
Translated description in German edition,
LÓCZY 1899: p. 42. Types probably lost

2. Triassic types

2.1. Triassic Radiolaria

Alatipicapora latoalata KOZUR, MOIX & OZSVÁRT,
2007
holotype, M 2008.89.1.
Carnian, Killik Formation
southwest of Tavşuşçayır Tepe, Turkey
Koz07c, p. 200, text-fig. 1c
Authors' inventory number: 23-9-04/III-95

Alatipicapora spinosa KOZUR, MOIX & OZSVÁRT,
2007
holotype, M 2008.88.1.
Carnian, Killik Formation
southwest of Tavşuşçayır Tepe, Turkey
Koz07c, p. 199, text-fig. 1b
Authors' inventory number: 23-9-04/II-75

Alatipicapora spinosa KOZUR, MOIX & OZSVÁRT,
2007
paratype, M 2008.88.2.
Carnian, Killik Formation
southwest of Tavşuşçayır Tepe, Turkey
Koz07c, p. 199, text-fig. 1a
Authors' inventory number: 23-9-04/VI-139

Alatipicapora tetrapedis KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.87.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 198, pl. 1, fig. 16
Authors' inventory number: 23-9-04/V-202

Archaeoacanthocircus angustiannulatus KOZUR,
MOIX & OZSVÁRT, 2007
holotype, M 2008.68.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 178, pl. 1, fig. 5
Authors' inventory number: 23-9-04/V-142

Archaeoacanthocircus angustiannulatus KOZUR,
MOIX & OZSVÁRT, 2007
paratype, M 2008.68.2.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 178, pl. 1, fig. 6
Authors' inventory number: 23-9-04/V-145

Archaeoacanthocircus latiannulatus KOZUR,
MOIX & OZSVÁRT, 2007
holotype, M 2008.67.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 178, pl. 1, fig. 1
Authors' inventory number: 23-9-04/VI-64

Archaeoacanthocircus latiannulatus KOZUR,
MOIX & OZSVÁRT, 2007
paratype, M 2008.67.2.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 178, pl. 1, fig. 2
Authors' inventory number: 23-9-04/III-98

Archaeoacanthocircus ovalis KOZUR, MOIX &
OZSVÁRT, 2007
holotype, M 2008.69.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 179, pl. 1, fig. 8
Authors' inventory number: 23-9-04/V-134

Archaeoacanthocircus rectangularis KOZUR,
MOIX & OZSVÁRT, 2007
holotype, M 2008.70.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 179, pl. 1, fig. 3
Authors' inventory number: 23-9-04/VI-44

Archaeoacanthocircus rectangularis KOZUR,
MOIX & OZSVÁRT, 2007
paratype, M 2008.70.2.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 179, pl. 1, fig. 4
Authors' inventory number: 23-9-04/VI-42

Archaeoacanthocircus transitus KOZUR, MOIX &
OZSVÁRT, 2007
holotype, M 2008.71.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 180, pl. 1, fig. 7
Authors' inventory number: 8-12-03/I-27

Huglusphaera aperta KOZUR, MOIX & OZSVÁRT,
2007
holotype, M 2008.73.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 182, pl. 1, fig. 13
Authors' inventory number: 23-9-04/I-22

Huglusphaera aspinosa KOZUR, MOIX & OZSVÁRT,
2007
holotype, M 2008.74.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 183, text-fig. 1b
Authors' inventory number: 8-12-03/I-32

Huglusphaera weemsi KOZUR, MOIX & OZSVÁRT,
2007
holotype, M 2008.72.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07b, p. 181, text-fig. 1a
Authors' inventory number: 23-9-04/V-133

22 TRIASSIC INVERTEBRATE TYPES

Huglusphaera weemsi KOZUR, MOIX & OZSVÁRT, 2007
paratype, M 2008.72.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 181, text-fig. 1c
Authors' inventory number: 23-9-04/V-148

Huglusphaera yini KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.75.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 183, pl. 1, fig. 10
Authors' inventory number: 23-9-04/V-138

Huglusphaera yini KOZUR, MOIX & OZSVÁRT, 2007
paratype, M 2008.75.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 183, pl. 1, fig. 12
Authors' inventory number: 23-9-04/V-143

Huglusphaera zakharovi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.76.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 183, pl. 1, fig. 11
Authors' inventory number: 23-9-04/VI-70

Huglusphaera zakharovi KOZUR, MOIX & OZSVÁRT, 2007
paratype, M 2008.76.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 183, pl. 1, fig. 9
Authors' inventory number: 23-9-04/V-144

Hypoxiphothecaella claviformis KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.63.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 166, pl. 2, fig. 11
Authors' inventory number: 23-9-04/II-77

Hypoxiphothecaella elongata KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.64.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 167, pl. 2, fig. 13
Authors' inventory number: 23-9-04/II-78

Hypoxiphothecaella foeyi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.65.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 167, pl. 2, fig. 12
Authors' inventory number: 23-9-04/II-87

Hypoxiphothecaella inflata KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.66.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 168, pl. 1, fig. 11
Authors' inventory number: 23-9-04/II-98

Hypoxiphothecaella mersinensis KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.62.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 166, pl. 2, fig. 10
Authors' inventory number: 23-9-04/II-86

Kahlerosphaera abnormis KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.77.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 185, pl. 2, fig. 1
Authors' inventory number: 23-9-04/II-174

Kahlerosphaera faludyi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.78.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07b, p. 185, pl. 2, fig. 3
Authors' inventory number: 23-9-04/VI-102

Kahlerosphaera hamvasi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.79.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 186, pl. 2, fig. 2
Authors' inventory number: 23-9-04/I-76

Kahlerosphaera karinthyi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.80.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 186, pl. 2, fig. 4
Authors' inventory number: 8-12-03/I-18

Kahlerosphaera kerteszi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.81.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 186, pl. 2, fig. 5
Authors' inventory number: 23-9-04/IV-105

Kahlerosphaera koestleri KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.82.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 187, pl. 2, fig. 6
Authors' inventory number: 23-9-04/I-125

Kahlerosphaera pamuki KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.83.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 187, pl. 2, fig. 7
Authors' inventory number: 23-9-04/III-6

Kahlerosphaera rejtoei KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.84.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 188, pl. 2, fig. 8
Authors' inventory number: 23-9-04/I-58

Kahlerosphaera szerbi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.85.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 188, pl. 2, fig. 9
Authors' inventory number: 23-9-04/I-36

Kahlerosphaera vonneguti KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.86.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07b, p. 189, pl. 2, fig. 10
Authors' inventory number: 23-9-04/III-49

Karnospongella multispinosa KOZUR, MOIX & MOSTLER, 2007
holotype, M 2008.107.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Moi07, p. 295, pl. 2, fig. 11
Authors' inventory number: 23-9-04/VI-33

Karnospongella multispinosa KOZUR, MOIX & MOSTLER, 2007
paratype, M 2008.107.2.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Moi07, p. 295, pl. 4, fig. 1
Authors' inventory number: 8-12-03/I-15

Pararuesticyrtium longispinosum KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.47.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07a, p. 154, pl. 1, fig. 1
Authors' inventory number: 23-9-04/III-79

Pararuesticyrtium mersinensis KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.48.1.
Carnian, Killik Formation
southwest of Tavusçayı Tepe, Turkey
Koz07a, p. 154, pl. 1, fig. 2
Authors' inventory number: 23-9-04/V-56

24 TRIASSIC INVERTEBRATE TYPES

Podobursa claviformis KOZUR, MOIX & OZSVÁRT,

2007

holotype, M 2008.93.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 205, pl. 1, fig. 1

Authors' inventory number: 23-9-04/I-134

Podobursa sceptrumides KOZUR, MOIX &

OZSVÁRT, 2007

holotype, M 2008.98.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 208, pl. 1, fig. 5

Authors' inventory number: 27-11-04/I-17

Podobursa fusiformis KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.94.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 206, pl. 1, fig. 2

Authors' inventory number: 23-9-04/II-97

Podobursa tuvalica KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.99.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 209, pl. 1, fig. 4

Authors' inventory number: 27-11-04/I-8

Podobursa fusiformis KOZUR, MOIX & OZSVÁRT, 2007

paratype, M 2008.94.2.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 206, pl. 1, fig. 3

Authors' inventory number: 23-9-04/II-66

Ruesticyrtium coronatum KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.49.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07a, p. 155, pl. 1, fig. 3

Authors' inventory number: 23-9-04/I-141

Podobursa longiceras longiceras KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.95.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 207, pl. 1, fig. 8

Authors' inventory number: 23-9-04/IV-208

Ruesticyrtium georgi georgi KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.50.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07a, p. 156, pl. 1, fig. 5

Authors' inventory number: 23-9-04/I-129

Podobursa longiceras tetraspinosa KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.96.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 207, pl. 1, fig. 9

Authors' inventory number: 23-9-04/V-140

Ruesticyrtium georgi georgi KOZUR, MOIX & OZSVÁRT, 2007

paratype, M 2008.50.2.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07a, p. 156, pl. 1, fig. 8

Authors' inventory number: 23-9-04/VI-169

Podobursa mersinensis KOZUR, MOIX & OZSVÁRT, 2007

holotype, M 2008.97.1.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07c, p. 207, pl. 1, fig. 6

Authors' inventory number: 23-9-04/II-90

Ruesticyrtium georgi georgi KOZUR, MOIX & OZSVÁRT, 2007

paratype, M 2008.50.3.

Carnian, Killik Formation

southwest of Tavuşçayı Tepe, Turkey

Koz07a, p. 156, pl. 1, fig. 9

Authors' inventory number: 23-9-04/III-113

- Ruesticyrtium georgi novemdentatum* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.51.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07a, p. 157, pl. 1, fig. 4
Authors' inventory number: 23-9-04/V-58
- Ruesticyrtium latidentatum* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.52.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07a, p. 157, pl. 1, fig. 7
Authors' inventory number: 23-9-04/VI-140
- Ruesticyrtium lobatum* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.54.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07a, p. 158, pl. 1, fig. 10
Authors' inventory number: 23-9-04/I-140
- Ruesticyrtium mostleri* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.53.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07a, p. 158, pl. 1, fig. 12
Authors' inventory number: 23-9-04/III-153
- Spinomersinella goricanae* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.90.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 201, pl. 1, fig. 14
Authors' inventory number: 23-9-04/I-64
- Spinomersinella multispinosa* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.91.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 202, pl. 1, fig. 15
Authors' inventory number: 23-9-04/II-84
- Spinoprotunuma? constricta* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.102.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 212, pl. 1, fig. 11
Authors' inventory number: 23-9-04/VII-15
- Spinoprotunuma? furcastostriata* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.103.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 212, pl. 1, fig. 12
Authors' inventory number: 27-11-04/II-143
- Spinoprotunuma triassica* KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.101.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Koz07c, p. 211, pl. 1, fig. 10
Authors' inventory number: 23-9-04/V-250
- Spongotorilispinus inaequispinosus* KOZUR, MOIX & MOSTLER, 2007
holotype, M 2008.109.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Moi07, p. 296, pl. 3, fig. 5
Authors' inventory number: 23-9-04/II-46
- Spongotorilispinus moixi* KOZUR & MOSTLER, 2007
holotype, M 2008.108.1.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Moi07, p. 295, pl. 2, fig. 10
Authors' inventory number: 23-9-04/I-188
- Spongotorilispinus moixi* KOZUR & MOSTLER, 2007
paratype, M 2008.108.2.
Carnian, Killik Formation
southwest of Tavuşçayı Tepe, Turkey
Moi07, p. 295, pl. 2, fig. 9
Authors' inventory number: 23-9-04/II-6

Spongotorilispinus moixi KOZUR & MOSTLER, 2007

paratype, M 2008.108.3.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 295, pl. 3, fig. 1

Authors' inventory number: 23-9-04/II-7

Spongotorilispinus moixi KOZUR & MOSTLER, 2007

paratype, M 2008.108.4.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 295, pl. 3, fig. 2

Authors' inventory number: 8-12-03/I-25H

Spongotorilispinus moixi KOZUR & MOSTLER, 2007

paratype, M 2008.108.5.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 295, pl. 3, fig. 3

Authors' inventory number: 8-12-03/I-59

Spongotorilispinus ozsvarti KOZUR, MOIX &

MOSTLER, 2007

holotype, M 2008.110.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 296, pl. 3, fig. 4

Authors' inventory number: 8-12-03/I-13

Spongotorilispinus turkensis KOZUR, MOIX & MOSTLER, 2007

holotype, M 2008.111.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 296, pl. 3, fig. 7

Authors' inventory number: 27-11-04/II-101

Spongotorilispinus turkensis KOZUR, MOIX & MOSTLER, 2007

paratype, M 2008.111.2.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 296, pl. 3, fig. 6

Authors' inventory number: 23-9-04/II-1

Spongotorilispinus tuvalicus MOIX & MOSTLER,

2007

holotype, M 2008.112.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Moi07, p. 296, pl. 3, fig. 9

Authors' inventory number: 23-9-04/II-37

Stampfliella tuvalica KOZUR, MOIX & OZSVÁRT,

2007

holotype, M 2008.92.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Koz07c, p. 204, pl. 1, fig. 13

Authors' inventory number: 23-9-04/III-166

Syringocapsa glabra KOZUR, MOIX & OZSVÁRT,

2007

holotype, M 2008.100.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Koz07c, p. 209, pl. 1, fig. 7

Authors' inventory number: 23-9-04/VII-3

Velouesticyrtium palfyi KOZUR, MOIX &

OZSVÁRT, 2007

holotype, M 2008.55.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Koz07a, p. 159, pl. 1, fig. 6

Authors' inventory number: 23-9-04/I-49

Xiphothecaella brevicaudata KOZUR, MOIX &

OZSVÁRT, 2007

holotype, M 2008.56.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Koz07a, p. 161, pl. 2, fig. 1

Authors' inventory number: 23-9-04/II-117

Xiphothecaella elegans KOZUR, MOIX & OZSVÁRT,

2007

holotype, M 2008.57.1.

Carnian, Killik Formation

southwest of Tavşusçayırı Tepe, Turkey

Koz07a, p. 162, pl. 2, fig. 2

Authors' inventory number: 23-9-04/II-94

Xiphothecaella longicaudata KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.58.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 162, pl. 2, fig. 3
Authors' inventory number: 23-9-04/V-158

Xiphothecaella longicaudata KOZUR, MOIX & OZSVÁRT, 2007
paratype, M 2008.58.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 162, pl. 2, fig. 4
Authors' inventory number: 23-9-04/II-80

Xiphothecaella orchardi KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.59.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 163, pl. 2, fig. 5
Authors' inventory number: 23-9-04/II-93

Xiphothecaella procera KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.60.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 164, pl. 2, fig. 7
Authors' inventory number: 23-9-04/II-68

Xiphothecaella procera KOZUR, MOIX & OZSVÁRT, 2007
paratype, M 2008.60.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 164, pl. 2, fig. 6
Authors' inventory number: 23-9-04/II-92

Xiphothecaella strigosa KOZUR, MOIX & OZSVÁRT, 2007
holotype, M 2008.61.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Koz07a, p. 164, pl. 2, fig. 9
Authors' inventory number: 23-9-04/II-116

Zhamojdasphaera latispinosa mersinensis KOZUR, MOIX & MOSTLER, 2007
holotype, M 2008.113.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Moi07, p. 297, pl. 4, fig. 3
Authors' inventory number: 23-9-04/IV-30

Zhamojdasphaera rigoi rigoi KOZUR, MOIX & MOSTLER, 2007
holotype, M 2008.114.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Moi07, p. 297, pl. 4, fig. 2
Authors' inventory number: 23-9-04/V-90

Zhamojdasphaera rigoi rigoi KOZUR, MOIX & MOSTLER, 2007
paratype, M 2008.114.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Moi07, p. 297, pl. 4, figs. 6a–b
Authors' inventory number: 23-9-04/VI-159

Zhamojdasphaera rigoi brevispinosa KOZUR, MOIX & MOSTLER, 2007
holotype, M 2008.115.1.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Moi07, p. 297, pl. 4, fig. 5
Authors' inventory number: 23-9-04/V-200

Zhamojdasphaera rigoi brevispinosa KOZUR, MOIX & MOSTLER, 2007
paratype, M 2008.115.2.
Carnian, Killik Formation
southwest of Tavuşçayırı Tepe, Turkey
Moi07, p. 297, pl. 4, fig. 4
Authors' inventory number: 8-12-03/I-12

2.2. Triassic Bivalvia

Cornucardia turcensis VÉGH-NEUBRANDT, 1976
holotype, 2004.46.1.
Carnian, "marl with reefal limestone blocks"
Tilkideligi Tepe, Korkuteli, Turkey
V-N76, p. 211, pl. 4, figs. 8–10
Inventory number in original publication: 245A

- Dicerocardium baconicum* VÉGH-NEUBRANDT, 1974
holotype, 2004.16.1.
Norian, Main Dolomite Formation
Nyírespuszta, Nagytárkány, Csabrendek
V-N74, p. 27 (18), pl. 2, fig. 2, pl. 10, figs. 1a-b
- Dicerocardium furcatolateratum* VÉGH-NEUBRANDT, 1974
holotype, 2004.21.1.
Norian, Main Dolomite Formation
trench east of Deákpuszta, Nyirág
V-N74, p. 27 (17), pl. 8, figs. 1a-b, pl. 9, fig. 1c
- Dicerocardium pteriforme* VÉGH-NEUBRANDT, 1963
holotype, 2004.14.1.
Norian, Dachstein Limestone Formation
Vinye Sándor-major, Cuha-völgy,
Bakonyzentlászló
V-N63, p. 339 (336), pl. 17, figs. 1a-e
Inventory number in original publication:
12.158
- Dicerocardium pteriforme* VÉGH-NEUBRANDT, 1963
paratype, 2004.33.1.
Norian, Dachstein Limestone Formation
Vinye Sándor-major, Cuha-völgy,
Bakonyzentlászló
V-N63, p. 339 (336)
- Lima chinensis* LÓCZY, 1897
syntypes
Middle Triassic, "yellow sandstone and shale"
Chung-tien Plateau (3500 m a.s.l.), China
Lóc97, p. 121, pl. 10, figs. 8-11
Translated description in German edition,
LÓCZY 1899: p. 142. Types probably lost
- Megalodon subcolumbella* KOKEN, 1913
neotype, 2004.18.1.
Norian, Main Dolomite Formation
quarry beside Sümeg-Tapolca road, Újdörög,
Zalahaláp
Kok13, p. 32
Rossiodus subcolumbella, V-N82
Neotype designated and figured in VÉGH-
NEUBRANDT 1982: p. 324, text-fig. 153b

- Megalodus arcuatus* VÉGH-NEUBRANDT, 1963
replica of holotype, 2006.105.1.
Norian, Dachstein Limestone Formation
Várbük, Csesznek
V-N63, p. 338 (335), pl. 15, figs. 1a-c
Neomegalodon arcuatus arcuatus, V-N82
Inventory number in original publication:
12.181
- Megalodus arcuatus inflatus* VÉGH-NEUBRANDT, 1963
holotype, 2004.1.1.
Norian, Dachstein Limestone Formation
Vinye Sándor-major, Cuha-völgy,
Bakonyzentlászló
V-N63, p. 339 (335), pl. 16, figs. 1a-c
Neomegalodon arcuatus inflatus, V-N82
Inventory number in original publication:
12.176
- Myophoria kreitneri* LÓCZY, 1897
syntypes
Middle Triassic, "yellow sandstone and shale"
Chung-tien Plateau (3500 m a.s.l.), China
Lóc97, p. 132, pl. 9, figs. 25, 26
Translated description in German edition,
LÓCZY 1899: p. 156. Types probably lost
- Myophoria radiata* LÓCZY, 1897
syntypes
Middle Triassic, "yellow sandstone and shale"
Chung-tien Plateau (3500 m a.s.l.), China
Lóc97, p. 131, pl. 9, figs. 21, 22
Translated description in German edition,
LÓCZY 1899: p. 155. Types probably lost
- Myophoria széchenyi* LÓCZY, 1897
syntypes
Middle Triassic, "yellow sandstone and shale"
Chung-tien Plateau (3500 m a.s.l.), China
Lóc97, p. 130, pl. 9, figs. 14-20
Translated description in German edition,
LÓCZY 1899: p. 153. Types probably lost
- Neomegalodon (Gemmelarodus) paronai*
praenoricus VÉGH-NEUBRANDT, 1974
holotype, 2004.34.1.
Carnian, Main Dolomite Formation
Aranyos-völgy, Veszprém
V-N74, p. 24 (12), pl. 4, figs. 5a-d (pl. 4,
figs. 7-8, pl. 7, fig. 3)

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratype, 2004.35.1.
Carnian, Main Dolomite Formation
Aranyos-völgy, Veszprém
V-N74, p. 24 (12), pl. 4, figs. 6a-d

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratype, 2004.36.1.
Carnian, Main Dolomite Formation
Aranyos-völgy, Veszprém
V-N74, p. 24 (12), pl. 4, figs. 4a-d

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratype, 2004.39.1.
Carnian, Main Dolomite Formation
Kisgellérthegy, Budapest
V-N74, p. 24 (12), pl. 3, figs. 3a-d

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratype, 2004.40.1.
Carnian, Main Dolomite Formation
Vörös-hegy, Gyermely
V-N74, p. 24 (12), pl. 3, figs. 1a-d

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratypes, 2004.41.1-4., 2004.45.1-22.
Carnian, Main Dolomite Formation
Aranyos-völgy, Veszprém
V-N74, p. 24 (12)

Neomegalodon (Gemmelarodus) paronai praenorius VÉGH-NEUBRANDT, 1974
paratype, 2006.88.1.
Carnian, Main Dolomite Formation
Gánt
V-N74, p. 24 (12), pl. 3, figs. 2a-d

Parallelodon baconicum VÉGH, 1964
holotype, 2004.37.1.
Norian, Main Dolomite Formation
500 m north of railway station, east side of
Hárskúti út, Márkó
Vég64, p. 339 (336), pl. 23, fig. 1
Inventory number in original publication:
12.186

Physocardia julii VÉGH-NEUBRANDT, 1974
holotype, 2004.20.1.
Carnian, Veszprém Marl Formation
Öregszőlök, Csákberény
V-N74, p. 28 (18), pl. 5, figs. 2a-c

Pinna transdanubica VÉGH, 1964
holotype, 2004.3.1.
Norian, Main Dolomite Formation
between Kajmát quarry and road to Balinka,
Bodajk
Vég64, p. 339 (336), pl. 23, figs. 2a-c
Inventory number in original publication:
12.187

Rhaetomegalodon acutareatus VÉGH-NEUBRANDT,
1974
holotype, 2004.2.1.
Rhaetian, Dachstein Limestone Formation
Héthézpuszta quarry, Bakonycsernye
V-N74, p. 26 (16), pl. 7, figs. 1a-c

Rhaetomegalodon bajotensis VÉGH-NEUBRANDT,
1969
holotype, M.97.67
Rhaetian, Dachstein Limestone Formation
Öreg-kő, Bajót
V-N69, p. 123
Inventory number in original publication:
12.212/a. Type figured in VIGH 1914: text-fig.
48, pl. 3, fig. 3 (as *Megalodus eupalliatius* FRECH).
Text-fig. 178 in VÉGH-NEUBRANDT 1982 shows
a different specimen

Rhaetomegalodon bajotensis VÉGH-NEUBRANDT,
1969
paratype, M.96.435
Rhaetian, Dachstein Limestone Formation
Öreg-kő, Bajót
V-N69, p. 123
Inventory number in original publication:
12.212/b

Rhaetomegalodon incisus gutnici VÉGH-
NEUBRANDT, 1976
paratype, 2004.47.1.
Rhaetian, Leylek Limestone Formation
Sorkun Yayla, Beyşehir, Turkey
V-N76, p. 209 (pl. 3, figs. 9-11)

Triadomegalodon rátóti VÉGH-NEUBRANDT, 1974
holotype, 2004.17.1.
Norian, Main Dolomite Formation
quarry between Rátót and Eplény, Eplény
V-N74, p. 22 (11), pl. 1, figs. 1a–b, pl. 2, fig. 1c

2.3. Triassic Gastropoda

Hungariella kutassyi SZABÓ, 2007
holotype, 2007.2.1.
Carnian, Dachstein Limestone Formation
Fazekas-hegy, Budapest
Szb07, p. 73, pl. 1, figs. 18–20
Referred to as lectotype in SZABÓ 2007

Hungariella stredae KUTASSY, 1933
lectotype, 2007.4.1.
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut33, p. 241 (pl. 1, figs. 10–14)
Lectotype designated and figured in SZABÓ
2007: pl. 1, figs. 12–13

Hungariella stredae KUTASSY, 1933
paralectotype, M.97.34
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut33, p. 241
Type figured in SZABÓ 2007: pl. 1, figs. 16–17

Hungariella stredae KUTASSY, 1933
paralectotypes, M.97.54
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut33, p. 241

Hungariella stredae KUTASSY, 1933
paralectotype, 2007.6.1.
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut33, p. 241
Type figured in SZABÓ 2007: pl. 1, figs. 14–15

Hungariella stredae KUTASSY, 1933
paralectotype, 2007.7.1.
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut33, p. 241
Type figured in SZABÓ 2007: pl. 1, figs. 10–11

Neritopsis rappi KUTASSY, 1927
neotype, 2007.8.1.
Norian, Dachstein Limestone Formation
Remete-hegy, Máriaremete, Budapest
Kut27, p. 153 (pl. 3, figs. 10a–d)
Hungariella rappi, Kut33; Szb07
Neotype designated and figured in SZABÓ
2007: pl. 1, figs. 6–9

2.4. Triassic Cephalopoda

Encoiloceras balatonicum VÖRÖS, 2001
holotype, M.99.105
Anisian, Felsőörs Limestone Formation
Farkó-kő I., Aszófő
VöA01, p. 4, text-figs. 3a, 3c, pl. 1, fig. 2

Encoiloceras balatonicum VÖRÖS, 2001
paratype, M.99.106
Anisian, Felsőörs Limestone Formation
Farkó-kő I., Aszófő
VöA01, p. 4, text-fig. 3b, pl. 1, figs. 3a–b

Encoiloceras lajosi VÖRÖS, 2001
holotype, M.99.107
Anisian, Felsőörs Limestone Formation
Farkó-kő I., Aszófő
VöA01, p. 5, text-fig. 4a, pl. 2, figs. 1a–b

Encoiloceras lajosi VÖRÖS, 2001
paratype, M.99.108
Anisian, Felsőörs Limestone Formation
Farkó-kő I., Aszófő
VöA01, p. 5, text-figs. 4b, 4d, pl. 1, fig. 4

Encoiloceras lajosi VÖRÖS, 2001
paratype, M.99.114
Anisian, Felsőörs Limestone Formation
Farkó-kő I., Aszófő
VöA01, p. 5, text-fig. 4c, pl. 1, figs. 5a–b

Encoiloceras lajosi VÖRÖS, 2001
paratype, M.99.115
Anisian, Felsőörs Limestone Formation
Farkó-kő II., Aszófő
VöA01, p. 5, pl. 1, figs. 6a–c

2.5. Triassic Brachiopoda

Spiriferina subfragilis LÓCZY, 1897

syntypes

Middle Triassic, "yellow sandstone and shale"
Chung-tien Plateau (3500 m a.s.l.), China
Lócz97, p. 133, pl. 10, figs. 27–34
Translated description in German edition,
LÓCZY 1899: p. 158. Types probably lost

Thecocystella horogensis PÁLFY, 2003

holotype, M.92.492

Anisian, Felsőrs Limestone Formation
Horog-hegy, Köveskál
Pál03, p. 148, pl. Br-1, figs. 23a–e

Thecocystella horogensis PÁLFY, 2003

paratype, M.2002.866

Anisian, Felsőrs Limestone Formation
Horog-hegy, Köveskál
Pál03, p. 148, pl. Br-1, fig. 34

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.52

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, fig. 9

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.53

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, fig. 2

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.54

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.55

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, fig. 3

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.56

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, figs. 7, 8

2.6. Triassic Ostracoda

Triadogigantocypris balatonica MONOSTORI,
1991

holotype, M.90.1

Ladinian, Buchenstein Formation
Öreg-hegy, P/IIa section, Vászoly
Mon91, p. 94, text-fig. 2

3.2. Jurassic Anthozoa

Isastraea hodnaensis COQUAND, 1880

holotype, M.63.3704

Kimmeridgian
road between d'Ain-Rich and Aïn-M'garnès,
Hodna, Algeria
Coq80, p. 354

Microsolena rotula pannonica KOLOSVÁRY, 1954

syntypes, M.57.3088, M.57.3089, M.57.3097

Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár

Kol54b, p. 239, pl. 40, figs. 1–12, pl. 41, figs. 1–4

Microsolena rotula pannonica KOLOSVÁRY, 1954

syntype, M.57.3094

Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár

Kol54b, p. 239, pl. 40, figs. 1–12, pl. 41, figs. 1–4
Type missing

3. Jurassic types

3.1. Jurassic Foraminiferida

Globuligerina geczyi GÖRÖG, 1994

holotype, M.93.51

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, figs. 1, 4

Globuligerina geczyi GÖRÖG, 1994

paratype, M.93.50

Hettangian, Pisznice Limestone Formation
Nagy-Pisznic, Lábatlan
Gör94, p. 256, pl. 1, figs. 5, 6

- Thecocystathus baconica* KOLOSVÁRY, 1954
holotype, M.57.3093
Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár
Kol54b, p. 237, pl. 38, figs. 3–4
- Thecocystathus mactraeformis* KOLOSVÁRY, 1954
syntypes, M.57.3091
Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár
Kol54b, p. 237, pl. 37, figs. 3–7
- Undulocystathus hungaricus* KOLOSVÁRY, 1954
holotype, M.57.3090
Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár
Kol54b, p. 238, pl. 38, fig. 1
Type missing
- Undulocystathus noszkyi* KOLOSVÁRY, 1954
holotype, M.57.3303
Tithonian, Pálihálás Limestone Formation
Pás kom-tető, Borzavár
Kol54b, p. 238, pl. 37, figs. 8–9
Type missing
- 3.3. Jurassic Bivalvia**
- Anomia jarnacensis* COQUAND, 1860
syntypes, M.60.1140
Tithonian
Jarnac, France
Coq60, p. 37
- Corbula carentonensis* COQUAND, 1860
syntypes, M.60.907
Tithonian–Berri asian
Champ-Blanc near Cognac, France
Coq60, p. 38
- Corbula condamyi* COQUAND, 1860
syntype, M.60.1471
Tithonian–Berri asian
Montgaud, France
Coq60, p. 38
- Corbula jarnacensis* COQUAND, 1860
syntype, M.60.906
Tithonian–Berri asian
Champ-Blanc near Cognac, France
Coq60, p. 38
- Corbula purbeckensis* COQUAND, 1860
syntypes, M.60.877
Tithonian–Berri asian
Champ-Blanc near Cognac, France
Coq60, p. 39
- Cyclas arnaudi* COQUAND, 1860
syntype, M.60.1472
Tithonian–Berri asian
Montgaud, France
Coq60, p. 39
- Cypricardia vatonnei* COQUAND, 1880
holotype, M.60.1670
Pliensbachian
Djebel-Ouarsenis, Algeria
Coq80, p. 115
- Gervillia pančići* RADOVANOVIC, 1900
syntype, M.63.3532
Aalenian, Vrška Čuka Formation
Vrška Čuka, Serbia
Rdv00, p. 64, pl. 1, fig. 5 (pl. 1, fig. 4)
Gervillella aviculoides, Rdv97
- Gervillia pančići* RADOVANOVIC, 1900
syntypes, M.63.3532
Aalenian, Vrška Čuka Formation
Vrška Čuka, Serbia
Rdv00, p. 64
- Isocardia maresi* COQUAND, 1880
holotype, M.60.1657
Pliensbachian
Djebel-Ouarsenis, Algeria
Coq80, p. 113

- Lima hodnensis* COQUAND, 1880
holotype, M.63.3711
Kimmeridgian
road between d'Aïn-Rich and Aïn-M'garnès,
Hodna, Algeria
Coq80, p. 140
- Lima (Plagiostoma) villei* COQUAND, 1880
holotype, M.60.1655
Pliensbachian
Ouarsenis, Algeria
Coq80, p. 139
- Limea (Pseudolimea) galaczi* SZENTE, 1995
holotype, M.95.1
Bathonian, Óbánya Limestone Formation
Hidasi-völgy, Hosszúhetény
Sze95, p. 100, pl. 3, figs. 1, 8 (pl. 2, figs. 9, 10,
pl. 3, figs. 2–6)
- Pecten jarnacensis* COQUAND, 1860
syntype, M.60.874
Tithonian
Chez-Ville, France
Coq60, p. 37
- Pecten jarnacensis* COQUAND, 1860
syntypes, M.60.1145
Tithonian
Jarnac, France
Coq60, p. 37
- Pecten menisa* COQUAND, 1880
holotype, M.63.3709
Kimmeridgian
road between d'Aïn-Rich and Aïn-M'garnès,
Hodna, Algeria
Coq80, p. 149
- Pholadomya schafarziki* PAPP, 1907
holotype, M.63.3198
Callovian–Oxfordian, “limestone”
Gunib, Daghestan, Russia
Pap07, p. 155, pl. 3, figs. 1, 1a, 1b
- Pleuromya merzbacheri* PAPP, 1907
holotype, M.63.3616
Callovian
Psebay, Russia
Pap07, p. 158, pl. 4, figs. 2, 2a–c

3.4. Jurassic Gastropoda

- Adeorbisina conica* SZABÓ, 1996
holotype, M.96.351
Bajocian, fissure fill, Eplény Limestone
Formation?
Som-hegy, Bakonybél
Szb96, p. 65, pl. 1, figs. 1–4
- Adeorbisina nitens* SZABÓ, 1996
holotype, M.96.352
Bajocian, fissure fill, Eplény Limestone
Formation?
Som-hegy, Bakonybél
Szb96, p. 65, pl. 1, figs. 5–8
- Ampullaria pisum* COQUAND, 1860
holotype, M.60.904
Tithonian–Berriasiian
Toinot near Cognac, France
Coq60, p. 38
- Helcion extlectorius* COQUAND, 1880
syntypes, M.60.1672
Pliensbachian
Djebel-Ouarsenis, Algeria
Coq80, p. 90
- Melania boreau* COQUAND, 1860
holotype, M.60.908
Tithonian–Berriasiian
Champ-Blanc near Cognac, France
Coq60, p. 38
- Nerinea perforans* COQUAND, 1880
holotype, M.63.3712
Kimmeridgian
road between d'Aïn-Rich and Aïn-M'garnès,
Hodna, Algeria
Coq80, p. 49
- Physa boreau* COQUAND, 1860
holotype, M.60.1469
Tithonian–Berriasiian
Montgaud, France
Coq60, p. 38
- Trochus guilhoti* COQUAND, 1860
holotype, M.60.1998
Toarcian
Saint-Gervais, France
Coq60, p. 7

Trochus ouarsenensis COQUAND, 1880
 syntype, M.60.1649
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 62

Turbo afer COQUAND, 1880
 syntype, M.60.1666
 Pliensbachian
 Ouarsenis, Algeria
 Coq80, p. 64

Turbo nabdalsæ COQUAND, 1880
 syntype, M.60.1673
 Pliensbachian
 Ouarsenis, Algeria
 Coq80, p. 65

Turbo nicaisei COQUAND, 1880
 syntypes, M.60.1667
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 64

3.5. Jurassic Cephalopoda

Alocolytoceras ophioneum toarcense GÉCZY, 1967
 paratype, M.66.596
 Toarcian, Tölgyhát Limestone Formation
 Bakonycsernye
 Géc67, p. 80 (text-fig. 83, pl. 23, fig. 6,
 pl. 64, fig. 33)
 Type missing

Ammonites letourneuxi COQUAND, 1880
 holotype, M.60.1658
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 6

Ammonites peraffinis COQUAND, 1880
 holotype, M.63.3721
 Oxfordian
 Foum-Islamen, near Batna, Algeria
 Coq80, p. 8

Aptychus beaumontii COQUAND, 1841
 holotype, M.60.664
 Oxfordian
 Vergons, Basses-Alpes, France
 Coq41, p. 388, pl. 9, fig. 12

Calliphylloceras altisulcatum quadratum GÉCZY, 1967
 holotype
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
 Géc67, p. 39, text-fig. 36, pl. 9, fig. 2, pl. 63,
 fig. 40
 Type in GIH collection as J.2395 but quoted in
 original publication as in HNHM

Calliphylloceras altisulcatum magnum GÉCZY, 1967
 paratypes, M.66.347, M.66.350, M.66.355,
 M.66.357, M.66.360, M.66.362, M.66.370,
 M.66.371, M.66.376, M.66.378, M.83.217
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
 Géc67, p. 38 (text-fig. 35, pl. 9, figs. 1, 3, pl. 63,
 figs. 37–39)

Calliphylloceras liasicum GÉCZY, 1967
 paratypes, M.66.458, M.66.496, M.66.510,
 M.66.517, M.66.600, M.66.641
 Pliensbachian, Túzkövesárok Limestone
 Formation
 Bakonycsernye
 Géc67, p. 25 (text-fig. 19, pl. 8, figs. 1, 2)

Costileioceras sinon viallii GÉCZY, 1967
 paratypes, M.66.699, M.83.198
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
 Géc67, p. 175 (text-fig. 178, pl. 38, figs. 4, 7,
 pl. 39, fig. 6, pl. 65, figs. 4, 5, 73)

Erycites ovatus GÉCZY, 1966
 holotype
 Toarcian–Aalenian, Tölgyhát Limestone
 Formation
 Bakonycsernye
 Géc66, p. 104, text-fig. 91, pl. 30, fig. 1, pl. 42,
 fig. 10
 Type in GIH collection as J.2315 but quoted in
 original publication as in HNHM

- Erycites ovatus? rogeri* GÉCZY, 1966
holotype
Toarcian-Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 105, text-fig. 92, pl. 30, fig. 3, pl. 42, fig. 11
Type in GIH collection as J.2316 but quoted in original publication as in HNHM
- Erycites subquadratus* GÉCZY, 1966
holotype
Toarcian-Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 95, text-fig. 81, pl. 27, fig. 1, pl. 42, fig. 2
Type in GIH collection as J.2306 but quoted in original publication as in HNHM
- Erycites subquadratus* GÉCZY, 1966
paratypes, M.66.855, M.66.862
Toarcian-Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 95
- Erycites telegdirothi amplus* GÉCZY, 1966
holotype
Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 89, text-fig. 74, pl. 24, fig. 3, pl. 41, fig. 10
Type in GIH collection as J.2301 but quoted in original publication as in HNHM
- Hammatoceras meneghinii raricostatum* GÉCZY, 1966
holotype
Toarcian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 84, text-fig. 71, pl. 23, fig. 1, pl. 41, fig. 3
Type in GIH collection as J.2298 but quoted in original publication as in HNHM
- Hammatoceras planiforme stredai* GÉCZY, 1966
holotype
Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 57, text-fig. 46, pl. 12, fig. 1, pl. 39, fig. 2
Type in GIH collection as J.2271 but quoted in original publication as in HNHM
- Hammatoceras planinsigne merlai* GÉCZY, 1966
holotype
Toarcian-Aalenian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 61, text-fig. 48, pl. 13, fig. 1, pl. 39, fig. 5
Type in GIH collection as J.2275 but quoted in original publication as in HNHM
- Hammatoceras tenerum szoerenyiæ* GÉCZY, 1966
holotype
Aalenian-Bajocian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 55, text-fig. 44, pl. 10, fig. 4, pl. 38, fig. 15
Type in GIH collection as J.2269 but quoted in original publication as in HNHM
- Hammatoceras tenerum szoerenyiæ* GÉCZY, 1966
paratype, M.66.733
Aalenian-Bajocian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 55
Type missing
- Hammatoceras tenerum szoerenyiæ* GÉCZY, 1966
paratype, M.66.734
Aalenian-Bajocian, Tölgyhát Limestone Formation
Bakonycsernye
Géc66, p. 55
- Ludwigia vaceki* GÉCZY, 1967
paratype, M.66.661
Aalenian?, Tölgyhát Limestone Formation
Bakonycsernye
Géc67, p. 197 (text-fig. 205, pl. 46, fig. 5, pl. 65, fig. 33)

- Lytoceras amplum kocsisi* GÉCZY, 1967
 paratypes, M.66.562, M.66.597
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
Géc67, p. 74 (text-fig. 77, pl. 20, fig. 7, pl. 21,
 figs. 1, 3, pl. 64, fig. 28)
- Lytoceras haasi* GÉCZY, 1967
 holotype
 Pliensbachian, Túzkővesárok Limestone
 Formation
 Bakonycsernye
Géc67, p. 59, text-fig. 57, pl. 16, fig. 3
 Type in GIH collection as J.2429 but quoted in
 original publication as in HNHM
- Lytoceras vaceki* GÉCZY, 1967
 paratypes, M.66.527, M.66.528, M.66.529,
 M.66.534, M.66.536, M.66.547, M.66.552,
 M.66.559, M.66.560, M.66.564, M.66.566
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
Géc67, p. 68 (text-fig. 71, pl. 22, fig. 3, pl. 23,
 figs. 2, 4, pl. 64, figs. 22, 23)
- Perisphinctes lóczyi* PAPP, 1907
 syntypes, M.63.3301
 Bajocian, "lower dark shale"
 Gunib, Daghestan, Russia
Pap07, p. 165, pl. 7, figs. 1, 2, 2a, 2b
- Phylloceras? baconicum mauveugei* GÉCZY, 1967
 paratypes, M.66.346, M.66.361, M.66.438,
 M.66.476, M.66.522, M.66.619, M.66.620,
 M.66.621, M.66.622, M.66.623, M.66.624,
 M.66.625, M.66.626, M.66.627, M.66.630,
 M.66.632, M.66.633, M.66.634, M.66.636,
 M.66.637, M.66.640, M.66.642, M.66.643,
 M.66.644, M.66.645, M.66.646, M.66.648,
 M.66.650, M.66.652, M.66.653, M.66.654,
 M.66.656, M.66.657, M.66.658, M.68.248,
 M.68.249, M.83.206, M.83.215, M.83.245,
 M.83.248
 Aalenian, Tölgyhát Limestone Formation
 Bakonycsernye
Géc67, p. 19 (text-fig. 14, pl. 4, figs. 2–4,
 pl. 43, figs. 12–14)
- Rhacophyllites ssemenovi* PAPP, 1907
 holotype, M.63.3634
 Early Jurassic
 Terek River, Russia
Pap07, p. 162, pl. 6, figs. 3, 3a, 3b

- Stephanoceras liechtensteinii* PAPP, 1907
 holotype, M.63.3610
 Bajocian, "dark ferruginous limestone boulder"
 Fiagdon Creek, west of Vladikavkaz, Russia
Pap07, p. 163, pl. 6, fig. 4

3.6. Jurassic Brachiopoda

- Dichotomosella galaczi* VÖRÖS, 1995
 holotype, M.94.92
 Bathonian, Óbánya Limestone Formation
 Hidasi-völgy, Zobákpuszta, Hosszúhetény
VöA95, p. 188, pl. 1, figs. 4a–c

- Dichotomosella galaczi* VÖRÖS, 1995
 paratype, M.94.93
 Bathonian, Óbánya Limestone Formation
 Márévári-völgy, Magyaregregy
VöA95, p. 188, text-fig. 3

- Karadagithyris eduardi* VÖRÖS, 1995
 holotype, M.94.109
 Bathonian, Óbánya Limestone Formation
 Hidasi-völgy, Zobákpuszta, Hosszúhetény
VöA95, p. 199, pl. 1, figs. 10a–c

- Karadagithyris eduardi* VÖRÖS, 1995
 paratype, M.94.110
 Bathonian, Óbánya Limestone Formation
 Sövér-árok, Magyaregregy
VöA95, p. 199, text-fig. 9

- Lokutella kondai* VÖRÖS, 1983
 holotype, M.82.1
 Pliensbachian, Hierlatz Limestone
 Kőris-hegy, Bakonyszücs
VöA83, p. 8, text-figs. 3a–c

- Rhynchonella guerrizila* COQUAND, 1880
 syntypes, M.60.1671
 Pliensbachian
 Djebel-Ouarsenis, Algeria
Coq80, p. 198

- Rhynchonella mazana* COQUAND, 1880
 syntypes, M.60.1654
 Pliensbachian
 Djebel-Ouarsenis, Algeria
Coq80, p. 199

Rhynchonella meridionalis COQUAND, 1862
 syntypes, M.60.644
 Pliensbachian
 Brignolles, France
 Coq62, p. 322, pl. 35, figs. 14, 15, 15a
Rhynchonella meridionalis DESLONGCHAMPS,
 1863 is a junior primary homonym,
Homoeorhynchia batalleri (DUBAR, 1931) is
 regarded as a subjective junior synonym
 (see ALMÉRAS 1996) of this species

Rhynchonella meridionalis COQUAND, 1862
 syntypes, M.60.957
 Pliensbachian
 Cuers, France
 Coq62, p. 322, pl. 35, figs. 14, 15, 15a

Rhynchonella meridionalis COQUAND, 1862
 syntypes, M.60.1372, M.60.1397
 Pliensbachian
 Mazaugues, France
 Coq62, p. 322, pl. 35, figs. 14, 15, 15a

Rhynchonella meridionalis COQUAND, 1862
 syntypes, M.60.1565
 Pliensbachian
 Plan-d'Aups, France
 Coq62, p. 322, pl. 35, figs. 14, 15, 15a

Rhynchonella meridionalis COQUAND, 1862
 syntypes, M.60.1668
 Pliensbachian
 Ouarsenis, Algeria
 Coq62, p. 322, pl. 35, figs. 14, 15, 15a

Rhynchonella nicaisei COQUAND, 1880
 syntypes, M.60.1647, M.60.1648
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 201

Rhynchonella zaboul COQUAND, 1880
 syntypes, M.60.1653
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 200

Spiriferina africana COQUAND, 1880
 syntypes, M.60.1651
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 197

Spiriferina nicaisei COQUAND, 1880
 holotype, M.60.1669
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 198

Terebratula (Waldheimia) antalas COQUAND, 1880
 syntypes, M.60.1650
 Pliensbachian
 Ouarsenis, Algeria
 Coq80, p. 202

Terebratula (Waldheimia) jabdas COQUAND, 1880
 holotype, M.60.1659
 Pliensbachian
 Djebel-Ouarsenis, Algeria
 Coq80, p. 204

Terebratula peyssoneli COQUAND, 1880
 syntypes, M.60.2024
 Kimmeridgian
 Seba-Liamone, Algeria
 Coq80, p. 207

4. Cretaceous types

4.1. Cretaceous Anthozoa

Ceratotrochus bakonyensis KOLOSVÁRY, 1954
 syntype, M.57.3130
 Aptian, Pénzeskút Marl Formation
 Pénzeskút, Hásagy, Pénzesgyőr
 Kol54a, p. 126 (98), pl. 8, figs. 17, 21
 Type missing

Ceratotrochus noszkyi KOLOSVÁRY, 1954
 syntypes, M.57.3226, M.57.3227
 Aptian, Pénzeskút Marl Formation
 Pénzeskút, Hásagy, Pénzesgyőr
 Kol54a, p. 126 (98), pl. 8, fig. 22, pl. 9, figs. 1–6
 Types missing

Cladocora schréteri KOLOSVÁRY, 1954
 syntypes, M.57.3164
 Late Cretaceous, Nekézseny Conglomerate
 Formation
 Jócsós-völgy, Nekézseny
 Kol54a, p. 125 (82), pl. 5, figs. 8–13

- Coelosmilia niobe* KOLOSVÁRY, 1954
syntype, M.57.3222
Late Cretaceous, Jákó Marl Formation
Öreghegy quarry, Sümeg
Kol54a, p. 127 (105), pl. 11, figs. 16–17, pl. 12,
figs. 1–3
- Helicopora neocomiensis* KOLOSVÁRY, 1954
holotype, M.57.3124
Albian, Lábatlan Sandstone Formation
Köszörűkő quarry, Lábatlan
Kol54a, p. 128 (115), pl. 6, fig. 7
- Montlivaltia obesaformis* KOLOSVÁRY, 1954
syntypes, M.57.3138
Aptian, Pénzeskút Marl Formation
Pénzeskút, Hársgy, Pénzesgyőr
Kol54a, p. 125 (79), pl. 5, figs. 1–4
- Phyllosmilia sümegensis* KOLOSVÁRY, 1954
holotype, M.57.3221
Late Cretaceous, Jákó Marl Formation
Öreghegy quarry, Sümeg
Kol54a, p. 128 (108), pl. 12, figs. 11–12
- Phyllosmilia upponensis* KOLOSVÁRY, 1954
holotype, M.57.3158
Late Cretaceous, Nekézseny Conglomerate
Formation
Jócsös-völgy, Nekézseny
Kol54a, p. 128 (107), pl. 12, figs. 5–6
- Placosmilia? hungarica* KOLOSVÁRY, 1954
holotype, M.57.3217
Late Cretaceous, Jákó Marl Formation
Ajka
Kol54a, p. 127 (105), pl. 11, figs. 7–8
- Stephanosmilia polydectes* KOLOSVÁRY, 1954
syntypes, M.57.3171
Late Cretaceous, Jákó Marl Formation
Öreghegy quarry, Sümeg
Kol54a, p. 127 (103), pl. 10, figs. 4–9
- Stylocyathus minutus* KOLOSVÁRY, 1954
holotype, M.57.3215
Albian–Cenomanian, Pénzeskút Marl
Formation
drill core, Olaszfalu
Kol54a, p. 126 (99), pl. 16, figs. 1–7
Type missing

4.2. Cretaceous Bivalvia

- Lucina zamma* COQUAND, 1880
syntypes, M.63.4064
Early Cretaceous
Djebel-Ouach, Algeria
Coq80, p. 385

4.3. Cretaceous Gastropoda

- Aporrhais augustini* COQUAND, 1880
holotype, M.63.3715
Late Cretaceous
Ras-Ouzina, Sétif, Algeria
Coq80, p. 72

- Buccinum cretaceum* COQUAND, 1862
holotype, M.63.3695
Turonian
Tébessa, Algeria
Coq62, p. 188, pl. 5, fig. 12
Aporrhais cretacea, Coq80

- Fusus conspicuus* COQUAND, 1862
holotype, M.63.3700
Late Cretaceous
Tébessa, Algeria
Coq62, p. 187, pl. 4, fig. 15

- Fusus julieni* COQUAND, 1862
syntype, M.63.3699
Late Cretaceous
Tébessa, Algeria
Coq62, p. 187, pl. 4, fig. 17

- Fusus strangulatus* COQUAND, 1862
syntype, M.60.2045
Late Cretaceous
Trik-Karretta, Algeria
Coq62, p. 187, pl. 4, fig. 14

- Fusus strangulatus* COQUAND, 1862
syntypes, M.63.3697
Late Cretaceous
Tébessa, Algeria
Coq62, p. 187, pl. 4, fig. 14

- Fusus tevesthenensis* COQUAND, 1862
syntypes, M.63.3701
Late Cretaceous
Tébessa, Algeria
Coq62, p. 187, pl. 4, fig. 13
- Natica gervaisi* COQUAND, 1862
syntypes, M.60.1151
Late Cretaceous
Kenchela, Algeria
Coq62, p. 180, pl. 4, fig. 1
- Nerinea gemmifera* COQUAND, 1862
syntype, M.63.4065
Late Cretaceous
col de Sfa, Biskra, Algeria
Coq62, p. 177, pl. 4, fig. 4
- Turritella leoperdites* COQUAND, 1862
holotype, M.63.3692
Late Cretaceous
Tébessa, Algeria
Coq62, p. 176, pl. 3, fig. 3
- 4.4. Cretaceous Cephalopoda**
- Aptychus blainvillei* COQUAND, 1841
holotype, M.60.2249
Early Cretaceous
Vérignon, Var, France
Coq41, p. 387, pl. 9, figs. 8, 9
- Deitanites labatlanensis* COMPANY, FÓZY,
SANDOVAL & TAVERA, 2006
holotype, 2004.170.1.
Barremian, Lábatlan Sandstone Formation
Bersek-hegy, Lábatlan
Com06, p. 9, pl. 1, fig. 17
- Deitanites labatlanensis* COMPANY, FÓZY,
SANDOVAL & TAVERA, 2006
paratypes, 2004.168.2., 2004.169.2., 2004.170.2–6.
Barremian, Lábatlan Sandstone Formation
Bersek-hegy, Lábatlan
Com06, p. 9
- Deitanites labatlanensis* COMPANY, FÓZY,
SANDOVAL & TAVERA, 2006
paratype, 2004.169.1.
Barremian, Lábatlan Sandstone Formation
Bersek-hegy, Lábatlan
Com06, p. 9, pl. 1, fig. 16
- Dufrenoyia katalinae* SZIVES, 2007
holotype, 2007.103.1.
Aptian, Tata Limestone Formation
Fazekas utca, Tata
Szi07a, p. 65, pl. 4, fig. 1
- Graysonites horvathi* SZIVES, 2007
holotype, 2007.68.1.
Albian, Pénzeskút Marl Formation
Jásd-1 quarry, Jásd
Szi07b, p. 109, pl. 24, fig. 12
- Heter ammonites ammoniticeras* COQUAND, 1880
paralectotype, M.63.3710
Turonian
Tébessa, Algeria
Coq80, p. 40
Lectotype fixed in PERVINQUIÈRE 1907: pl. 25,
fig. 3
- Menabites (Delawarella) suemegensis* FÓZY, 2001
holotype, M.63.1355
Campanian, Polány Marl Formation
Községi quarry, Sümeg
Fóz01, p. 32, pl. 4
- ?*Metascaphites kashaii* SZIVES, 2007
holotype, 2007.70.1.
Albian, Pénzeskút Marl Formation
Tilos-erdő, Pénzesgyör
Szi07b, p. 119, pl. 15, fig. 2
- ?*Metascaphites scholzi* SZIVES, 2007
holotype, 2007.69.1.
Albian, Pénzeskút Marl Formation
Tilos-erdő, Pénzesgyör
Szi07b, p. 120, pl. 16, fig. 18 (pl. 16, fig. 17)

Scaphites (Scaphites) evanicsi SZIVES, 2007
holotype, 2007.71.1.
Albian, Pénzeskút Marl Formation
Tilos-erdő, Pénzesgyör
Szi07b, p. 121, pl. 23, fig. 10 (pl. 18, figs. 3, 4,
10, 11)

Tonohamites boldii SZIVES & MONKS, 2002
holotype, 2007.20.1.
Aptian-Albian, Tata Limestone Formation
Eperkés-hegy, Olaszfalu
Szi02, p. 1140, text-fig. 4a

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.5.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5j-k

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.6.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5l-n

5. Paleocene types

5.1. Paleocene Brachiopoda

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
holotype, 2005.1.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 4a-c

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.2.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5a-c

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.3.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5d-f

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.4.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5g-i

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.8.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5q-s

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2005.9.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 5t-v

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2006.14.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-fig. 6

Basiliocostella kambueheli DULAI, BITNER &
MÜLLER, 2008
paratype, 2006.15.1.
Danian, Kambühel Formation
Kambühel, Austria
Dul08, p. 195, text-figs. 7a-b

6. Eocene types

6.1. Eocene Foraminiferida

Astrorhiza bakonycsernyensis OZSVÁRT, 2007
holotype, M 2008.116.1.

Bartonian, Padrag Marl Formation
borehole B-18, Bakonycsernye
Ozs07, p. 27, pl. 1, fig. 1

Clavulina cylindrica HANTKEN, 1875
paratype, M.99.47
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han75a (Han75b), p. 18 (14), pl. 1, fig. 8
Cylindroclavulina rutilosa, Hag56; Hor02
Type figured in HORVÁTH 2002. Type lost

Clavulina szaboi HANTKEN, 1868
paratype, M.99.43
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han68, p. 83, pl. 1, figs. 4a-b, 6a-b, 7a-b
Clavulinoides szaboi, Ozs07
Type figured in HORVÁTH 2002

Clavulinoides procerus OZSVÁRT, 2007
holotype, M 2008.117.1.
Bartonian, Padrag Marl Formation
borehole B-18, Bakonycsernye
Ozs07, p. 37, pl. 2, fig. 20

Cristellaria schwageri HANTKEN, 1875
paratype, M.99.65
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han75a (Han75b), p. 49 (42), pl. 5, fig. 11
Percultazonaria schwageri, Hor03
Type figured in HORVÁTH 2003

Dentalina gigantea HANTKEN, 1875
lectotype, 2006.93.1.
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han75a (Han75b), p. 34 (29), pl. 3, fig. 15
Nodosaria gigantea, Hor03
Lectotype designated in HORVÁTH 2003
under inventory number M.01.15

Dentalina intermedia HANTKEN, 1875
neotype, M.99.56
Priabonian, Buda Marl Formation
Síkfőkút, Noszvaj
Han75a (Han75b), p. 30 (25), pl. 3, figs. 4, 8
Laevidentalina intermedia, Hor03
Neotype designated in HORVÁTH 2003

Discocyclina concentrica KECSKEMÉTI, 1959
holotype, M.58.289
Bartonian, Szőc Limestone Formation
upper part of Köleskepe-árok, Ajka
Kec59, p. 45, text-fig. 10, pl. 2, figs. 6, 7, 9

Discocyclina concentrica KECSKEMÉTI, 1959
paratypes, M.61.641, M.61.646, M.61.720,
M.61.727
Bartonian, Szőc Limestone Formation
Köleskepe-árok, Ajka
Kec59, p. 45

Discocyclina hungarica KECSKEMÉTI, 1959
holotype, M.58.290
Bartonian, Szőc Limestone Formation
Csékúti-árok, Padrag, Ajka
Kec59, p. 51, text-fig. 14, pl. 3, figs. 5, 6, 8

Discocyclina hungarica KECSKEMÉTI, 1959
paratypes, M.66.159, M.66.160, M.66.161,
M.66.178, M.66.179, M.66.181
Bartonian, Szőc Limestone Formation
Csékúti-árok, Padrag, Ajka
Kec59, p. 51

Epistomaroides costatus OZSVÁRT, 2003
holotype, 2004.52.1.
Priabonian, Padrag Marl Formation
borehole Dv-4, Devecser
Ozs03, p. 3, pl. 1, figs. 6-7
Inventory numbers in original publication:
OP-4398, OP-4399, OP-4510, Collection of
Dept. of Paleontology, Eötvös University

Flabellina striata HANTKEN, 1875
neotype, M.99.61
Priabonian, Buda Marl Formation
Síkfőkút, Noszvaj
Han75a (Han75b), p. 43 (36), pl. 13, fig. 13
Plectofrondicularia striata, Maj62; Hor03
Neotype designated in HORVÁTH 2003

Gaudryina textilaroides HANTKEN, 1875

paralectotype, M.99.44

Priabonian, Buda Marl Formation

Vár-hegy, Budapest

Han75a (Han75b), p. 15 (12), pl. 1, fig. 6

Type figured in HORVÁTH 2002

Lingulina glabra HANTKEN, 1875

lectotype, M.99.59

Priabonian, Buda Marl Formation

Vár-hegy, Budapest

Han75a (Han75b), p. 42 (35), pl. 13, fig. 14

Lectotype designated in HORVÁTH 2003

Globigerina appanata HANTKEN, 1883

lectotype, M.72.553

Priabonian, "marl"

I'Escarène (Scarena), France

Han83a (Han83b), p. 11 (132), pl. 2, figs. 7a-c

Turborotalia appanata, Szt73

Lectotype designated in SZTRÁKOS 1973. Type missing

Nummulina bactchisaraiensis ROZLOZSNIK, 1929

lectotype, M 2008.6.1.

Lutetian

Bakhchisaray, Ukraine

Roz29, p. 118, 213, pl. 2, fig. 8, pl. 5, fig. 7

(pl. 2, figs. 11, 20, 27, 29, pl. 3, figs. 4, 17, 18, pl. 7, fig. 16)

Form A. Lectotype designated in SCHAUB 1981, using unjustifiably emended species name *bakhchisaraiensis*

Globigerina appanata HANTKEN, 1883

paralectotypes, M.72.554, M.72.555, M.72.556, M.72.557

Priabonian, "marl"

I'Escarène (Scarena), France

Han83a (Han83b), p. 11 (132), pl. 2, figs. 7a-c

Turborotalia appanata, Szt73

Types missing

Nummulina bactchisaraiensis ROZLOZSNIK, 1929

paralectotypes, M 2008.7.1.

Lutetian

Bakhchisaray, Ukraine

Roz29, p. 118, 213

Globigerina globosa HANTKEN, 1883

neotype, M.72.548

Priabonian, "marl"

I'Escarène (Scarena), France

Han83a (Han83b), p. 11 (132), pl. 2, figs. 3a-b

Globigerinatheca globosa, Szt73

Neotype designated in SZTRÁKOS 1973. Type missing

Nummulina böckhi ROZLOZSNIK, 1924

syntypes, M.61.782

Bartonian, Szőc Limestone Formation

Zsellér-dűlő, Tatabánya

Roz24a, p. 49

Form B

Gyroidinoides koestleri OZSVÁRT, 2003

holotype, 2004.53.1.

Priabonian, Padrag Marl Formation

borehole Cs-61, Csetény

Ozs03, p. 4, pl. 1, figs. 8-9

Inventory numbers in original publication:

OP-4467, OP-4468, Collection of Dept. of Paleontology, Eötvös University

Nummulina böckhi ROZLOZSNIK, 1924

syntypes, M.61.787

Bartonian, Szőc Limestone Formation

Zsellér-dűlő, Tatabánya

Roz24a, p. 49

Form A

Lagena clava OZSVÁRT, 2003

holotype, 2004.51.1.

Priabonian, Padrag Marl Formation

borehole Dv-4, Devecser

Ozs03, p. 2, pl. 1, fig. 5

Inventory number in original publication:

OP-4150, Collection of Dept. of Paleontology, Eötvös University

Nummulina boussaci ROZLOZSNIK, 1924

syntype, 2006.104.1.

Lutetian

Slani potak, Kossavín, Croatia

Roz24a, p. 49, 77, text-fig. 33c

Form B. Referred to as *N. lorioli* in German

edition, ROZLOZSNIK 1927: p. 87, text-fig. 33c.

More formal description in ROZLOZSNIK 1929:

p. 161. Type figured in ROZLOZSNIK 1929: pl. 6, fig. 9

Nummulina boussaci ROZLOZSNIK, 1924
 syntype, M 2008.8.1.
 Lutetian
 Slani potak, Kossavin, Croatia
 Roz24a, p. 49, 77, text-fig. 33c
 Form B. Type figured in ROZLOZSNIK 1929:
 pl. 4, fig. 9

Nummulina bronniarti d'archiaci ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.15.1.
 Bartonian, Szőc Limestone Formation
 Csékút, Ajka
 Roz24b, p. 171 (pl. 3, figs. 3a-d, 3f-g, pl. 5,
 figs. 9a-b)

Nummulina bronniarti d'archiaci ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.16.1., M 2008.17.1.,
 M 2008.18.1., M 2008.19.1.
 Bartonian, Szőc Limestone Formation
 Nagyganna, Ganna
 Roz24b, p. 171
 Form A

Nummulina bronniarti d'archiaci ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.20.1., M 2008.21.1.
 Bartonian, Szőc Limestone Formation
 Nagykovácsi
 Roz24b, p. 171

Nummulina bronniarti d'archiaci ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.22.1.
 Bartonian, Szőc Limestone Formation
 Porva
 Roz24b, p. 171
 Form A

Nummulina bronniarti d'archiaci ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.23.1.
 Bartonian, Szőc Limestone Formation
 Porva
 Roz24b, p. 171

Nummulina granifera frentanaeformis
 ROZLOZSNIK, 1929
 syntype, M.62.6299
 Lutetian
 Qafë Glumakë, Berat, Albania
 Roz29, p. 209 (pl. 2, figs. 1, 6, 28)
 Form B

Nummulina granifera frentanaeformis
 ROZLOZSNIK, 1929
 syntype, M.62.6300
 Lutetian
 Qafë Glumakë, Berat, Albania
 Roz29, p. 209
 Form A

Nummulina kovácsensis ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.24.1., M 2008.25.1.,
 M 2008.30.1.
 Bartonian, Szőc Limestone Formation
 Nagykovácsi
 Roz24b, p. 186 (pl. 4, figs. 10a-g)
 Form B

Nummulina kovácsensis ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.26.1., M 2008.27.1.,
 M 2008.28.1., M 2008.29.1., M 2008.31.1.
 Bartonian, Szőc Limestone Formation
 Nagykovácsi
 Roz24b, p. 186

Nummulina kovácsensis ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.32.1.
 Bartonian, Szőc Limestone Formation
 Domonkospuszta, Nagysáp
 Roz24b, p. 186
 Form A

Nummulina kovácsensis ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 syntypes, M 2008.33.1.
 Bartonian, Szőc Limestone Formation
 Tokod
 Roz24b, p. 186
 Form A

- Nummulina lászlói* ROZLOZSNIK, 1924
syntype, M.62.6244
Bartonian, Szőc Limestone Formation
Sas-hegy, Tokod
Roz24a, p. 21, 50, 62, 63, 64, 65, 66, 68, 70, 72,
73, 110 (text-figs. 1, 14a, 25–27, 29)
Form B. Referred to as *N. gizehensis* race *lászlói*
in German edition, ROZLOZSNIK 1927: p. 25 etc.
More formal description as *N. gizehensis* race
lászlói in ROZLOZSNIK 1929: p. 170. Type figured
in ROZLOZSNIK 1929: pl. 4, fig. 14
- Nummulina oppenheimi* ROZLOZSNIK, 1929
syntype, M 2008.9.1.
Ypresian
Purga di Bolca, Bolca, Italy
Roz29, p. 154, pl. 1, fig. 27 (pl. 7, fig. 7, pl. 8,
fig. 18)
Form A
- Nummulina striata pannonica* ROZLOZSNIK, 1924
syntypes, M.62.6275
Bartonian, Szőc Limestone Formation
Tokod
Roz24b, p. 179 (pl. 5, figs. 11b, 13a)
Form B
- Nummulina striata pannonica* ROZLOZSNIK, 1924
syntype, M 2008.12.1.
Bartonian, Szőc Limestone Formation
Piszke, Lábatlan
Roz24b, p. 179
Form A. Type figured in ROZLOZSNIK 1929:
pl. 6, fig. 20
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotypes, M.61.389, M.62.6251, M.62.9397,
M.62.9398, M.62.9399, M.62.9400, M.62.9401,
M.62.9402, M.62.9404, M.62.9405, M.62.9406,
M.62.9407, M.62.9408, M.62.9409, M.62.9410,
M.62.9411, M.62.9412, M.62.9413, M 2008.46.1.
Bartonian, Szőc Limestone Formation
Tokod
Roz24a, p. 24, 33, 37, 43, 45, 46, 48, 66, 67, 68, 69,
71, 72, 75, 77, 96, 105, 109, 128 (text-figs. 3, 10a,
11e–g, 21, 32b, pl. 1, fig. 1)
More formal description in ROZLOZSNIK 1924b:
p. 184. Lectotype designated in SCHAUB 1981,
based on drawings in ROZLOZSNIK 1924b: pl. 4,
figs. 5a–f, pl. 5, fig. 17. Original specimens not
identified but may be present in collection
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotypes, M.61.390, M.61.391,
M.62.6263, M.62.6294, M.62.6296,
M.62.6297, M.62.6327, M.62.6328,
M 2008.34.1., M 2008.35.1., M 2008.36.1.,
M 2008.37.1., M 2008.38.1.
Bartonian, Szőc Limestone Formation
Dorog
Roz24a, p. 24 etc.
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotypes, M.62.6247, M.62.6289,
M.62.6290, M.62.6295, M.62.6326
Bartonian, Szőc Limestone Formation
Reimann shaft, Csolnok
Roz24a, p. 24 etc.
Form B
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotypes, M.62.6308
Bartonian, Szőc Limestone Formation
Csolnok
Roz24a, p. 24 etc.
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotypes, M.62.9403
Bartonian, Szőc Limestone Formation
Dorog
Roz24a, p. 24 etc.
Types missing
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotype, M.62.9414
Bartonian, Szőc Limestone Formation
Solymár
Roz24a, p. 24 etc.
- Nummulina subplanulata* ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
paralectotype, M.62.9415
Bartonian, Szőc Limestone Formation
Sárisáp
Roz24a, p. 24 etc.

Nummulina subplanulata ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 paralectotypes, M 2008.39.1., M 2008.40.1.,
 M 2008.41.1.
 Bartonian, Szőc Limestone Formation
 Nagykovácsi
 Roz24a, p. 24 etc.

Nummulina subplanulata ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 paralectotypes, M 2008.42.1.
 Bartonian, Szőc Limestone Formation
 Piszke, Lábatlan
 Roz24a, p. 24 etc.
 Form A

Nummulina subplanulata ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 paralectotypes, M 2008.43.1., M 2008.44.1.
 Bartonian, Szőc Limestone Formation
 Piszke, Lábatlan
 Roz24a, p. 24 etc.

Nummulina subplanulata ROZLOZSNIK
ex HANTKEN & MADARÁSZ, 1924
 paralectotypes, M 2008.45.1.
 Bartonian, Szőc Limestone Formation
 Sárisáp
 Roz24a, p. 24 etc.
 Form A

Nummulites anomaloides KECSKEMÉTI, 1974
 holotype, M.73.41
 Bartonian, Szőc Limestone Formation
 borehole Pgvt-31, Pénzesgyőr
 Kec74, p. 41, pl. 1, fig. 4
 Form A. Inventory number given erroneously
 in original publication as M.73.42

Nummulites anomaloides KECSKEMÉTI, 1974
 paratype, M.73.42
 Bartonian, Szőc Limestone Formation
 borehole Pgvt-31, Pénzesgyőr
 Kec74, p. 41, pl. 1, fig. 3
 Form A. Original publication erroneously
 quotes this inventory number for holotype

Nummulites anomaloides KECSKEMÉTI, 1974
 paratype, M.73.129
 Bartonian, Szőc Limestone Formation
 borehole Bcs-9, Bakonycsernye
 Kec74, p. 41
 Form A

Nummulites bombitus HOTTINGER, 1977
 holotype, M 2008.3.1.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129, pl. 61, fig. 1 (text-figs. 52a-b,
 pl. 61, figs. 2-4)
 Form B

Nummulites bombitus HOTTINGER, 1977
 paratype, M 2008.4.1.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129, text-fig. 54a
 Form A

Nummulites bombitus HOTTINGER, 1977
 paratype, M 2008.4.2.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129, text-fig. 54b
 Form A

Nummulites bombitus HOTTINGER, 1977
 paratype, M 2008.4.3.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129, text-fig. 54c
 Form A

Nummulites bombitus HOTTINGER, 1977
 paratype, M 2008.4.4.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129, text-fig. 54d
 Form B

Nummulites bombitus HOTTINGER, 1977
 paratypes, M 2008.5.1-3.
 Ypresian
 Dikilitas, Varna, Bulgaria
 Hot77, p. 129

Nummulites dudarensis KECSKEMÉTI, 1970
holotype, M.69.508
Lutetian, Szőc Limestone Formation
Weimpuszta, Nagyesztergár
Kec70, p. 62, pl. 1, figs. 11, 12
Form A

Nummulites dudarensis KECSKEMÉTI, 1970
paratype, M.69.509
Lutetian, Szőc Limestone Formation
Weimpuszta, Nagyesztergár
Kec70, p. 62, pl. 1, fig. 13
Form A

Nummulites kopeki KECSKEMÉTI, 1974
holotype, M.73.29
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 5
Form A

Nummulites kopeki KECSKEMÉTI, 1974
paratype, M.73.31
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 6
Form B

Nummulites kopeki KECSKEMÉTI, 1974
paratypes, M.73.130, M.73.131, M.73.132,
M.73.133, M.73.134
Bartonian, Szőc Limestone Formation
borehole Ba-219, Balinka
Kec74, p. 42
Form A

Nummulites kopeki KECSKEMÉTI, 1974
paratypes, M.73.136, M.73.137
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42
Form A

Nummulites kopeki KECSKEMÉTI, 1974
paratype, M.73.140
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 10
Form A

Nummulites kopeki KECSKEMÉTI, 1974
paratype, M.73.141
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 9
Form B

Nummulites kopeki KECSKEMÉTI, 1974
paratype, M.73.142
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 8
Form A

Nummulites kopeki KECSKEMÉTI, 1974
paratype, M.73.143
Bartonian, Szőc Limestone Formation
borehole D-218, Dudar
Kec74, p. 42, pl. 1, fig. 7
Form A

Nummulites madarászi HANTKEN, 1875
syntypes, M 2008.14.1.
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han75a (Han75b), p. 86 (75) (pl. 16,
figs. 7a–c)

Nummulites majzoni KECSKEMÉTI, 1970
holotype, M.69.506
Bartonian, Szőc Limestone Formation
Ree-hegy, Pénzesgyőr
Kec70, p. 60, pl. 1, figs. 7, 8
Form A

Nummulites penzesgyoerensis KECSKEMÉTI,
1970
holotype, M.69.504
Bartonian, Szőc Limestone Formation
Ree-hegy, Pénzesgyőr
Kec70, p. 57, pl. 1, figs. 2, 4
Form A

Nummulites penzesgyoerensis KECSKEMÉTI,
1970
paratype, M.69.503
Bartonian, Szőc Limestone Formation
Ree-hegy, Pénzesgyőr
Kec70, p. 57, pl. 1, figs. 1, 3
Form B

- Nummulites schaubi* KECSKEMÉTI, 1970
holotype, M.69.505
Bartonian, Szőc Limestone Formation
Ree-hegy, Pénzesgyőr
Kec70, p. 59, pl. 1, figs. 5, 6
Form A
- Nummulites subtilis maior* KECSKEMÉTI, 1972
holotype, M.71.176
Bartonian, Szőc Limestone Formation
borehole Bj-30, Bajót
Kec72, p. 127, pl. 28, fig. 13, pl. 29, fig. 1
Form A
- Nummulites subtilis maior* KECSKEMÉTI, 1972
paratypes, M.71.174
Bartonian, Szőc Limestone Formation
borehole Cs-692, Csolnok
Kec72, p. 127
Form A
- Nummulites subtilis maior* KECSKEMÉTI, 1972
paratype, M.71.175
Bartonian, Szőc Limestone Formation
borehole Cs-692, Csolnok
Kec72, p. 127, pl. 29, fig. 2
Form A
- Nummulites suemegensis* KECSKEMÉTI, 1974
holotype, M.73.4
Lutetian, Szőc Limestone Formation
Darvas-tó No. VI., Sümeg
Kec74, p. 39, pl. 1, figs. 1, 2
Form A
- Nummulites zircensis* KECSKEMÉTI, 1970
holotype, M.69.507
Bartonian, Szőc Limestone Formation
Ree-hegy, Pénzesgyőr
Kec70, p. 61, pl. 1, figs. 9, 10
Form A
- Pyrgo microstriatus* OZSVÁRT, 2003
holotype, 2004.50.1.
Bartonian, Csolnok Clay Marl Formation
borehole Csbr-89, Csákberény
Ozs03, p. 2, pl. 1, figs. 3–4
Inventory number in original publication:
OP-4300, Collection of Dept. of Paleontology,
Eötvös University
- Robulina granulata* HANTKEN, 1875
paralectotype, M.99.64
Bartonian, Padrag Marl Formation
Porva
Han75a (Han75b), p. 57 (49), pl. 14, fig. 15
Lenticulina granulata, Dar99; Hor03
Type figured in HORVÁTH 2003. Type lost
- Textilaria subflabelliformis* HANTKEN, 1875
paralectotype, M.99.42
Priabonian, Buda Marl Formation
Kis-Sváb-hegy, Budapest
Han75a (Han75b), p. 66 (57), pl. 15, fig. 2
Vulvulina subflabelliformis, Hag60; Hor02
Type figured in HORVÁTH 2002
- Textularia tuberosus* OZSVÁRT, 2003
holotype, 2004.49.1.
Priabonian, Buda Marl Formation
Mátyás-hegy, western quarry, Budapest
Ozs03, p. 2, pl. 1, figs. 1–2
Inventory number in original publication:
OP-4448, Collection of Dept. of Paleontology,
Eötvös University
- Triloculina porvaensis* HANTKEN, 1875
neotype, M.99.49
Bartonian, Csolnok Clay Marl Formation
borehole Csordakút-2, Nagyegyháza, Óbarok
Han75a (Han75b), p. 21 (76), pl. 13, fig. 3
Neotype designated in HORVÁTH 2002
- Vaginulinopsis hagni* OZSVÁRT, 2007
holotype, M 2008.118.1.
Bartonian, Padrag Marl Formation
borehole Dv-4, Devecser
Ozs07, p. 55, pl. 5, fig. 26
- Vaginulinopsis hagni* OZSVÁRT, 2007
paratype, M 2008.118.2.
Bartonian, Padrag Marl Formation
borehole Dv-4, Devecser
Ozs07, p. 55, pl. 5, fig. 25
- Vaginulinopsis hagni* OZSVÁRT, 2007
paratype, M 2008.118.3.
Bartonian, Padrag Marl Formation
borehole Dv-4, Devecser
Ozs07, p. 55, pl. 5, fig. 27

Vulvulina pectinata HANTKEN, 1875
 paralectotype, M.99.41
 Priabonian, Buda Marl Formation
 Kis-Sváb-hegy, Budapest
Han75a (*Han75b*), p. 68 (58), pl. 7, fig. 10
 Type figured in HORVÁTH 2002

Cyathophyllia hantkeni REUSS, 1870
 syntype
 Lutetian–Bartonian, Csernye Formation?
 Piszke, Lábatlan
Res70, p. 48, pl. 4, fig. 1
 Type probably destroyed in 1956

6.2. Eocene Anthozoa

Archicoenopsammia hungarica KOLOSVÁRY,
 1949
 syntype, M.57.2291
 Lutetian?, Csernye Formation?
 Felsőgalla, Tatabánya
Kol49b, p. 229 (192)

Flabellum rotundum KOLOSVÁRY, 1949
 holotype, M.57.2516
 Lutetian–Bartonian, Csernye Formation?
 Mogyorós, Mogyorósbánya
Kol49b, p. 229 (193), pl. 21, fig. c

Archicoenopsammia hungarica KOLOSVÁRY,
 1949
 syntype, M.62.663
 Lutetian?, Csernye Formation?
 Felsőgalla, Tatabánya
Kol49b, p. 229 (192), pl. 21, fig. b
 Type missing

Flabellum szőtsi KOLOSVÁRY, 1949
 holotype, M.57.2515
 Lutetian–Bartonian, Csernye Formation?
 Bajót
Kol49b, p. 229 (193), pl. 21, fig. d

Calamophyllia curvicostata KOLOSVÁRY, 1949
 syntypes, M.57.2336
 Lutetian, Csernye Formation?
 below Nagy-kő, Ótokod, Tokod
Kol49b, p. 227 (190), pl. 20, fig. 2a

Rhabdophyllia budense KOLOSVÁRY, 1949
 syntype, M.57.2716
 Priabonian?, Szépvölgy Limestone Formation?
 Zugliget, Budapest
Kol49b, p. 228 (190), pl. 20, fig. 2c

Calamophyllia curvicostata KOLOSVÁRY, 1949
 syntype, M.57.2343
 Lutetian–Bartonian, Csernye Formation?
 Pusztavám
Kol49b, p. 227 (190), pl. 20, fig. 2b

Rhabdophyllia budense KOLOSVÁRY, 1949
 syntype, M.57.2717
 Priabonian?, Szépvölgy Limestone Formation?
 Endrődy utca 61., Budapest
Kol49b, p. 228 (190)

Calamophyllia curvicostata KOLOSVÁRY, 1949
 syntype, M.57.2344
 Lutetian–Bartonian, Csernye Formation?
 Pusztavám
Kol49b, p. 227 (190)

Stephanosmilia dendricola KOLOSVÁRY, 1949
 syntype, M.57.2710
 Lutetian–Bartonian, Csernye Formation?
 Hosszúharasztos, Csákvár
Kol49b, p. 231 (195), pl. 22, figs. h–k
 Type missing

Cyathophyllia hantkeni REUSS, 1870
 syntypes
 Lutetian–Bartonian, Csernye Formation?
 Mogyorósbánya
Res70, p. 48, pl. 4, fig. 1
 Types probably destroyed in 1956

Stephanosmilia vadászi KOLOSVÁRY, 1949
 syntype, M.57.2711
 Lutetian–Bartonian, Csernye Formation?
 “new outcrop”, Csákvár
Kol49b, p. 230 (195), pl. 22, figs. b–g
 Type missing

Stylacropora hungarica KOLOSVÁRY, 1949
 syntypes
 Lutetian?, Csernye Formation?
 Felsőgalla, Tatabánya
Kol49b, p. 231 (197), pl. 22, fig. 1
 Types probably destroyed in 1956

- Trochocyathus longus* REUSS, 1870
 syntype
 Lutetian–Bartonian, Csernye Formation?
 Mogyorósbánya
Res70, p. 42, pl. 2, figs. 2, 3
 Type probably destroyed in 1956
- Trochosmilia 4-cingulata* KOLOSVÁRY, 1949
 syntype, M.57.2898
 Lutetian–Bartonian, Csernye Formation?
 Bakonybél
Kol49b, p. 230 (194), pl. 21, fig. i, pl. 22, fig. a
- Trochosmilia aequalis* REUSS, 1870
 syntypes
 Lutetian–Bartonian, Csernye Formation?
 Mogyorósbánya
Res70, p. 45, pl. 3, figs. 3–5
 Types probably destroyed in 1956
- Trochosmilia brachypoda* REUSS, 1870
 holotype, M.57.2899
 Lutetian–Bartonian, Csernye Formation?
 Piszke, Lábatlan
Res70, p. 46, pl. 3, fig. 6
 Locality on specimen label: Bajót
- ?*Turbinoseris noszkyi* KOLOSVÁRY, 1949
 holotype
 Lutetian–Bartonian, Csernye Formation?
 Sűrű-hegy, Nagyesztergár
Kol49b, p. 228 (191), pl. 20, fig. 2e
 Type probably destroyed in 1956
- ?*Turbinoseris vadászi* KOLOSVÁRY, 1949
 syntypes, M.57.2899
 Lutetian?, Csernye Formation?
 Csákánygödör, Felsőgalla, Tatabánya
Kol49b, p. 228 (191), pl. 21, fig. a
- 6.3. Eocene Bivalvia**
- Abra pannonica* SZŐTS, 1953
 syntypes, M.59.7642
 Bartonian, Forna Formation
 Gánti szőlök, Zámoly
Sz653, p. 219 (95) (pl. 10, figs. 45–48)
- Abra pannonica* SZŐTS, 1953
 syntypes, M.59.7643
 Bartonian, Forna Formation
 “new outcrop”, Csákvár
Sz653, p. 219 (95)
- Arca caillatiformis* SZŐTS, 1953
 syntypes, M.59.7585
 Bartonian, Forna Formation
 Gánti szőlök, Zámoly
Sz653, p. 207 (84) (pl. 8, figs. 19–21)
- Arca vértesensis* SZŐTS, 1953
 syntypes, M.59.7575, M.59.7577,
 M.59.7579, M.59.7581
 Bartonian, Forna Formation
 Gánti szőlök, Zámoly
Sz653, p. 206 (84) (pl. 8, figs. 11–18)
- Arca vértesensis* SZŐTS, 1953
 syntypes, M.59.7576
 Bartonian, Forna Formation
 “new outcrop”, Csákvár
Sz653, p. 206 (84)
- Arca vértesensis* SZŐTS, 1953
 syntypes, M.59.7578
 Bartonian, Forna Formation
 Szőlőhegy, Csákberény
Sz653, p. 206 (84)
- Arca vértesensis* SZŐTS, 1953
 syntypes, M.59.7580, M.59.7582
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
Sz653, p. 206 (84)
- Arcopagia majeri* SZŐTS, 1953
 syntype, M.59.7639
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
Sz653, p. 219 (94) (pl. 10, figs. 41–44)
- Arcopagia majeri* SZŐTS, 1953
 syntypes, M.59.7640
 Bartonian, Forna Formation
 “new outcrop”, Csákvár
Sz653, p. 219 (94)

- Arcopagia majeri* SZŐTS, 1953
 syntypes, M.59.7641
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 219 (94)
- Beguina taegeri* SZŐTS, 1953
 syntypes, M.59.7606
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 212 (88) (pl. 8, figs. 37–42)
- Corculum subrotundatum* SZŐTS, 1953
 syntypes, M.59.7624, M.59.7629
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 217 (93) (pl. 10, figs. 22–27)
- Corculum subrotundatum* SZŐTS, 1953
 syntypes, M.59.7627, M.59.7628
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 217 (93)
- Corculum subrotundatum* SZŐTS, 1953
 syntypes, M.59.7630
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 217 (93)
 Specimens from this locality reported in publication as lost
- Lucina vogli* SZŐTS, 1953
 syntypes, M.59.7621
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 216 (92) (pl. 10, figs. 10–17)
- Lucina vogli* SZŐTS, 1953
 syntypes, M.59.7622
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 216 (92)
- Pholas (Martesia) pappi* SZŐTS, 1938
 holotype, M.77.81
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 31, pl. 1, figs. 6a–c
- Trinacria gántensis* SZŐTS, 1953
 syntype, M.59.7571
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 204 (82) (pl. 8, figs. 7–8)
- Trinacria gántensis* SZŐTS, 1953
 syntype, M.59.7572
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 204 (82)
- Trinacria mórensis* SZŐTS, 1953
 syntypes, M.59.7567
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 204 (82) (pl. 8, figs. 1–6)
- Trinacria mórensis* SZŐTS, 1953
 syntypes, M.59.7568, M.59.7569
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 204 (82)
- Trinacria mórensis* SZŐTS, 1953
 syntypes, M.59.7570
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 204 (82)
- 6.4. Eocene Scaphopoda**
- Cadulus pseudohungaricus* SZŐTS, 1953
 syntypes, M.61.1536
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 203 (81) (pl. 7, figs. 53–54)
- Cadulus pseudohungaricus* SZŐTS, 1953
 syntypes, M.61.1537
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 203 (81)
- Cadulus pseudohungaricus* SZŐTS, 1953
 syntypes, M.61.1577
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 203 (81)

6.5. Eocene Gastropoda

Acera aspirata SZŐTS, 1953
syntypes, M.59.7543
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 200 (77) (pl. 7, fig. 38)

Acera aspirata SZŐTS, 1953
syntypes, M.59.7544
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 200 (77)

Actaeon vitálisi SZŐTS, 1953
syntype, M.59.7538
Bartonian, Forna Formation
Gánti szőlők, Zámoly
Szö53, p. 199 (76) (pl. 7, fig. 35)
Specimen from this locality not mentioned
in publication

Actaeon vitálisi SZŐTS, 1953
syntype, M.59.7539
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 199 (76)

Anisus bicarinatus SZŐTS, 1953
syntypes, M.59.7557
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 202 (79) (pl. 7, figs. 47–49)

Anisus pseudosubangulatus SZŐTS, 1953
syntypes, M.59.7555
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 202 (79) (pl. 7, figs. 44–46)

Anisus pseudosubangulatus SZŐTS, 1953
syntypes, M.59.7556
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 202 (79)

Assiminea gránásensis SZŐTS, 1953
syntypes, M.59.7152
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 151 (38) (pl. 2, fig. 32)

Assiminea gránásensis SZŐTS, 1953
syntypes, M.59.7153
Bartonian, Forna Formation
Gánti szőlők, Zámoly
Szö53, p. 151 (38)

Assiminea gránásensis SZŐTS, 1953
syntypes, M.59.7154
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 151 (38)

Assiminea quadrangulata SZŐTS, 1953
syntypes, M.59.7155
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 151 (39) (pl. 2, fig. 33)

Assiminea quadrangulata SZŐTS, 1953
syntypes, M.59.7157
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 151 (39)

Asthenotoma graniformis SZŐTS, 1953
syntypes, M.59.7525
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 195 (73) (pl. 7, fig. 23)

Asthenotoma graniformis SZŐTS, 1953
syntypes, M.59.7526
Bartonian, Forna Formation
Hosszúharasztos, Csákvár
Szö53, p. 195 (73)

Asthenotoma graniformis SZŐTS, 1953
syntypes, M.59.7527
Bartonian, Forna Formation
Gánti szőlők, Zámoly
Szö53, p. 195 (73)

Bayania boussaci SZŐTS, 1953
syntypes, M.59.7204
Bartonian, Forna Formation
“new outcrop”, Csákvár
Szö53, p. 160 (46) (pl. 3, figs. 1–2)

- Bayania boussaci* SZŐTS, 1953
 syntypes, M.59.7205
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 160 (46)
- Bayania boussaci* SZŐTS, 1953
 syntypes, M.59.7206
 Bartonian, Forna Formation
 Csákberény
 Sző53, p. 160 (46)
- Bayania supravarians* SZŐTS, 1953
 syntypes, M.59.7198
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 160 (46) (pl. 2, figs. 67–68)
- Bayania supravarians* SZŐTS, 1953
 syntypes, M.59.7202
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 160 (46)
- Bayania supravarians* SZŐTS, 1953
 syntypes, M.59.7203
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 160 (46)
- Bayania variocostata* SZŐTS, 1953
 syntypes, M.59.7195
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 160 (45) (pl. 2, fig. 66)
- Bayania variocostata* SZŐTS, 1953
 syntypes, M.59.7196
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 160 (45)
- Bayania variocostata* SZŐTS, 1953
 syntypes, M.59.7197
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 160 (45)
- Bythinella auriculata* SZŐTS, 1953
 syntypes, M.59.7135
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 148 (35) (pl. 2, figs. 23–24)
- Bythinella gracillima* SZŐTS, 1953
 syntypes, M.59.7133
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 148 (35) (pl. 2, fig. 22)
- Bythinella gracillima* SZŐTS, 1953
 syntypes, M.59.7134
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 148 (35)
- Bythinella gracillima* SZŐTS, 1953
 syntypes, M.59.7136
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 148 (35)
- Bythinella pulcherrima* SZŐTS, 1953
 syntypes, M.59.7130
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 147 (35) (pl. 2, fig. 21)
- Bythinella pulcherrima* SZŐTS, 1953
 syntypes, M.59.7131
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 147 (35)
- Bythinella pulcherrima* SZŐTS, 1953
 syntypes, M.59.7132
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 147 (35)
- Cantharus vértesensis* SZŐTS, 1953
 syntypes, M.59.7460
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 183 (63) (pl. 6, figs. 12–13)

- Cantharus vértesensis* SZŐTS, 1953
 syntypes, M.59.7461
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 183 (63)
- Cantharus vértesensis* SZŐTS, 1953
 syntypes, M.59.7462
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 183 (63)
- Cerithiella rara* SZŐTS, 1953
 syntypes, M.59.7293
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 169 (51) (pl. 4, fig. 22)
- Cerithiella rara* SZŐTS, 1953
 syntypes, M.59.7294
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 169 (51)
- Cerithiella rara* SZŐTS, 1953
 syntypes, M.59.7295
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 169 (51)
- Cerithium calcaratum montis-antonii* SZŐTS, 1938
 syntypes, M.74.270
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 32, pl. 1, fig. 11
- Cingula pseudomumiola* SZŐTS, 1953
 syntypes, M.59.7137
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 149 (36) (pl. 2, fig. 26)
- Cingula pseudomumiola* SZŐTS, 1953
 syntypes, M.59.7138
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 149 (36)
- Collonia vértesensis* SZŐTS, 1953
 syntypes, M.59.7112
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 139 (27) (pl. 1, figs. 34–36)
- Collonia vértesensis* SZŐTS, 1953
 syntypes, M.59.7113
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 139 (27)
- Collonia vértesensis* SZŐTS, 1953
 syntypes, M.59.7114
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 139 (27)
- Cyclostrema csákvárensis* SZŐTS, 1953
 syntypes, M.59.7111
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 138 (27) (pl. 1, figs. 31–33)
- Cylichna gántensis* SZŐTS, 1953
 syntypes, M.59.7545
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 200 (78) (pl. 7, fig. 39)
- Cylichna gántensis* SZŐTS, 1953
 syntypes, M.59.7546
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 200 (78)
- Cylichna gántensis* SZŐTS, 1953
 syntypes, M.59.7547, M.59.7548
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 200 (78)

Cylichna hantkeni SZŐTS, 1953
 syntypes, M.59.7552
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 201 (78) (pl. 7, fig. 41)

Cylichna hantkeni SZŐTS, 1953
 syntypes, M.59.7553
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 201 (78)

Cylichna vértesensis SZŐTS, 1953
 syntypes, M.59.7549
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 200 (78) (pl. 7, fig. 40)

Cylichna vértesensis SZŐTS, 1953
 syntypes, M.59.7550
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 200 (78)

Cylichna vértesensis SZŐTS, 1953
 syntypes, M.59.7551
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 200 (78)

Cythara vértesensis SZŐTS, 1953
 syntypes, M.59.7528
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 195 (73) (pl. 7, fig. 24)

Cythara vértesensis SZŐTS, 1953
 syntypes, M.59.7529
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 195 (73)

Cythara vértesensis SZŐTS, 1953
 syntypes, M.59.7530
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 195 (73)

Deshayesia naticoides SZŐTS, 1938
 syntypes, M.74.229
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 33, pl. 1, fig. 9

Eulimella guttulina SZŐTS, 1953
 syntypes, M.59.7319
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 174 (56) (pl. 4, fig. 39)

Eulimella guttulina SZŐTS, 1953
 syntypes, M.59.7320
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 174 (56)

Eulimella guttulina SZŐTS, 1953
 syntypes, M.59.7321
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 174 (56)

Lacuna umbonata SZŐTS, 1953
 syntype, M.59.7128
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 146 (34) (pl. 2, fig. 18)

Lacuna umbonata SZŐTS, 1953
 syntypes, M.59.7129
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 146 (34)

Marginella crassula humilispira SZŐTS, 1938
 holotype, M.62.1834
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 34, pl. 1, fig. 12

Marginella frequens SZŐTS, 1953
 syntypes, M.59.7510
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 193 (71) (pl. 7, fig. 20)

- Marginella frequens* SZŐTS, 1953
 syntypes, M.59.7511
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 193 (71)
- Marginella frequens* SZŐTS, 1953
 syntypes, M.59.7512
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 193 (71)
- Marginella hantkeni* SZŐTS, 1953
 syntypes, M.59.7504
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 191 (70) (pl. 7, fig. 17)
- Marginella hantkeni* SZŐTS, 1953
 syntypes, M.59.7505
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 191 (70)
- Marginella hantkeni* SZŐTS, 1953
 syntypes, M.59.7506
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 191 (70)
- Marginella pannonica* SZŐTS, 1953
 syntypes, M.59.7513, M.59.7516
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 193 (72) (pl. 7, fig. 21)
- Marginella pannonica* SZŐTS, 1953
 syntypes, M.59.7514
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 193 (72)
- Marginella pannonica* SZŐTS, 1953
 syntypes, M.59.7515
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 193 (72)
- Marginella pseudonana* SZŐTS, 1953
 syntypes, M.59.7495
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 190 (69) (pl. 7, fig. 14)
- Marginella pseudonana* SZŐTS, 1953
 syntypes, M.59.7496
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 190 (69)
- Marginella pseudonana* SZŐTS, 1953
 syntypes, M.59.7497
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 190 (69)
- Marginella subcylindrica* SZŐTS, 1953
 syntypes, M.59.7507
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 192 (71) (pl. 7, figs. 18–19)
- Marginella subcylindrica* SZŐTS, 1953
 syntypes, M.59.7508
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 192 (71)
- Marginella subcylindrica* SZŐTS, 1953
 syntypes, M.59.7509
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 192 (71)
- Marginella vértesensis* SZŐTS, 1953
 syntypes, M.59.7517
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 194 (72) (pl. 7, fig. 22)
- Marginella vértesensis* SZŐTS, 1953
 syntype, M.59.7519
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 194 (72)

Marginella vértesensis SZŐTS, 1953
 syntypes, M.59.7520
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 194 (72)

Mathilda frequens SZŐTS, 1953
 syntypes, M.59.7167
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 153 (41) (pl. 2, figs. 41–42)

Mathilda frequens SZŐTS, 1953
 syntypes, M.59.7168
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 153 (41)

Mathilda frequens SZŐTS, 1953
 syntypes, M.59.7169
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 153 (41)

Mitra subcrebricosta SZŐTS, 1953
 syntypes, M.59.7488
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 188 (67) (pl. 7, fig. 9)

Murex gántensis SZŐTS, 1953
 syntypes, M.59.7443
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 181 (62) (pl. 6, figs. 3–5)

Murex gántensis SZŐTS, 1953
 syntypes, M.59.7444
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 181 (62)

Natica boggschi SZŐTS, 1938
 syntypes, M.62.1826
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 33, pl. 1, fig. 8

Natica gránásensis SZŐTS, 1953
 syntypes, M.59.7428, M.59.7429
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 179 (60) (pl. 5, figs. 14–15)

Natica gránásensis SZŐTS, 1953
 syntypes, M.59.7430
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 179 (60)

Natica gránásensis SZŐTS, 1953
 syntypes, M.59.7435
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 179 (60)

Nerita hantkeni SZŐTS, 1953
 syntypes, M.59.7122
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 142 (30) (pl. 2, figs. 6–7)

Nerita hantkeni SZŐTS, 1953
 syntypes, M.59.7123
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 142 (30)

Nerita héberti SZŐTS, 1953
 syntypes, M.59.7118, M.59.7119, M.59.7121
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 141 (30) (pl. 2, figs. 3–5)

Nerita héberti SZŐTS, 1953
 syntypes, M.59.7120
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 141 (30)

Odostomia pannonica SZŐTS, 1953
 syntypes, M.59.7309
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 172 (54) (pl. 4, fig. 31)

- Odostomia pannonica* SZŐTS, 1953
 syntypes, M.59.7310
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 172 (54)
- Odostomia pannonica* SZŐTS, 1953
 syntypes, M.59.7311
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 172 (54)
- Odostomia pseudoruellensis* SZŐTS, 1953
 syntypes, M.59.7313
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 172 (55) (pl. 4, fig. 33)
- Odostomia pseudoruellensis* SZŐTS, 1953
 syntypes, M.59.7314
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 172 (55)
- Odostomia pseudoruellensis* SZŐTS, 1953
 syntypes, M.59.7315
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 172 (55)
- Odostomia semistriata* SZŐTS, 1953
 syntypes, M.59.7312
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 172 (54) (pl. 4, fig. 32)
- Odostomia submisera* SZŐTS, 1953
 syntypes, M.59.7316
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 173 (55) (pl. 4, figs. 34–36)
- Odostomia submisera* SZŐTS, 1953
 syntypes, M.62.6903
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 173 (55)
- Odostomia supravariabilis* SZŐTS, 1953
 syntypes, M.59.7317
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 173 (55) (pl. 4, figs. 37–38)
- Odostomia supravariabilis* SZŐTS, 1953
 syntypes, M.59.7318
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 173 (55)
- Parvisipho nudus* SZŐTS, 1953
 syntypes, M.59.7463
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 184 (64) (pl. 6, fig. 15)
- Parvisipho nudus* SZŐTS, 1953
 syntypes, M.59.7464
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 184 (64)
- Parvisipho nudus* SZŐTS, 1953
 syntypes, M.59.7465
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 184 (64)
- Persicula pseudoallixi* SZŐTS, 1953
 syntypes, M.59.7489
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 189 (68) (pl. 7, fig. 12)
- Persicula pseudoallixi* SZŐTS, 1953
 syntypes, M.59.7490
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 189 (68)
- Persicula pseudoallixi* SZŐTS, 1953
 syntypes, M.59.7491
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 189 (68)

Rissoa munieri SZŐTS, 1953
 syntypes, M.59.7140
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 149 (37) (pl. 2, fig. 27)

Rissoa munieri SZŐTS, 1953
 syntypes, M.59.7141
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 149 (37)

Rissoa munieri SZŐTS, 1953
 syntypes, M.59.7142
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 149 (37)

Scissurella hungarica SZŐTS, 1953
 syntypes, M.59.7102
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 133 (22) (pl. 1, figs. 5-7)

Scissurella hungarica SZŐTS, 1953
 syntypes, M.59.7103, M.60.1
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 133 (22)

Seila quadricostata SZŐTS, 1953
 syntypes, M.59.7300
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 170 (52) (pl. 4, figs. 25-26)

Seila quadricostata SZŐTS, 1953
 syntypes, M.59.7301
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 170 (52)

Seila quadricostata SZŐTS, 1953
 syntypes, M.59.7302
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 170 (52)

Seila subtrifaria SZŐTS, 1953
 syntypes, M.59.7298
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 170 (52) (pl. 4, fig. 24)
 Specimens from this locality not mentioned in publication

Seila subtrifaria SZŐTS, 1953
 syntypes, M.59.7299
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 170 (52)

Seila vértesensis SZŐTS, 1953
 syntypes, M.59.7296
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 170 (52) (pl. 4, fig. 23)

Seila vértesensis SZŐTS, 1953
 syntypes, M.59.7297
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 170 (52)

Solariella gántensis SZŐTS, 1953
 syntype, M.59.7105
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 137 (26) (pl. 1, figs. 21-23)
 Single specimen reported in publication.
 The photographed syntype is in GIH collection (E.11.), see BODA 1964

Terebellum vértesensis SZŐTS, 1953
 syntypes, M.59.7325
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 175 (57) (pl. 4, figs. 43-44)

Terebellum vértesensis SZŐTS, 1953
 syntypes, M.59.7326
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 175 (57)

Terebellum vértesensis SZŐTS, 1953
 syntypes, M.59.7327
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 175 (57)

Tricolia colorata SZŐTS, 1953
 syntypes, M.59.7115
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 140 (29) (pl. 1, fig. 39)

Triphora tricostata SZŐTS, 1953
 syntypes, M.59.7303
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 171 (53) (pl. 4, figs. 27–28)

Triphora tricostata SZŐTS, 1953
 syntypes, M.59.7304
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 171 (53)

Triphora tricostata SZŐTS, 1953
 syntypes, M.59.7305
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 171 (53)

Turbanilla conica SZŐTS, 1953
 syntypes, M.59.7323
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 175 (57) (pl. 4, fig. 41)

Turritella (Mesalia) berenderi SZŐTS, 1938
 syntypes, M.62.1807
 Lutetian, Dorog Formation
 Antal-hegy, Mór
 Sző38, p. 33, pl. 1, fig. 10

Turritella rómeri SZŐTS, 1953
 syntypes, M.59.7163
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 153 (40) (pl. 2, figs. 38–39)

Turritella rómeri SZŐTS, 1953
 syntypes, M.59.7164
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 153 (40)

Turritella rómeri SZŐTS, 1953
 syntypes, M.59.7165, M.59.7166
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 153 (40)

Tymanotonus rozloznskii SZŐTS, 1953
 syntypes, M.59.7282
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 162 (48) (pl. 3, figs. 19–20)

Tymanotonus rozloznskii SZŐTS, 1953
 syntypes, M.59.7283, M.59.7284
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 162 (48)

Vermetus ornatissimus SZŐTS, 1953
 syntypes, M.59.7171
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 154 (42) (pl. 2, figs. 46–47)

Vexillum böckhi SZŐTS, 1953
 syntypes, M.59.7484
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 187 (66) (pl. 7, figs. 5–6)

Vexillum böckhi SZŐTS, 1953
 syntypes, M.59.7485
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Sző53, p. 187 (66)

Vexillum csákvárense SZŐTS, 1953
 syntypes, M.59.7482
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Sző53, p. 187 (66) (pl. 7, fig. 4)

Vexillum csákvárense SZŐTS, 1953
 syntypes, M.59.7483
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Sző53, p. 187 (66)

Vexillum splendidum SZŐTS, 1953
 syntypes, M.59.7486
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Szö53, p. 188 (67) (pl. 7, figs. 7–8)

Zebina hungarica SZŐTS, 1953
 syntypes, M.59.7143, M.59.7144
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Szö53, p. 149 (37) (pl. 2, fig. 28)

Zebina hungarica SZŐTS, 1953
 syntypes, M.59.7145
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Szö53, p. 149 (37)

Zebina hungarica SZŐTS, 1953
 syntypes, M.59.7146
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Szö53, p. 149 (37)

Zebina zitteli SZŐTS, 1953
 syntypes, M.59.7147
 Bartonian, Forna Formation
 Gánti szőlők, Zámoly
 Szö53, p. 150 (37) (pl. 2, fig. 29)

Zebina zitteli SZŐTS, 1953
 syntypes, M.59.7148
 Bartonian, Forna Formation
 Hosszúharasztos, Csákvár
 Szö53, p. 150 (37)

Zebina zitteli SZŐTS, 1953
 syntypes, M.59.7149
 Bartonian, Forna Formation
 "new outcrop", Csákvár
 Szö53, p. 150 (37)

6.6. Eocene Cephalopoda

Archaeosepia naefi SZÖRÉNYI, 1933
 syntype, M.61.3120
 Bartonian, Csolnok Clay Marl Formation
 Tatabánya
 Szö33, p. 186, pl. 6, fig. 1 (pl. 6, fig. 2)

Sepia agriensis WAGNER, 1938
 holotype, M.63.6909
 Priabonian, Buda Marl Formation
 Eged-hegy, Eger
 Wag38, p. 196 (187), text-fig. 3 (left side)
 Type missing

Sepia agriensis WAGNER, 1938
 paratypes
 Priabonian, Buda Marl Formation
 Eged-hegy, Eger
 Wag38, p. 196 (187), text-fig. 3 (right side)
 Types missing, not found in inventory

6.7. Eocene Brachiopoda

Argyrotheca tokodensis BITNER & DULAI, 2008
 holotype, 2006.37.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-figs. 7.6–9.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.35.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 7.1.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.36.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 7.12.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.38.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-figs. 7.2–3.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.39.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-figs. 7.4–5.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.40.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-figs. 7.10–11.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.42.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.1.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.43.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.2.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.44.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.3.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.45.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.4.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.46.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-figs. 8.5–6.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.47.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.7.

Argyrotheca tokodensis BITNER & DULAI, 2008
 paratype, 2006.48.1.
 Bartonian, Padrag Marl Formation
 Tokod
 Bit08, p. 37, text-fig. 8.8.

6.8. Eocene Cirripedia

Balanus hantkeni KOLOSVÁRY, 1947
 syntype, M.57.3492
 Lutetian–Bartonian, Csernye Formation?
 quarry, Nagykovácsi
 Kol47, p. 307, text-fig. 1
 Type missing

6.9. Eocene Decapoda

Actaeites lobatus MÜLLER & COLLINS, 1991
 holotype, M.91.153
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 70, text-fig. 4c, pl. 4, figs. 9, 10

Branchioplax sulcata MÜLLER & COLLINS, 1991
 holotype, M.91.199
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 82, text-fig. 5a, pl. 6, fig. 14
 Paratypes under same inventory number
 (pl. 6, figs. 16, 18)

Branchioplax sulcata MÜLLER & COLLINS, 1991
 paratypes, M.91.200
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 82, text-fig. 5a

Branchioplax sulcata MÜLLER & COLLINS, 1991
 paratypes, M.91.201, M.91.202, M.91.203
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 82, text-fig. 5a

Budapanopeus denticulatus MÜLLER & COLLINS,
 1991
 holotype, M.91.155
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 72, text-fig. 4b, pl. 4, fig. 15
 Paratypes under same inventory number
 (pl. 4, fig. 16)

Budapanopeus denticulatus MÜLLER & COLLINS,
 1991
 paratypes, M.91.154
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 72, text-fig. 4b, pl. 5, figs. 2, 7

Budapanopeus denticulatus MÜLLER & COLLINS,
 1991
 paratype, M.91.156
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 72, text-fig. 4b, pl. 5, fig. 14

Budapanopeus denticulatus MÜLLER & COLLINS, 1991
 paratype, M.91.157
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 72, text-fig. 4b

Budapanopeus denticulatus MÜLLER & COLLINS, 1991
 paratypes, M.91.158, M.91.159
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 72, text-fig. 4b

Caprocancer altus MÜLLER & COLLINS, 1991
 holotype, M.91.204
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 84, text-fig. 5e, pl. 8, fig. 1
 Paratypes under same inventory number
 (pl. 8, figs. 2, 3)

Corallicarcinus planus MÜLLER & COLLINS, 1991
 holotype, M.91.207
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 85, text-fig. 5b, pl. 7, figs. 2, 3
 Paratypes under same inventory number
 (pl. 7, fig. 8)

Cynonomus primitivus MÜLLER & COLLINS, 1991
 holotype, M.91.135
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 63, text-fig. 3g, pl. 3, fig. 6

Daragrapus trispinosus MÜLLER & COLLINS, 1991
 holotype, M.91.209
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 88, text-fig. 5h, pl. 7, figs. 9, 12, 13, 14
 Paratypes under same inventory number
 (pl. 7, fig. 10)

Daragrapus trispinosus MÜLLER & COLLINS, 1991
 paratypes, M.91.208
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 88, text-fig. 5h

Daragrapus trispinosus MÜLLER & COLLINS, 1991
 paratypes, M.91.210, M.91.211, M.91.212
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 88, text-fig. 5h

Daragrapus trispinosus MÜLLER & COLLINS, 1991
 paratypes, M.91.213, M.91.214
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 88, text-fig. 5h

Dardanus curtimanus MÜLLER & COLLINS, 1991
 holotype, M.91.99
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 51, text-fig. 2b, pl. 1, figs. 4, 5

Dardanus curtimanus MÜLLER & COLLINS, 1991
 paratype, M.91.98
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 51, text-fig. 2b, pl. 1, fig. 6

Diogenes longimanus MÜLLER & COLLINS, 1991
 holotype, M.91.101
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 52, text-fig. 2d, pl. 1, fig. 11
 Paratype under same inventory number
 (pl. 1, fig. 10)

Dromilites fossata MÜLLER & COLLINS, 1991
 holotype, M.91.131
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 62, text-fig. 3b, pl. 2, figs. 13, 16

Dromilites fossata MÜLLER & COLLINS, 1991
 paratype, M.91.130
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 62, text-fig. 3b

Dromilites subglobosa MÜLLER & COLLINS, 1991
 holotype, M.91.127
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 62, text-fig. 3c, pl. 3, figs. 1, 2, 3
 Paratypes under same inventory number

- Dromilites subglobosa* MÜLLER & COLLINS, 1991
 paratypes, M.91.128
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 62, text-fig. 3c
- Dromilites subglobosa* MÜLLER & COLLINS, 1991
 paratype, M.91.129
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 62, text-fig. 3c
- Eomaldivia pannonica* MÜLLER & COLLINS, 1991
 holotype, M.91.190
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 81, text-fig. 4k, pl. 6, figs. 10, 13
 Paratypes under same inventory number
- Eomaldivia trispinosa* MÜLLER & COLLINS, 1991
 holotype, M.91.191
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 81, text-fig. 4l, pl. 6, figs. 8, 11
 Paratypes under same inventory number
- Eomaldivia trispinosa* MÜLLER & COLLINS, 1991
 paratypes, M.91.192
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 81, text-fig. 4l
- Eoplax minima* MÜLLER & COLLINS, 1991
 holotype, M.91.219
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 85, text-fig. 5d, pl. 8, figs. 6, 7
- Eoplax minima* MÜLLER & COLLINS, 1991
 paratypes, M.91.220
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 85, text-fig. 5d
- Ethusa evae* MÜLLER & COLLINS, 1991
 holotype, M.91.144
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 66, text-fig. 3h, pl. 4, figs. 1, 2
- Galathea (Acanthogalathea) parva* MÜLLER & COLLINS, 1991
 holotype, M.91.106
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 56, text-fig. 2h, pl. 2, fig. 3
- Gemmacarcinus fossatus* MÜLLER & COLLINS, 1991
 holotype, M.91.145
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 67, text-fig. 3j, pl. 3, fig. 13
- Goniocypoda transsilvanica* BITTNER, 1893
 holotype, M.68.1887
 Priabonian, Cluj Limestone Formation
 quarry, Suceagu (Szucság), Romania
 Btt93, p. 24, pl. 2, fig. 4
- Harpactocarcinus hungaricus* TOMOR-THIRRING, 1936
 syntype, M.74.145
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 65, text-fig. 15, pl. 2, figs. 11–12
- Harpactocarcinus hungaricus* TOMOR-THIRRING, 1936
 syntype, M.74.167
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 65
- Harpactocarcinus telegdi-rothi* TOMOR-THIRRING, 1936
 syntype, M.74.143
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 61, text-fig. 14, pl. 2, fig. 9
- Harpactocarcinus telegdi-rothi* TOMOR-THIRRING, 1936
 syntypes, M.74.168
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 61

Harpactocarcinus telegdi-rothi baconica TOMOR-THIRRING, 1936
 syntype, M.74.144
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 63, pl. 2, fig. 10

Harpactocarcinus telegdi-rothi baconica TOMOR-THIRRING, 1936
 syntype, M.74.172
 Bartonian, Szőc Limestone Formation
 Sűrű-hegy, Dúdar
 Tom36, p. 63
 Type missing

Kromtisis pentagonalis MÜLLER & COLLINS, 1991
 holotype, M.91.133
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 63, text-fig. 3e, pl. 3, figs. 4, 8
 Paratypes under same inventory number

Kromtisis pentagonalis MÜLLER & COLLINS, 1991
 paratypes, M.91.132
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 63, text-fig. 3e

Kromtisis pentagonalis MÜLLER & COLLINS, 1991
 paratype, M.91.134
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 63, text-fig. 3e, pl. 3, fig. 5

Longoporellana denticulata MÜLLER & COLLINS, 1991
 holotype, M.91.120
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 57, text-fig. 2j, pl. 2, figs. 4, 5
 Paratype under same inventory number;
 author expressed doubt in identifying with
 holotype

Longoporellana denticulata MÜLLER & COLLINS, 1991
 paratype, M.91.121
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 57, text-fig. 2j

Mesolambrus declinatus MÜLLER & COLLINS, 1991
 holotype, M.91.151
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 69, text-fig. 3f, pl. 3, figs. 11, 12
 Paratype under same inventory number

Nanomaja simplex MÜLLER & COLLINS, 1991
 holotype, M.91.149
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 68, text-fig. 3d, pl. 4, figs. 5, 6
 Paratypes under same inventory number

Nanomaja simplex MÜLLER & COLLINS, 1991
 paratypes, M.91.147
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 68, text-fig. 3d

Nanomaja simplex MÜLLER & COLLINS, 1991
 paratypes, M.91.148
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 68, text-fig. 3d

Ovamene franciae MÜLLER & COLLINS, 1991
 holotype, M.91.143
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 66, text-fig. 3i, pl. 3, fig. 7

Ovocarcinus elongatus MÜLLER & COLLINS, 1991
 holotype, M.91.126
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 60, text-fig. 3a, pl. 2, figs. 9, 10, 11

Paguristes oligotuberculatus MÜLLER & COLLINS, 1991
 holotype, M.91.104
 Priabonian, Szépvölgy Limestone Formation
 Francia quarry, Budapest
 Mül91a, p. 54, text-fig. 2e, pl. 1, fig. 2

Pagurus latidactylus MÜLLER & COLLINS, 1991
holotype, M.91.100
Priabonian, Szépvölgy Limestone Formation
Francia quarry, Budapest
Mül91a, p. 52, text-fig. 2c, pl. 1, fig. 9
Paratypes under same inventory number
(pl. 1, figs. 7, 8). Author expressed doubt in
identifying specimen figured on fig. 7 with
holotype

Palaeograpsus parvus MÜLLER & COLLINS, 1991
holotype, M.91.227
Priabonian, Szépvölgy Limestone Formation
Francia quarry, Budapest
Mül91b, p. 140
Replacement name for *Palaeograpsus bittneri*
MÜLLER & COLLINS, 1991. Description in
MÜLLER & COLLINS 1991a: p. 9, text-fig. 5i,
pl. 8, figs. 11, 12, 15

Panopeus granulineatus MÜLLER & COLLINS, 1991
holotype, M.91.167
Priabonian, Szépvölgy Limestone Formation
Gugger-hegy, Budapest
Mül91a, p. 74, text-fig. 4d, pl. 5, figs. 4, 6
Paratypes under same inventory number
(pl. 5, fig. 3)

Panopeus granulineatus MÜLLER & COLLINS, 1991
paratypes, M.91.165
Priabonian, Szépvölgy Limestone Formation
Ruprecht quarry, Felső-Kecske-hegy, Budapest
Mül91a, p. 74, text-fig. 4d

Panopeus granulineatus MÜLLER & COLLINS, 1991
paratype, M.91.166
Priabonian, Szépvölgy Limestone Formation
Bimbó út 185., Budapest
Mül91a, p. 74, text-fig. 4d

Paraxanthosia budensis MÜLLER & COLLINS, 1991
holotype, M.91.168
Priabonian, Szépvölgy Limestone Formation
Ruprecht quarry, Felső-Kecske-hegy, Budapest
Mül91a, p. 75, text-fig. 4f, pl. 4, fig. 12
Paratypes under same inventory number
(pl. 4, figs. 13, 14)

Paraxanthosia budensis MÜLLER & COLLINS, 1991
paratype, M.91.169
Priabonian, Szépvölgy Limestone Formation
Bimbó út 185., Budapest
Mül91a, p. 75, text-fig. 4f
Author expressed doubt in identifying with
holotype

Petrolisthes? striatissimus MÜLLER & COLLINS,
1991
holotype, M.91.123
Priabonian, Szépvölgy Limestone Formation
Gugger-hegy, Budapest
Mül91a, p. 59, text-fig. 2l, pl. 2, fig. 8

Pilumnomimus planidentatus MÜLLER &
COLLINS, 1991
holotype, M.91.170
Priabonian, Szépvölgy Limestone Formation
Ruprecht quarry, Felső-Kecske-hegy, Budapest
Mül91a, p. 75, text-fig. 4h, pl. 5, fig. 10

Pilumnomimus planidentatus MÜLLER &
COLLINS, 1991
paratype, M.91.171
Priabonian, Szépvölgy Limestone Formation
Gugger-hegy, Budapest
Mül91a, p. 75, text-fig. 4h, pl. 5, figs. 5, 8

Polyonyx arcuatus MÜLLER & COLLINS, 1991
holotype, M.91.122
Priabonian, Szépvölgy Limestone Formation
Ruprecht quarry, Felső-Kecske-hegy, Budapest
Mül91a, p. 59, text-fig. 2k, pl. 2, figs. 6, 7

Priabonocarcinus gallicus MÜLLER & COLLINS,
1991
holotype, M.91.180
Priabonian, Szépvölgy Limestone Formation
Francia quarry, Budapest
Mül91a, p. 78, text-fig. 4e, pl. 5, figs. 12, 15
Paratypes under same inventory number
(pl. 5, fig. 16)

Prochlorodius ellipticus MÜLLER & COLLINS, 1991
holotype, M.91.181
Priabonian, Szépvölgy Limestone Formation
Francia quarry, Budapest
Mül91a, p. 78, text-figs. 4i-j, pl. 6, figs. 4, 7
Paratypes under same inventory number

Prochlorodius ellipticus MÜLLER & COLLINS, 1991
 paratypes, M.91.182
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 78, text-figs. 4i-j, pl. 6, fig. 17

Sculptoplax rigida MÜLLER & COLLINS, 1991
 holotype, M.91.228
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 90, text-fig. 5g, pl. 8, fig. 13
 Paratype under same inventory number

Prochlorodius ellipticus MÜLLER & COLLINS, 1991
 paratypes, M.91.183
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 78, text-figs. 4i-j, pl. 5, fig. 11
 Juvenile specimens, author expressed doubt in identifying with holotype

Telphusograpsus laevis LÖRENTHEY, 1901
 syntype, M.68.1888
 Priabonian, Cluj Limestone Formation
 quarry, Suceagu (Szucság), Romania
 Lőr01, p. 819, pl. 2, figs. 2a-c (pl. 2, fig. 3)
 Translated description in German edition, LÖRENTHEY 1903: p. 115

Prochlorodius ellipticus MÜLLER & COLLINS, 1991
 paratypes, M.91.184, M.91.185
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 78, text-figs. 4i-j

Trapezia loerentheyi MÜLLER, 1975
 holotype
 Priabonian, Szépvölgy Limestone Formation
 Felső-Kecske-hegy, Budapest
 Mül75b, p. 520 (517), pl. 1, fig. 1
Tetralia loerentheyi, Mül91a
 Author's inventory number: EK-1-1. Type missing, not found in inventory

Protomunida pentacantha MÜLLER & COLLINS, 1991
 holotype, M.91.116
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 57, text-fig. 2i, pl. 1, fig. 16
 Paratypes under same inventory number

Trapezia loerentheyi MÜLLER, 1975
 paratype, M.91.194
 Priabonian, Szépvölgy Limestone Formation
 Felső-Kecske-hegy, Budapest
 Mül75b, p. 520 (517), pl. 1, fig. 2
Tetralia loerentheyi, Mül91a
 Author's inventory number: EK-1-1

Protomunida pentacantha MÜLLER & COLLINS, 1991
 paratypes, M.91.115
 Priabonian, Szépvölgy Limestone Formation
 Ruprecht quarry, Felső-Kecske-hegy, Budapest
 Mül91a, p. 57, text-fig. 2i, pl. 1, fig. 15

6.10. Eocene Echinoidea

Protomunida pentacantha MÜLLER & COLLINS, 1991
 paratypes, M.91.117
 Priabonian, Szépvölgy Limestone Formation
 Bimbó út 185., Budapest
 Mül91a, p. 57, text-fig. 2i

Conoclypus platysoma SZÖRÉNYI, 1929
 holotype, M.59.5094
 Priabonian, Buda Marl Formation
 Kecske-hegy, Óbuda, Budapest
 Szö29, p. 14, pl. 1, fig. 2
 Inventory number in original publication: 566/1860

Protomunida pentacantha MÜLLER & COLLINS, 1991
 paratypes, M.91.118, M.91.119
 Priabonian, Szépvölgy Limestone Formation
 Gugger-hegy, Budapest
 Mül91a, p. 57, text-fig. 2i

Echinolampas plesiobathystoma SZÖRÉNYI, 1929
 holotype, M.61.2293
 Priabonian, Buda Marl Formation
 Várhegy tunnel, Budapest
 Szö29, p. 18, pl. 1, fig. 8
 Inventory number in original publication: K. 949

- Rabdocidaris posthumus* PÁVAY, 1874
holotype, V.61.893
Priabonian, Buda Marl Formation
Várhegy tunnel, Budapest
Páv74a (Páv74b), p. 87 (243), pl. 10, fig. 1
Correct spelling of genus name is *Rhabdocidaris*
- Spatagoides (Toxopatagus) várhegyensis*
SZÖRÉNYI, 1929
holotype, M.61.2292
Priabonian, Buda Marl Formation
Várhegy tunnel, Budapest
Szö29, p. 20, pl. 1, fig. 5
Inventory number in original publication:
K. 950
- Titanaster labiostoma* SZÖRÉNYI, 1929
syntypes, M.61.2187
Priabonian, Buda Marl Formation
Várhegy tunnel, Budapest
Szö29, p. 19 (pl. 1, fig. 10)
- 7. Oligocene types**
- 7.1. Oligocene Foraminifera**
- Amphicoryne marginuliniformis* NYÍRŐ, 1961
holotype, M.63.13
Egerian, Törökálint Sandstone Formation
road cut below the vineyard called "Világos",
Törökálint
Nyí61, p. 49, text-fig. 1
Form A. Inventory number in original
publication: M.60.580
- Amphicoryne marginuliniformis* NYÍRŐ, 1961
paratype, M.63.29
Kiscellian, Kiscell Clay Formation
brickyard, Törökálint
Nyí61, p. 49, text-fig. 2
Form B. Inventory number in original
publication: M.60.581
- Bulimina parvula* FRANZENAU, 1892
syntypes
Kiscellian, Kiscell Clay Formation
Romhány
Fra92, p. 139 (108)
Types probably destroyed in 1956, figured
in FRANZENAU & MAJZON 1956: p. 215, pl. 1,
figs. 44–46
- Bulimina triquetra* FRANZENAU, 1892
syntypes
Kiscellian, Kiscell Clay Formation
Romhány
Fra92, p. 139 (108)
Reusella triquetra, Fra56
Types probably destroyed in 1956,
figured in FRANZENAU & MAJZON 1956:
p. 215, pl. 1, figs. 31–33
- Cristellaria (Robulina) arcuatostriata*
HANTKEN, 1868
paralectotype, 2006.96.1.
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 93, pl. 2, figs. 30a–b
Lenticulina arcuatostriata, Bra75; Hor03
Type figured in HORVÁTH 2003 under
inventory number M.01.18
- Cristellaria (Robulina) kubinyi* HANTKEN,
1868
paralectotype, M.99.77
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 92, pl. 2, figs. 29a–b
Planularia kubinyii, Cuv49; Hor03
Type figured in HORVÁTH 2003
- Cristellaria minuta* HANTKEN, 1875
neotype, M.99.76
Kiscellian, Kiscell Clay Formation
brickyard, Pilisborosjenő
Han75a (Han75b), p. 50 (43), pl. 14, fig. 7
Vaginulinopsis minutus, Szt79; Hor03
Neotype designated in HORVÁTH 2003
- Cristellaria propinqua* HANTKEN, 1875
neotype, M.99.66
Kiscellian, Kiscell Clay Formation
Nyárjas-tető, Novaj
Han75a (Han75b), p. 52 (45), pl. 5, fig. 4
Saracenaria propinqua, Maj62; Hor03
Neotype designated in HORVÁTH 2003
- Cristellaria (Marginulina) tunicata*
HANTKEN, 1868
neotype, M.99.70
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 91, pl. 2, figs. 24a–c
Amphicoryna tunicata, Maj62; Hor03
Neotype designated in HORVÁTH 2003

Dentalina gümbeli HANTKEN, 1875
 neotype, 2006.90.1.
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 38 (32), pl. 4, fig. 1
 Neotype designated in HORVÁTH 2003
 under inventory number M.01.12

Dentalina semilaevis HANTKEN, 1875
 paralectotype, 2006.91.1.
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 39 (32), pl. 4, fig. 6,
 pl. 12, fig. 13
 Type figured in HORVÁTH 2003 under
 inventory number M.01.13

Dentalina setosa HANTKEN, 1875
 neotype, 2006.95.1.
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 39 (33), pl. 13, fig. 9
Nodosaria setosa, Hor03
 Neotype designated in HORVÁTH 2003
 under inventory number M.01.17

Eponides majzoni NYÍRÓ, 1960
 holotype, M.58.99
 Kiscellian, Kiscell Clay Formation
 borehole Demjén-11, Demjén
 Nyí60, p. 36, pl. 1, figs. 4a-b
 Type missing

Flabellina budensis HANTKEN, 1875
 neotype, M.99.69
 Kiscellian, Kiscell Clay Formation
 brickyard, Pilisborosjenő
Han75a (*Han75b*), p. 44 (37), pl. 4, fig. 17
Palmula budensis, Maj62; Hor03
 Neotype designated in HORVÁTH 2003

Frondicularia superba HANTKEN, 1875
 neotype, M.99.68
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 42 (36), pl. 4, fig. 16
Frondovaginulina superba, Hor03
 Neotype designated in HORVÁTH 2003

Gaudryina irregularis HANTKEN, 1875
 paralectotype, 2006.99.1.
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 15 (12), pl. 1, fig. 7
Eggerella irregularis, Szt79; Hor02
 Type figured in HORVÁTH 2002 under
 inventory number M.00.00

Haplophragmium acutidorsatum HANTKEN,
 1868
 paralectotype, M.99.39
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han68, p. 82, pl. 1, figs. 1a-b
Reticulophragmium acutidorsatum, Cic98; Hor02
 Type figured in HORVÁTH 2002

Haplophragmium rotundidorsatum HANTKEN,
 1875
 lectotype, M.99.40
 Kiscellian, Kiscell Clay Formation
 Újlak, Budapest
Han75a (*Han75b*), p. 12 (10), pl. 1, fig. 2
Reticulophragmium rotundidorsatum, Cic98;
 Hor02
 Lectotype designated in HORVÁTH 2002

Heterostegina margaritacea MAJZON, 1960
 holotype
 Egerian, Eger Formation
 borehole No. 1, Bogács
 Maj60, p. 362 (360), pl. 18, fig. 1
 Type probably lost

Heterostegina rupelica MAJZON, 1960
 holotype
 Egerian, Eger Formation
 borehole No. 1, Bogács
 Maj60, p. 362 (360), pl. 18, fig. 2
 Type probably lost

Lepidocyclina (Eulepidina) hungarica
 KECSKEMÉTI, 1961
 holotype, M.60.10962
 Egerian, Eger Formation
 Nyárjas-tető, Novaj
 Bál61, p. 85, pl. 3, figs. 2-3
Lepidocyclina (Eulepidina) dilatata, Les91
 Paratypes under same inventory number

- Lepidocyclina (Eulepidina) hungarica*
KECSKEMÉTI, 1961
paratypes, M.61.6
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 85, pl. 3, fig. 1
Lepidocyclina (Eulepidina) dilatata, Les91
- Lingulina costata seminuda* HANTKEN, 1875
neotype, M.99.60
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han75a (Han75b), p. 41 (35), pl. 4, figs. 8a-b
Lingulina seminuda, Maj62; Hor03
Neotype designated in HORVÁTH 2003
- Marginulina budensis* HANTKEN, 1875
neotype, 2006.97.1.
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han75a (Han75b), p. 47 (40), pl. 14, fig. 5
Astacolus budensis, Hor03
Neotype designated in HORVÁTH 2003
under inventory number M.01.19
- Marginulina indifferens* HANTKEN, 1875
neotype, M.99.71
Kiscellian, Kiscell Clay Formation
borehole VIII/1, Budapest
Han75a (Han75b), p. 47 (40), pl. 4, fig. 14
Astacolus indifferens, Hor03
Neotype designated in HORVÁTH 2003
- Marginulina splendens* HANTKEN, 1875
neotype, M.99.75
Kiscellian, Kiscell Clay Formation
brickyard, Pilisborosjenő
Han75a (Han75b), p. 87 (40), pl. 4, fig. 11
Hemirobulina splendens, Hor03
Neotype designated in HORVÁTH 2003.
Type lost
- Nodosaria acuminata* HANTKEN, 1875
neotype, M.99.50
Kiscellian, Kiscell Clay Formation
Kis-Eged-hegy, road cut, Eger
Han75a (Han75b), p. 28 (23), pl. 2, fig. 9,
pl. 13, fig. 5
Dentalina acuminata, Hor03
Neotype designated in HORVÁTH 2003
- Nodosaria bacilloides* HANTKEN, 1868
neotype, M.99.51
Kiscellian, Kiscell Clay Formation
brickyard, Pilisborosjenő
Han68, p. 86, pl. 1, fig. 9
Dentalina bacilloides, Hor03
Neotype designated in HORVÁTH 2003
- Nodosaria budensis* HANTKEN, 1875
neotype, M.99.52
Kiscellian, Kiscell Clay Formation
brickyard, Pilisborosjenő
Han75a (Han75b), p. 28 (23), pl. 2, fig. 10
Dentalina budensis, Maj62; Hor03
Neotype designated in HORVÁTH 2003
- Nodosaria (Dentalina) contorta* HANTKEN,
1868
paralectotype, 2006.89.1.
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 89, pl. 1, fig. 16
Dentalina contorta, Han75; Hor03
Type figured in HORVÁTH 2003 under
inventory number M.01.11
- Nodosaria (Dentalina) majzoni* JASKÓ, 1940
syntypes, M 2008.1.1.
Egerian, Szécsény Formation
between the Rima and Tarna rivers,
northern Hungary or southern Slovakia
Jas40, p. 370 (304), pl. 9, figs. 21-22
- Nodosaria (Dentalina) reitzi* HANTKEN, 1868
neotype, 2006.94.1.
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 88, pl. 1, fig. 13
Neotype designated in HORVÁTH 2003
under inventory number M.01.16
- Nodosaria (Dentalina) simplex* HANTKEN,
1868
neotype, 2006.92.1.
Kiscellian, Kiscell Clay Formation
brickyard, Pilisborosjenő
Han68, p. 87, pl. 1, fig. 11
Laevidentalina simplex, Hor03
Neotype designated in HORVÁTH 2003
under inventory number M.01.14

Patellina legányii KENAWY & NYÍRŐ, 1967
holotype, M.66.1054
Egerian, Eger Formation
Wind brickyard, Eger
Ken67, p. 103, pl. 1, figs. 1-2
Type missing

Plecanium elegans HANTKEN, 1868
paralectotype, M.99.46
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han68, p. 83, pl. 1, fig. 5
Type figured in HORVÁTH 2002.
Type lost

Pseudopatellina plana KENAWY & NYÍRŐ, 1967
holotype, M.66.1055
Egerian, Eger Formation
Wind brickyard, Eger
Ken67, p. 104, pl. 1, figs. 3-4
Type missing

Pulvinulina romhányensis FRANZENAU, 1892
syntypes
Kiscellian, Kiscell Clay Formation
Romhány
Fra92, p. 142 (112)
Cibicides romhanyensis, Fra56
Types probably destroyed in 1956,
figured in FRANZENAU & MAJZON 1956:
p. 216, pl. 1, figs. 15-1

Robulina budensis HANTKEN, 1875
paralectotype, M.99.63
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Han75a (Han75b), p. 58 (49), pl. 7, fig. 1
Lenticulina budensis, Hor03
Type figured in HORVÁTH 2003

Siphonina pseudocarinata NYÍRŐ, 1961
holotype, M.60.10964
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 76, pl. 1, figs. 3a-b
Inventory number in original publication:
M.60.10963

Textilaria elongata HANTKEN, 1875
neotype, M.99.45
Kiscellian, Tard Clay Formation
borehole Kiscell-1, Budapest
Han75a (Han75b), p. 67 (57), pl. 15, fig. 3
Textularia elongata, Szt79; Hor02
Neotype designated in HORVÁTH 2002.
Type lost

Textularia carinata mucronata JASKÓ, 1940
syntypes, M 2008.2.1.
Egerian, Szécsény Formation
between the Rima and Tarna rivers,
northern Hungary or southern Slovakia
Jas40, p. 370 (305), pl. 9, figs. 23-25
Referred to in plate explanation as
Textularia carinata schréteri

Tritaxilina hantkeni CUSHMAN, 1936
paratype, M.99.48
Kiscellian, Kiscell Clay Formation
Újlak, Budapest
Cus36, p. 41 (pl. 3, fig. 13)
Type figured in HORVÁTH 2002

Turborotalia munda franzenauí SZTRÁKOS, 1974
holotype, M.73.178
Kiscellian, Kiscell Clay Formation
borehole Egerszalók-2a, Egerszalók
Szt74, p. 48, pl. 4, figs. 1a-c

Turborotalia munda franzenauí SZTRÁKOS, 1974
paratype, M.73.177
Kiscellian, Kiscell Clay Formation
Vár-hegy, Budapest
Szt74, p. 48, pl. 4, figs. 2a-c

Vaginulina plana NYÍRŐ, 1961
holotype, M.60.10963
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 74, pl. 1, fig. 4

7.2. Oligocene Bivalvia

Aequipecten deleatus harmati NOSZKY, 1939
holotype, M.59.5039
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 110 (38), pl. 1, fig. 3

- Aequipecten deletus sasselloniensis* NOSZKY, 1939
holotype, M.59.5038
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 110 (38)
- Aequipecten oligosquamosus szörényii* NOSZKY, 1939
holotype, M.59.5040
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 131 (86), pl. 1, fig. 1
- Aequipecten raulini tasnádii* NOSZKY, 1939
holotype, M.59.5090
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 111 (39), pl. 1, fig. 2
Paratype under same inventory number
- Aequipecten scabrellus zsivonyi* NOSZKY, 1939
holotype, M.59.5037
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 110 (38), pl. 1, fig. 6
- Amussium bronni zimányii* NOSZKY, 1939
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 110 (37), pl. 1, fig. 7
Type probably destroyed in 1956
- Amussium costulatum* NOSZKY, 1939
holotype, M.59.5036
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 109 (36), pl. 1, fig. 4
- Amussium semiradiatum tricosta* NOSZKY, 1939
holotype, M.59.5035
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 108 (36), pl. 1, fig. 5
- Angulus (Peronidia) nysti subfallax* BÁLDI, 1963
paratype, M.62.9347
Egerian, Törökbálint Sandstone Formation
Törökbálint
Bál63, p. 82, pl. 4, fig. 9
Angulus (Peronidia) nysti, Bál73
- Angulus (Peronidia) nysti subfallax* BÁLDI, 1963
paratype, M.62.9353
Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint
Bál63, p. 82, pl. 4, figs. 10–11
Angulus (Peronidia) nysti, Bál73
- Angulus (Peronidia) planatus ancestralis* BÁLDI, 1973
holotype, M.65.647
Egerian, Törökbálint Sandstone Formation
borehole Pomáz-22, Pomáz
Bál73, p. 227, pl. 20, fig. 3
- Angulus (Peronidia) planatus ancestralis* BÁLDI, 1973
paratype, M.97.205
Egerian, Törökbálint Sandstone Formation
borehole Dj-3, Diósjenő
Bál73, p. 227
- Arca (Acar) dactylus brevis* NOSZKY, 1939
syntypes, M.59.3981
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 116 (48)
- Arca (Acar) dactylus erecta* NOSZKY, 1939
holotype, M.59.5050
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 116 (49), pl. 2, fig. 1
Barbatia (Acar) dactylus erecta, Bál86
Type figured in BÁLDI 1986: pl. 3, fig. 29
- Arcopsis (Arcopsis) hofmanni* BÁLDI, 1986
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 74, pl. 4, fig. 30
Type missing, not found in inventory

Avicula hirundo elongata NOSZKY, 1939
 holotype, M.59.4010
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 115 (45)
Pteria hirundo wemmelensis, Bál86

Avicula hirundo vogli NOSZKY, 1939
 holotype, M.59.5047
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 115 (45), pl. 1, fig. 11
Pteria hirundo wemmelensis, Bál86
 Type figured in BÁLDI 1986: pl. 4, fig. 33

Avicula hirundo vogli NOSZKY, 1939
 paratype, M.59.4009
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 115 (45)
Pteria hirundo wemmelensis, Bál86

Axinus (Cryptodon) subangulatus pongráczi
 NOSZKY, 1939
 holotype, M.59.5181
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 120 (59), pl. 3, fig. 15
Thyasira vara, Bál86
 Paratype under same inventory number.
 Types figured in BÁLDI 1986: pl. 7, figs. 67–69

Cardiomya noszkyi BÁLDI, 1986
 holotype, M.59.4371
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Bál86, p. 83, pl. 9, fig. 100

Cardita (Cyclocardia) orbicularis subparvocostata
 BÁLDI, 1963
 holotype, M.62.9328
 Egerian, Törökbálint Sandstone Formation
 outcrop T2, Törökbálint
 Bál63, p. 77, pl. 2, fig. 8 (pl. 2, fig. 11)

Cardita (Cyclocardia) orbicularis subparvocostata
 BÁLDI, 1963
 paratypes, M.62.9329
 Egerian, Törökbálint Sandstone Formation
 outcrop T2, Törökbálint
 Bál63, p. 77, pl. 2, figs. 6–7

Cardium (Ringicardium) bükkianum serratulum
 NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 110 (88)
 Type destroyed in 1956

Cardium (Ringicardium) grateloupi oligocenicum
 NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 111 (88), pl. 6, fig. 7
 Type destroyed in 1956

Cardium (?Trachycardium) neglectum
intersulcatum BÁLDI, 1973
 syntype
 Egerian, Törökbálint Sandstone Formation
 borehole So-72, Solymár
 Bál73, p. 208
 Type missing, not found in inventory

Cardium (Parvicardium) praepapillosum BÁLDI,
 1966
 holotype, M.65.1080
 Egerian, Eger Formation
 Wind brickyard, Eger
 Bál66, p. 86, pl. 1, fig. 12

Cardium scobinula boggschi NOSZKY, 1939
 holotype, M.59.5061
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 119 (56), pl. 2, fig. 12
Poromya (Myoporomya) bicarinata boggschi, Bál86
 Paratype under same inventory number.
 Types figured in BÁLDI 1986: pl. 8, figs. 91, 92

Cardium (Parvicardium) transversale szontaghi
 NOSZKY, 1939
 holotype
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 119 (57), pl. 2, fig. 13
 Type probably destroyed in 1956

- Chlamys agriensis* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2848
Egerian, Eger Formation
well of "Vincellér" school, Eger
C-M60, p. 31, pl. 20, fig. 15 (pl. 20, fig. 16)
Inventory number in original publication:
M.59.284
- Chlamys (Aequipecten) csepreghy-meznericsae* BÁLDI, 1961
holotype, M.60.10787
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 90, pl. 3, fig. 6
"*Aequipecten*" *csepreghymeznericsae*, Bál99
- Chlamys (Aequipecten) csepreghy-meznericsae* BÁLDI, 1961
paratype, M.60.10788
Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint
Bál61, p. 90, pl. 3, fig. 7
"*Aequipecten*" *csepreghymeznericsae*, Bál99
- Crassatella (Crassatina) bosqueti minor* BÁLDI, 1961
holotype, M.60.10780
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 93, pl. 3, figs. 4a-b
- Crassatella (Crassatina) bosqueti minor* BÁLDI, 1961
paratypes, M.60.10781
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 93, pl. 4, figs. 1a-b
- Crassatella (Crassatina) bosqueti minor* BÁLDI, 1961
paratypes, M.60.10782
Egerian, Eger Formation
Nyárjas-tető, Novaj
Bál61, p. 93
- Cultellus budensis* BÁLDI, 1973
syntypes, M.97.571, M.97.576, M.97.588,
M.97.612, 2007.1.1.
Egerian, Törökbálint Sandstone Formation
borehole So-72, Solymár
Bál73, p. 229, text-figs. 54.1-3.
- Cuspidaria neoscalarina* BÁLDI, 1966
holotype, M.65.1081
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 86, pl. 1, fig. 9
- Cyrena sirena pygmea* NOSZKY, 1939
holotype, M.59.5065
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 124 (68), pl. 3, fig. 4
- Cytherea (Meretrix) incrassata koeneni* NOSZKY, 1939
holotype, M.59.5066
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 124 (69), pl. 3, fig. 3
- Cytherea (Callista) taurorugosa soósi* NOSZKY, 1939
holotype, M.59.5067
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 124 (71), pl. 3, fig. 11
- Cytherocardia oligocenica* NOSZKY, 1939
holotype, M.59.5064
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 124 (67), pl. 3, fig. 12
Paratype under same inventory number
- Dimyodon similis interstricta* NOSZKY, 1939
syntypes, M.59.5095
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 107 (34)
- Dymia fragilis punctata* NOSZKY, 1939
holotype, M.59.5089
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 132 (87)

Flabellipecten burdigalensis angustiformis

BÁLDI, 1963

holotype, M.62.9312

Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint

Bál63, p. 74, pl. 1, fig. 3

Flabellipecten angustiformis, Bál73

Flabellipecten burdigalensis angustiformis

BÁLDI, 1963

paratype, M.62.9313

Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint

Bál63, p. 74, pl. 1, fig. 4

Flabellipecten angustiformis, Bál73

Flabellipecten telegdi-rothi CSEPREGHY-

MEZNERICS, 1960

holotype, M.59.2841

Egerian, Eger Formation

Wind brickyard, Eger

C-M60, p. 16, pl. 11, fig. 7 (pl. 11, fig. 6)

"Aequipecten" *telegdirothi*, Bál99

Left valve

Flabellipecten telegdi-rothi CSEPREGHY-

MEZNERICS, 1960

holotype, M.59.2841

Egerian, Eger Formation

Wind brickyard, Eger

C-M60, p. 16, pl. 11, fig. 8

"Aequipecten" *telegdirothi*, Bál99

Right valve

Flabellipecten telegdi-rothi CSEPREGHY-

MEZNERICS, 1960

paratypes, M.62.5152

Egerian, Eger Formation

Wind brickyard, Eger

C-M60, p. 16

"Aequipecten" *telegdirothi*, Bál99

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

holotype, M.61.8471

Egerian, Törökbálint Sandstone Formation

Törökbálint

Bál62, p. 103, pl. 4, figs. 1a-c

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8467

Egerian, Törökbálint Sandstone Formation
outcrop T1, Törökbálint

Bál62, p. 103, pl. 4, figs. 2-6, pl. 5, figs. 1-5,
pl. 8, fig. 6

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8469

Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint

Bál62, p. 103, pl. 1, figs. 2d-e

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8470

Egerian, Törökbálint Sandstone Formation
outcrop T2, Törökbálint

Bál62, p. 103, pl. 1, figs. 2a-c

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8516, M.61.8517, M.61.8518,
M.61.8519, M.61.8528, M.61.8529, M.61.8530,
M.61.8532, M.61.8533, M.61.8534, M.61.8536,
M.63.5138, M.63.5139, M.63.5140, M.63.5141,
M.63.5142, M.63.5143, M.63.5144, M.63.5145,
M.63.5146

Egerian, Törökbálint Sandstone Formation
outcrop T1, Törökbálint

Bál62, p. 103

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8520, M.61.8531, M.61.8535

Egerian, Törökbálint Sandstone Formation
Törökbálint

Bál62, p. 103

Glycymeris (Glycymeris) latiradiata obovatooides

BÁLDI, 1962

paratypes, M.61.8521, M.61.8522, M.61.8523,
M.61.8524, M.61.8525, M.61.8526, M.61.8527

Egerian, Törökbálint Sandstone Formation
Pacsirta-hegy, Budapest

Bál62, p. 103

- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
holotype, M.61.8474
Egerian, Kováčov Formation
Kováčspatak, Kamenica nad Hronom
(Garamkövesd), Slovakia
Bál62, p. 106, pl. 6, figs. 1a-d (pl. 6, figs. 2, 7)
- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
paratype, M.61.8468
Egerian, Eger Formation
Wind brickyard, Eger
Bál62, p. 106, pl. 1, fig. 3
- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
paratype, M.61.8472
Egerian, Eger Formation
Wind brickyard, Eger
Bál62, p. 106, pl. 5, figs. 6a-c
- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
paratype, M.61.8473
Egerian, Eger Formation
Wind brickyard, Eger
Bál62, p. 106, pl. 6, fig. 3
- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
paratypes, M.61.8539, M.61.8540
Egerian, Kováčov Formation
Kováčspatak, Kamenica nad Hronom
(Garamkövesd), Slovakia
Bál62, p. 106
- Glycymeris (Glycymeris) latiradiata subfichteli*
BÁLDI, 1962
paratypes, M.61.8543
Egerian, Eger Formation
Wind brickyard, Eger
Bál62, p. 106
- Gryphaea brongniarti exalata* NOSZKY, 1939
syntypes, M.59.5028
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 105 (32)
Pycnodonte brongniarti, Bál86

- Leda (Lembulus) elata kiscelliensis* NOSZKY, 1939
holotype, M.59.5058
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 118 (53)
- Leda (Lembulus) gracilis* NOSZKY, 1939
holotype, M.59.5057
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 118 (52), pl. 2, fig. 5
- Leda modesta trigonalis* NOSZKY, 1939
holotype, M.59.5056
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 117 (52), pl. 2, fig. 4
Type missing
- Leda (Yoldia) perovalis pannonica* NOSZKY, 1939
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 118 (53)
Type probably destroyed in 1956
- Leda (Costatoleda) psammobiaeformis* TELELDI
ROTH, 1914
syntype, M.75.216
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 61 (52) (pl. 6, figs. 10-15)
Costatoleda psammobiaeformis, Bál86
- Leda (Costatoleda) psammobiaeformis prisca*
NOSZKY, 1939
holotype, M.59.5055
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 117 (51), pl. 2, fig. 9
Costatoleda psammobiaeformis prisca, Bál86
Type figured in BÁLDI 1986: pl. 3, fig. 28
- Leda (Yoldia) schréteri* NOSZKY, 1939
holotype, M.59.5059
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 118 (54), pl. 2, fig. 2

Lima cancellata angusta NOSZKY, 1939

syntypes, M.59.5041

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 111 (40)

Kiscellima cancellata, Bál86

Types figured in BÁLDI 1986: pl. 5, figs. 49,

50–52

Lima (Acesta) miocenica hantkeni NOSZKY, 1939

holotype, M.59.5042

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 111 (41), pl. 2, fig. 7

Lima (Acesta) miocenica hofmanni NOSZKY, 1939

holotype, M.59.5043

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 113 (43), pl. 2, fig. 25

Lima (Acesta) miocenica marginatostrigata NOSZKY, 1939

holotype, M.59.5044

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 113 (43), pl. 2, fig. 16

Limaria (Limatulella) langhiana zelenkai

BÁLDI, 1986

holotype, 2006.101.1.

Kiscellian, Kiscell Clay Formation

borehole RM-61, Recsk

Bál86, p. 78, pl. 5, fig. 53

Limaria (Limatulella) langhiana zelenkai

BÁLDI, 1986

paratype, 2006.102.1.

Kiscellian, Kiscell Clay Formation

borehole RM-28, Recsk

Bál86, p. 78, pl. 5, fig. 54

Limopsis retifera elongata NOSZKY, 1939

holotype, M.59.5049

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 116 (47)

Limopsis retifera jaworskii NOSZKY, 1939

holotype, M.59.5048

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 116 (47), pl. 2, fig. 6

Lucina (Megaxinus) bellardiana depressa NOSZKY, 1939

holotype, M.59.5027

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 121 (62), pl. 2, fig. 20

Lucina (Megaxinus) bellardiana depressa NOSZKY, 1939

paratype, M.61.5350

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 121 (62)

Lucina (Megaxinus) elliptica altialata NOSZKY, 1939

holotype, M.59.5024

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 121 (62), pl. 2, fig. 15

Lucina (Megaxinus) elliptica erecta NOSZKY, 1939

holotype, M.59.5018

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 121 (63)

Lucina (Myrthea) elongata NOSZKY, 1939

holotype, M.59.5017

Kiscellian, Kiscell Clay Formation

Holtzpach brickyard, Szépvölgy, Budapest

Nos39, p. 122 (64), pl. 2, fig. 26

Gibbolucina (Eomiltha) rectangulata, Bál86

Type figured in BÁLDI 1986: pl. 6, fig. 58

Lucina (Myrthea) elongata dregeri NOSZKY, 1939

holotype, M.59.5020

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos39, p. 122 (64), pl. 2, fig. 27

Gibbolucina (Eomiltha) rectangulata, Bál86

Paratypes under same inventory number.

Types figured in BÁLDI 1986: pl. 6, figs. 56, 57

Lucina globulosa kulcsári NOSZKY, 1939
holotype, M.59.5015
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 120 (60), pl. 2, fig. 28

Lucina (Dentilucina) meneghinii földvárii
NOSZKY, 1939
holotype, M.59.5025
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 121 (62), pl. 2, fig. 17

Lucina (Myrthea) rectangulata orbicularis
NOSZKY, 1939
syntype, M.59.5016
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 121 (64)

Lucina (Myrthea) spinifera lóczyi NOSZKY, 1939
holotype, M.59.5019
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 122 (65), pl. 2, fig. 21
Paratype under same inventory number

Lucina (Myrthea) spinifera peralta NOSZKY, 1939
holotype, M.59.5021
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 123 (65), pl. 2, fig. 18
Saxolucina heberti spissistriata, Bál86
Type figured in BÁLDI 1986: pl. 6, fig. 59

Lucina spinifera profunda NOSZKY, 1939
holotype, M.59.5023
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 132 (87), pl. 2, fig. 22
Saxolucina heberti spissistriata, Bál86
Type figured in BÁLDI 1986: pl. 6, fig. 62

Lucina (Myrthea) spinifera sinuosa NOSZKY, 1939
holotype, M.59.5022
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 123 (65), pl. 2, fig. 19

Lucina spissistriata peracuta NOSZKY, 1939
holotype, M.59.5026
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 120 (60), pl. 2, fig. 24

Lucina (Cardiolucina) striatula sandbergeri
NOSZKY, 1939
holotype, M.59.5014
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 120 (61), pl. 2, fig. 23

Lyonsia thraciaeformis NOSZKY, 1939
holotype, M.59.5082
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 129 (81), pl. 3, fig. 2

Malletia caterini convexa NOSZKY, 1939
holotype, M.59.5060
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 118 (54), pl. 2, fig. 14

Malletia harmati BÁLDI, 1986
holotype, 2006.100.1.
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 73, pl. 3, fig. 21

Malletia harmati BÁLDI, 1986
paratype, 2006.100.2.
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 73

Modiola mytiloides kovácsi NOSZKY, 1939
holotype, M.59.5045
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos39, p. 114 (44), pl. 1, fig. 17

Neaera (Cuspidaria) scalarina laevior NOSZKY,
1939
holotype, M.59.5083
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 129 (82)

- Neaera (Cuspidaria) sulcata* NOSZKY, 1939
holotype, M.59.5084
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 129 (82), pl. 3, fig. 18
Cuspidaria (Cuspidaria) rostrata, Bál86
Type figured as *Cuspidaria cuspidata* in
BÁLDI 1986: pl. 8, fig. 94
- Neaera (Tergulina) sulcosa* NOSZKY, 1939
holotype, M.59.5085
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 130 (83), pl. 3, fig. 20
Cuspidaria (Tergulina) sulcosa, Bál86
Type figured in BÁLDI 1986: pl. 9, fig. 99
- Nucula ledaeformis* NOSZKY, 1939
holotype, M.59.5051
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 116 (50), pl. 2, fig. 8
- Nucula paeelongata nysti* NOSZKY, 1939
holotype, M.59.5054
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 117 (51), pl. 2, fig. 3
- Nucula sulcifera peralta* NOSZKY, 1939
holotype, M.59.5053
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 117 (50), pl. 2, fig. 11
- Nucula sulcifera retifera* NOSZKY, 1939
holotype, M.59.5052
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 116 (50), pl. 2, fig. 10
- Nuculana solymarensis* BÁLDI, 1973
paratype, M.65.948
Egerian, Törökbálint Sandstone Formation
borehole Csordakút-5, Nagyegyháza, Óbarok
Bál73, p. 162, text-fig. 49.1.
- Nuculana solymarensis* BÁLDI, 1973
paratype, M.68.2088
Egerian, Törökbálint Sandstone Formation
borehole So-72, Solymár
Bál73, p. 162, text-fig. 49.1., pl. 2, fig. 5
- Panopaea (Glycymeris) kochi* NOSZKY, 1939
holotype, M.59.5076
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 127 (77), pl. 3, fig. 7
- Panopaea (Glycymeris) lőrentheyi* NOSZKY, 1939
holotype, M.59.5077
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 127 (78), pl. 3, fig. 1
- Panopaea (Pleuromya) vadászi* NOSZKY, 1939
holotype, M.59.5078
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 128 (78), pl. 3, fig. 8
Pholadomya vadászi, Bál86
Paratypes under same inventory number.
Types figured in BÁLDI 1986: pl. 8, figs. 85–87
- Pecchiolia argentea brachiumbonata* NOSZKY, 1939
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 123 (67), pl. 3, fig. 5
Type missing, not found in inventory.
Type figured in BÁLDI 1986: pl. 9, fig. 101
- Pecchiolia argentea brachiumbonata* NOSZKY, 1939
paratype, M.59.5063
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 123 (67)

- Pecten (Amusiopecten) burdigalensis minor*
TELEGDI ROTH, 1914
syntype, M.75.218
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 65 (55) (pl. 5, figs. 12–14)
Flabellipecten burdigalensis, Bál73
- Pecten (Aequipecten) schréteri* NOSZKY, 1936
holotype, M.59.2846
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 110 (84), pl. 5, fig. 3
Chlamys schreteri, C-M60
Type refigured in CSEPREGHY-MEZNERICS
1960: p. 30, pl. 20, fig. 9
- Perna lamarcki vendli* NOSZKY, 1939
holotype, M.59.5092
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 114 (45), pl. 1, fig. 15
- Perna maxillata schafarziki* NOSZKY, 1939
holotype, M.59.5046
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 114 (44), pl. 1, fig. 14
- Pholadomya (Procardia) canavarii hungarica*
NOSZKY, 1939
holotype, M.59.5080
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 128 (80), pl. 3, fig. 26
- Pholadomya (Procardia) canavarii rozloznskii*
NOSZKY, 1939
holotype, M.59.5079
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 128 (79), pl. 3, fig. 21
- Placenta saccoi* NOSZKY, 1939
holotype, M.59.5091
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 105 (29), pl. 1, fig. 13
- Portlandia korobkovi* BÁLDI, 1986
holotype, 2006.103.1.
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 73, pl. 3, fig. 26
- Portlandia korobkovi* BÁLDI, 1986
paratype, 2006.103.2.
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 73, pl. 3, fig. 27
- Portlandia korobkovi* BÁLDI, 1986
paratypes, 2006.103.3–6.
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 73
- Psammobia (Psammocula) tauroplana lata*
NOSZKY, 1939
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 125 (74)
Type probably destroyed in 1956
- Saxicava gaáli* NOSZKY, 1939
holotype, M.59.5075
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 127 (77), pl. 3, fig. 24
Thyasira vara, Bál86
- Saxicava rugosa medioumbonata* NOSZKY, 1939
holotype, M.59.5074
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 126 (77), pl. 3, fig. 28
- Semipecten mayeri franzenai* NOSZKY, 1939
holotype, M.59.5033
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 107 (35), pl. 1, fig. 8
Type missing

Semipecten spinicosta NOSZKY, 1939
 holotype, M.59.5034
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 108 (35), pl. 1, fig. 12
 Type missing

Siliqua harmati NOSZKY, 1939
 holotype, M.59.5070
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 125 (74), pl. 3, fig. 17
Aquinosiliqua harmati, Bál86
 Paratypes under same inventory number.
 Types figured in BÁLDI 1986: pl. 7, figs. 70-73

Siliqua harmati NOSZKY, 1939
 paratype, M.59.5073
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 125 (74)
Aquinosiliqua harmati, Bál86
 Type figured in BÁLDI 1986: pl. 7, fig. 74

Siliqua harmati compressa NOSZKY, 1939
 holotype, M.59.5072
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 126 (75), pl. 3, fig. 22
Aquinosiliqua harmati, Bál86

Siliqua harmati lata NOSZKY, 1939
 holotype, M.59.5071
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 126 (75), pl. 3, fig. 23
Aquinosiliqua harmati, Bál86
 Paratypes under same inventory number.
 Types figured in BÁLDI 1986: pl. 7, figs. 75-77

Sphaenia benghami szalaii NOSZKY, 1939
 holotype, M.59.5086
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 131 (84), pl. 3, fig. 10
Poromya szalaii, Bál86
 Type figured in BÁLDI 1986: pl. 8, fig. 93

Spondylus crenellaformis NOSZKY, 1939
 holotype, M.59.5031
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 106 (33), pl. 1, fig. 9
 Paratypes under same inventory number

Spondylus longispina NOSZKY, 1939
 holotype, M.59.5032
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 107 (34), pl. 1, fig. 16

Spondylus rogeri CSEPREGHY-MEZNERICS, 1961
 holotype, M.61.1
 Egerian, Eger Formation
 well of "Vincellér" school, Eger
 C-M61, p. 133, pl. 1, figs. 1, 2

Spondylus sulcosus NOSZKY, 1939
 holotype, M.59.5030
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 106 (33), pl. 1, fig. 10
Spondylus crenellaformis, Bál86

Spondylus tenuispina densicosta NOSZKY, 1939
 holotype, M.59.5029
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 106 (33)

Syndesmia elongata NOSZKY, 1939
 holotype, M.59.5177
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 133 (88), pl. 3, fig. 25
 Paratype under same inventory number

Tauraxinus budensis NOSZKY, 1939
 holotype, M.59.5062
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 119 (58), pl. 3, fig. 14

Tellina (Peronidea) budensis majzoni NOSZKY,
 1939
 holotype, M.59.5069
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 125 (73), pl. 3, fig. 16
 Paratypes under same inventory number

Tellina (Capsa) lamellosa NOSZKY, 1939
 holotype, M.59.5068
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos39, p. 125 (72), pl. 3, fig. 9

Tellina (Peronea) nitida recta NOSZKY, 1939
holotype, M.59.4277
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 125 (73), pl. 3, fig. 13

Teredo anguinea nodosa NOSZKY, 1939
holotype, M.59.5087
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 131 (85), pl. 3, fig. 27
Paratypes under same inventory number

Thracia teleldirothi NOSZKY, 1939
holotype, M.59.5081
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos39, p. 128 (80), pl. 3, fig. 6

Thyasira vara angusta BÁLDI, 1966
holotype, M.65.1079
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 85, pl. 1, fig. 4

Trinacria oligocaenica BÁLDI, 1986
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Bál86, p. 74, pl. 4, fig. 31
Type missing, not found in inventory

7.3. Oligocene Scaphopoda

Dentalium densitextum dejtarensense BÁLDI, 1973
holotype, M.68.2075
Egerian, Törökbálint Sandstone Formation
Dejtár
Bál73, p. 337, pl. 50, fig. 9

Dentalium haeringense densitexta NOSZKY, 1940
syntypes, M.59.5178
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 73 (51)
Dentalium densitextum, Bál73
Types figured in BÁLDI 1986: pl. 11, figs. 132–134

7.4. Oligocene Gastropoda

Acamptochetus clatratus BÁLDI, 1966
holotype, M.65.1089
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 89, pl. 3, fig. 11

Ampullina (Globularia) telegdirothi callosa
NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 102 (62), pl. 5, fig. 14
Globularia gibberosa callosa, Bál73
Type destroyed in 1956

Asthenotoma noszkyi BÁLDI, 1966
holotype, M.65.1095
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 92, pl. 3, fig. 13

Asthenotoma noszkyi BÁLDI, 1966
paratype, M.63.6614
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 92

Bittium reticulatum densespiratum BÁLDI, 1966
holotype, M.65.1084
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 87, pl. 2, fig. 3

Bittium spina agriense BÁLDI, 1966
holotype, M.65.1083
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 87, pl. 2, fig. 4

Bittium spina agriense BÁLDI, 1966
paratypes, M.83.73, M.83.77, M.83.79
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 87

Callistoma (Ampullo trochus) elegantulum
hegeduesi BÁLDI, 1973
holotype, M.68.2079
Egerian, Törökbálint Sandstone Formation
Dejtár
Bál73, p. 239, pl. 23, figs. 4–5
Paratype under same inventory number

*Calliostoma (Ampullotrochus) elegantulum
hegeduesi* BÁLDI, 1973
paratypes, M.68.2101, 2006.87.1.
Egerian, Törökbálint Sandstone Formation
Dejtár
Bál73, p. 239

Calyptera pseudodeformis BÁLDI, 1966
holotype, M.65.1085
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 88, pl. 2, figs. 6a–b

Calyptera pseudodeformis BÁLDI, 1966
paratype, M.65.1086
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 88

Cassidaria nodosa semicostata NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 102 (62), pl. 5, fig. 17
Cassidaria depressa, Bál73
Type destroyed in 1956

Charonia (Sassia) tarbelliana transiens BÁLDI,
1973
holotype, M.68.2095
Egerian, Eger Formation
Wind brickyard, Eger
Bál73, p. 282, pl. 35, figs. 6–7

Chenopus callosus TELELDI ROTH, 1914
syntypes, M.75.206
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 38 (32) (pl. 4, figs. 12–15)
Aporrhais callosa, Bál73

Chenopus callosus latialatus NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 102 (60)
Aporrhais callosa, Bál73
Type destroyed in 1956

Chenopus mathiasi NOSZKY, 1940
holotype
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos40, p. 59 (14), pl. 1, fig. 11
Type probably destroyed in 1956

Chenopus mathiasi retifera NOSZKY, 1940
holotype
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos40, p. 59 (14), pl. 1, fig. 10
Type probably destroyed in 1956

Chenopus speciosus digitata TELELDI ROTH, 1914
syntypes, M.75.205
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 38 (31) (pl. 4, figs. 7–8)
Drepanocheilus speciosus digitatus, Bál73

Chenopus uttingerianus éhiki NOSZKY, 1940
holotype, M.59.5109
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 60 (15), pl. 1, fig. 12
Paratypes under same inventory number

Chrysodomus dobói NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 103 (66), pl. 5, fig. 5
Pisanella doboi, Bál73
Type destroyed in 1956

Chrysodomus legányii NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 104 (66), pl. 5, fig. 8
Type destroyed in 1956

Clavella vighi NOSZKY, 1940
holotype, M.59.5143
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 67 (32), pl. 2, fig. 10

Cominella hungarica acuta NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 104 (67), pl. 6, fig. 4
 Type destroyed in 1956

Cominella hungarica elongata NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 104 (67), pl. 6, fig. 6
 Type destroyed in 1956

Cominella hungarica simplex NOSZKY, 1936
 syntypes
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 104 (67)
 Types destroyed in 1956

Conus (Leptoconus) dujardini brevispiratus
 NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 110 (81), pl. 5, fig. 13
Conus (Conolithus) dujardini egerensis, Bál73
 Type destroyed in 1956

Cythara (Margovoluta) bellardii laevis NOSZKY,
 1940
 holotype
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 62 (21), pl. 1, fig. 27
 Type probably destroyed in 1956

Daphnella franciscae NOSZKY, 1940
 holotype, M.59.5161
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 70 (43), pl. 2, fig. 22

Delphinula scobina basilaevia NOSZKY, 1940
 syntypes, M.59.5099
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 4

Dolichotoma collaris NOSZKY, 1940
 holotype, M.59.5163
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 70 (44), pl. 2, fig. 24
Epalxis (Bathytoma) collaris, Bál86
 Paratype under same inventory number. Types
 figured in BÁLDI 1986: pl. 11, figs. 123, 124

Dolichotoma collaris nudata NOSZKY, 1940
 syntypes, M.59.5164
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 44

Dolichotoma subdenticulata reticulata
 NOSZKY, 1940
 holotype, M.59.5162
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 70 (43), pl. 2, fig. 23

Drillia collectiva NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 106 (74), pl. 5, fig. 22
 Type destroyed in 1956

Drillia crispata oligocenica NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 107 (74), pl. 6, fig. 12
 Type destroyed in 1956

Drillia obtusa elongata NOSZKY, 1936
 holotype
 Egerian, Eger Formation
 Wind brickyard, Eger
 Nos36, p. 107 (74), pl. 5, fig. 18
 Type destroyed in 1956

Echinophoria intermedia andreatsi NOSZKY, 1940
 holotype, M.59.5120
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 62 (20), pl. 1, fig. 22

- Echinophoria intermedia retifera* NOSZKY, 1940
holotype, M.59.5121
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 20
- Echinophoria rondeletii longinodosa* NOSZKY, 1940
syntypes, M.59.5119
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 19
- Egerea collectiva nassaeformis* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 106 (73), pl. 5, fig. 12
Type destroyed in 1956
- Egerea collectiva pyruloides* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 106 (73)
Type destroyed in 1956
- Emarginula elongata raricosta* NOSZKY, 1940
holotype, M.59.5098
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 57 (4), pl. 1, fig. 9
- Eudolium fasciatum kiscelliensis* NOSZKY, 1940
holotype, M.59.5135
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 62 (22), pl. 1, fig. 26
Paratype under same inventory number
- Euthria kochi* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 103 (65), pl. 5, fig. 6
Type destroyed in 1956
- Euthriofusus szontaghi* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 105 (70), pl. 5, fig. 2
Type destroyed in 1956
- Fasciolaria jablonszkyi* NOSZKY, 1940
holotype, M.59.5141
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 66 (32), pl. 2, fig. 7
- Ficula condita longicauda* NOSZKY, 1940
holotype, M.59.5122
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 22
- Ficula tenuis arcuato-caudata* NOSZKY, 1940
holotype, M.59.5123
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (23), pl. 1, fig. 28
Paratypes under same inventory number
- Ficula tenuis effusa* NOSZKY, 1940
holotype, M.59.5124
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (23), pl. 1, fig. 29
Paratype under same inventory number
- Fusus (Aptyxis) lóczyi* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 105 (71), pl. 5, fig. 1
Aquilofusus loczyi, Bál73
Type destroyed in 1956
- Fusus (Aptyxis) lóczyi densicostatus* NOSZKY,
1936
syntypes
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 105 (71)
Aquilofusus loczyi, Bál73
Types destroyed in 1956

Fusus longiroster laevior NOSZKY, 1940
 syntypes, M.59.5140
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 30

Fusus pergracilis crassa NOSZKY, 1940
 holotype, M.59.5138
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 66 (29), pl. 2, fig. 5
 Paratypes under same inventory number

Fusus retrorsicosta longicauda NOSZKY, 1940
 holotype
 Kiscellian, Kiscell Clay Formation
 Szépvölgy, Budapest
 Nos40, p. 31
 Type destroyed in 1956

Fusus wenzi NOSZKY, 1940
 holotype
 Kiscellian, Kiscell Clay Formation
 Szépvölgy, Budapest
 Nos40, p. 66 (30), pl. 2, fig. 6
 Type destroyed in 1956

Galeodea echinophora clarae NOSZKY, 1940
 holotype, M.59.5117
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 61 (17), pl. 1, fig. 19

Galeodea echinophora margarethae NOSZKY, 1940
 holotype, M.59.5116
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 61 (17), pl. 1, fig. 17

Galeodea sconsoides luisae NOSZKY, 1940
 holotype, M.59.5115
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 61 (17), pl. 1, fig. 18

Galeodea taurinensis helenae NOSZKY, 1940
 holotype, M.59.5118
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 62 (18), pl. 1, fig. 21

Galeodocassis fuchsii saccoi NOSZKY, 1940
 holotype
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 61 (17), pl. 1, fig. 20
 Type probably destroyed in 1956

Gibbula dubia BÁLDI, 1973
 holotype, M.65.964
 Egerian, Törökbálint Sandstone Formation
 borehole N-3, Nagyegyháza, Óbarok
 Bál73, p. 242, pl. 23, fig. 3

Gibbula dubia BÁLDI, 1973
 paratype, M.65.964
 Egerian, Törökbálint Sandstone Formation
 borehole N-3, Nagyegyháza, Óbarok
 Bál73, p. 242, pl. 23, fig. 8

Globularia gibberosa callosa BÁLDI, 1964
 holotype, M.68.2096
 Egerian, Törökbálint Sandstone Formation
 borehole Dj-8, Jenői-tó, Diósjenő
 Bál64, p. 172, pl. 2, figs. 4a-b
Globularia gibberosa callosa (NOSZKY, 1936),
 Bál73
 Subsequently regarded identical to subspecies
 of same name of NOSZKY (1936)

Globularia gibberosa callosa BÁLDI, 1964
 paratypes, M.63.9474, M.63.9838, M.63.9839,
 M.63.9848, M.63.9849, M.97.190
 Egerian, Törökbálint Sandstone Formation
 borehole Dj-8, Jenői-tó, Diósjenő
 Bál64, p. 172
Globularia gibberosa callosa (NOSZKY, 1936),
 Bál73

Globularia ovata BÁLDI, 1963
 holotype, M.62.9365
 Egerian, Törökbálint Sandstone Formation
 outcrop T1, Törökbálint
 Bál63, p. 87, pl. 5, figs. 14a-b
Globularia rothi, Bál73

Globularia ovata BÁLDI, 1963
 paratype, M.62.9366
 Egerian, Törökbálint Sandstone Formation
 outcrop T1, Törökbálint
 Bál63, p. 87, pl. 5, fig. 13
Globularia rothi, Bál73

Harpa bellardii madáchi NOSZKY, 1940
 holotype, M.59.5145
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 67 (34), pl. 2, fig. 11

Hinia fortectorata edentata BÁLDI, 1966
 holotype, M.65.1091
 Egerian, Eger Formation
 Wind brickyard, Eger
 Bál66, p. 90, pl. 2, fig. 14

Hinia fortectorata edentata BÁLDI, 1966
 paratypes, M.83.64
 Egerian, Eger Formation
 Wind brickyard, Eger
 Bál66, p. 90

Hinia (Telasco) schlotheimi noszkyi BÁLDI, 1961
 holotype, M.60.10786
 Egerian, Eger Formation
 Wind brickyard, Eger
 Bál61, p. 101, pl. 4, fig. 6
Hinia schlotheimi, Bál73

Hinia (Telasco) schlotheimi noszkyi BÁLDI, 1961
 paratypes, M.60.10770
 Egerian, Eger Formation
 Nyárjas-tető, Novaj
 Bál61, p. 101, pl. 4, fig. 7
Hinia schlotheimi, Bál73

Jujubinus (Strigosella) multicingulatus praestrigosus BÁLDI, 1966
 holotype, M.64.104
 Egerian, Eger Formation
 Wind brickyard, Eger
 Bál66, p. 87, pl. 4, fig. 3

Latirus cognatus rarilamellosa NOSZKY, 1940
 holotype, M.59.5144
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 67 (33), pl. 2, fig. 9

Latrunculus (Peridipsaccus) caronis multisulcata
 NOSZKY, 1940
 holotype, M.59.5136
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 64 (26), pl. 1, fig. 25

Latrunculus (Peridipsaccus) eburnoides umbilicosiformis TELELDI ROTH, 1914
 syntypes, M.75.194
 Egerian, Eger Formation
 Wind brickyard, Eger
 Tel14a (Tel14b), p. 24 (20) (pl. 1, figs. 29–32)
Babylonia eburnoides umbilicosiformis, Bál73

Lithoconus ineditus acuticaudata NOSZKY, 1940
 syntypes, M.59.5173
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 48

Lithoconus ineditus parvispira NOSZKY, 1940
 syntypes, M.59.5172
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 48

Lyria collaris NOSZKY, 1940
 holotype, M.59.5147
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 68 (35), pl. 2, fig. 13

Lyria collaris szontaghi NOSZKY, 1940
 holotype, M.59.5148
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Nos40, p. 68 (35), pl. 2, fig. 14
 Paratype under same inventory number

Lyria taurinia mitraeformis NOSZKY, 1940
 holotype
 Kiscellian, Kiscell Clay Formation
 Szépvölgy, Budapest
 Nos40, p. 67 (35), pl. 2, fig. 12
 Type probably destroyed in 1956

Mangeliella (Enatoma) boggschi BÁLDI, 1961
 holotype, M.60.10755
 Egerian, Eger Formation
 Nyárjas-tető, Novaj
 Bál61, p. 107, pl. 4, figs. 17a–b

Marginella (Volvarina) vadászi BÁLDI, 1961
 holotype, M.60.10763
 Egerian, Eger Formation
 Nyárjas-tető, Novaj
 Bál61, p. 104, pl. 4, fig. 12

- Melanella naumannii depressosuturata* BÁLDI, 1966
holotype, M.65.1100
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 94, pl. 4, fig. 9
- Murex (Pteronotus) detritus venzoi* NOSZKY, 1940
holotype, M.59.5132
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 64 (28), pl. 2, fig. 2
Paratypes under same inventory number
- Melongena noszkyi* GAÁL, 1938
holotype, M.63.6483
Egerian, Törökbalint Sandstone Formation
Fehér-hegy, Balassagyarmat
Gaá38a, p. 53 (7), text-fig. 2
Galeodes basilica, Bál73
- Murex guembeli longispina* NOSZKY, 1940
holotype, M.60.4355
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 64 (27), pl. 1, fig. 31
- Metula reticulata incisa* NOSZKY, 1940
holotype, M.59.5142
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 67 (32), pl. 2, fig. 8
Paratypes under same inventory number
- Murex (Alipurpura) holocristatus* NOSZKY, 1940
holotype, M.59.5134
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 65 (28), pl. 2, fig. 3
- Murex (Phyllonotus) rufus michelotti* NOSZKY, 1940
holotype, M.59.5131
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 64 (28), pl. 1, fig. 32
- Murex (Haustellum) sismondae rarivaricosa* NOSZKY, 1940
holotype, M.59.5130
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 27
- Murex (Harmatia) stephani* NOSZKY, 1940
holotype, M.59.5137
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 65 (28), pl. 2, fig. 4
- Murex trigonalis nudus* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 88, pl. 3, fig. 7
- Murex (Antimitra) telegdirothi* BÁLDI, 1966
holotype, M.65.1096
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 92, pl. 3, fig. 10
- Murex (Foveomurex) trigonalis*, Bál73
Type destroyed in 1956
- Murex (Pteronotus) detritus bellardii* NOSZKY, 1940
holotype, M.59.5133
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 65 (28), pl. 2, fig. 1
- Nassa (Telasco) neuvillei elongata* NOSZKY, 1936
syntypes
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 105 (68)
- Hinia schlotheimi*, Bál73
Types destroyed in 1956

- Ocinebrina crassilabiata trivaricosa* BÁLDI, 1964
holotype, M.73.128
Egerian, Törökbálint Sandstone Formation
Sárisáp
Bál64, p. 173, pl. 2, figs. 6a–b
Inventory number quoted in BÁLDI 1973:
M. 55/730
- Ocinebrina crassilabiata trivaricosa* BÁLDI, 1964
paratype, M.63.9476
Egerian, Törökbálint Sandstone Formation
borehole Dj-3, Diósjenő
Bál64, p. 173
- Ocinebrina crassilabiata trivaricosa* BÁLDI, 1964
paratype, M.73.127
Egerian, Törökbálint Sandstone Formation
borehole Dj-1, Diósjenő
Bál64, p. 173, pl. 2, fig. 8
- Oliva (Olivella) inflata elongatula* NOSZKY, 1940
syntypes, M.59.5159
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 40
- Patella protea stredai* NOSZKY, 1940
holotype
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos40, p. 58 (8), pl. 1, fig. 7
Type probably destroyed in 1956
- Phos hevesensis* BÁLDI, 1966
holotype, M.65.1090
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 89, pl. 2, fig. 13
- Phos hevesensis* BÁLDI, 1966
paratypes, M.83.41
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 89
- Pleurotoma coronata erecta* NOSZKY, 1940
syntypes, M.59.5160
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 42
Turris (Gemmula) coronata, Bál73
- Pleurotoma duchatelli incostata* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 107 (75), pl. 5, fig. 19
Turris (Fusiturris) duchasteli, Bál73
Type destroyed in 1956
- Pleurotoma flexicostata ventricosa* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 107 (75), pl. 6, fig. 10
Turris (Fusiturris) duchasteli, Bál73
Type destroyed in 1956
- Pleurotoma rotata gracilis* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 107 (76), pl. 6, fig. 11
Type destroyed in 1956
- Pleurotomaria retifera* NOSZKY, 1940
holotype, M.59.5097
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 57 (3), pl. 1, fig. 1
Perotrochus budensis, Bál86
Type missing
- Pleurotomaria retifera* NOSZKY, 1940
paratype, M.59.4391
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos40, p. 57 (3)
Perotrochus budensis, Bál86
- Pleurotomaria sismondae densicosta* NOSZKY, 1940
syntypes, M.59.5096
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 3
Perotrochus budensis, Bál86
- Pseudotoma orbignyi cingulata* NOSZKY, 1940
holotype, M.59.5171
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 72 (47), pl. 2, fig. 29

- Pterocera (Alaria) rochatiana tergolaevis* NOSZKY, 1940
holotype, M.59.5139
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 60 (15), pl. 1, fig. 13
- Pugilina aequalis lathyroides* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 103 (65), pl. 6, fig. 13
Type destroyed in 1956
- Pyrula (Melongena) semseyiana* ERDŐS, 1900
holotype, M.69.226
Egerian, Törökbálint Sandstone Formation
Kő-hegy, Pomáz
Erd00, p. 298 (263), pl. 1
Galeodes semseyiana, Bál73
- Ranella (Apollon) gigantea budensis* NOSZKY, 1940
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (24), pl. 1, fig. 30
Type probably destroyed in 1956
- Ranella (Apollon) gigantea budensis* NOSZKY, 1940
paratypes, M.59.5126
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (24)
- Ranella (Apollon) gigantea rarinodosa* NOSZKY, 1940
holotype, M.59.5125
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 23
- Ranella (Apollon) semilaevis simplex* NOSZKY, 1940
holotype, M.59.5127
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 24
- Raphitoma pseudonassoides* BÁLDI, 1966
holotype, M.65.1099
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 94, pl. 3, fig. 16
- Raphitoma roemeri agriensis* BÁLDI, 1966
holotype, M.65.1098
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 93, pl. 3, fig. 15
- Raphitoma roemeri agriensis* BÁLDI, 1966
paratypes, M.83.46
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 93
- Raphitoma valdecarinata* BÁLDI, 1966
holotype, M.65.1097
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 93, pl. 3, fig. 1
- Rostellaria bicarinata* BÁLDI, 1966
holotype, M.65.1087
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 88, pl. 4, fig. 5
- Rostellaria bicarinata* BÁLDI, 1966
paratypes, M.83.57
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 88
- Rostellaria hungarica* NOSZKY, 1940
holotype, M.59.5110
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 60 (15), pl. 1, fig. 14
Tibia (Tibia) haueri hungarica, Bál86
Paratype under same inventory number,
figured in BÁLDI 1986: pl. 10, fig. 109
- Rostellaria hungarica* NOSZKY, 1940
paratypes, M.59.4994, M.59.5002, M.59.5003
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 60 (15)
Tibia (Tibia) haueri hungarica, Bál86

Rostellaria hungarica calcarata NOSZKY, 1940

holotype

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 61 (16), pl. 1, fig. 16

Tibia (Tibia) haueri hungarica, Bál86

Type probably destroyed in 1956

Rostellaria hungarica gracilis NOSZKY, 1940

holotype, M.59.5111

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 60 (16), pl. 1, fig. 15

Tibia (Tibia) haueri hungarica, Bál86

Type figured in BÁLDI 1986: pl. 10, fig. 110

Solarium mariae BÁLDI, 1961

holotype, M.60.10777

Egerian, Eger Formation

Nyárjas-tető, Novaj

Bál61, p. 97, pl. 4, figs. 10a-c

Architectonica mariae, Bál73

Styliola maxima raricostata NOSZKY, 1940

holotype, M.60.4374

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 73 (51), pl. 2, fig. 32

Praehyaloclysis raricostata, Zor97

Paratypes under same inventory number.

Types figured in ZORN & BOHN-HAVAS 1997:
pl. 1, figs. 3-4

Sulcogladus collegnroi acuta NOSZKY, 1940

syntypes, M.59.5112

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 16

Sulcogladus collegnroi brevis NOSZKY, 1940

syntypes, M.59.5114

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 16

Sulcogladus collegnroi elongata NOSZKY, 1940

syntypes, M.59.5113

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 16

Surcula beyrichi gracilis NOSZKY, 1936

holotype

Egerian, Eger Formation

Wind brickyard, Eger

Nos36, p. 108 (77), pl. 5, fig. 20

Turricula regularis, Bál73

Type destroyed in 1956

Surcula beyrichi simplex NOSZKY, 1936

holotype

Egerian, Eger Formation

Wind brickyard, Eger

Nos36, p. 108 (77), pl. 6, fig. 8

Turricula regularis, Bál73

Type destroyed in 1956

Surcula bulbosa NOSZKY, 1936

holotype

Egerian, Eger Formation

Wind brickyard, Eger

Nos36, p. 109 (79), pl. 6, fig. 2

Turricula regularis, Bál73

Type destroyed in 1956

Surcula effusa NOSZKY, 1940

holotype, M.59.5165

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 71 (45), pl. 2, fig. 25

Surcula lörentheyi NOSZKY, 1936

holotype

Egerian, Eger Formation

Wind brickyard, Eger

Nos36, p. 108 (77), pl. 5, fig. 16

Turricula regularis, Bál73

Type destroyed in 1956

Surcula moulinsii laevis NOSZKY, 1940

syntypes, M.59.5169

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 46

Surcula moulinsii raristriata NOSZKY, 1940

holotype, M.59.5166

Kiscellian, Kiscell Clay Formation

Újlaki brickyard, Kiscell, Budapest

Nos40, p. 71 (45), pl. 2, fig. 26

Turricula (Knefastia) leganyii, Bál86

Paratypes under same inventory number.

Types figured in BÁLDI 1986: pl. 11, figs. 127-129

- Surcula regularis arcuatospirata* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 109 (78), pl. 6, fig. 9
Turricula regularis, Bál73
Type destroyed in 1956
- Surcula regularis bulbiformis* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 109 (79), pl. 6, fig. 1
Turricula regularis, Bál73
Type destroyed in 1956
- Surcula regularis robusta* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 109 (78), pl. 6, fig. 3
Turricula regularis, Bál73
Type destroyed in 1956
- Surcula telegdirothi* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 108 (77), pl. 5, fig. 10
Turricula telegdirothi, Bál73
Type destroyed in 1956
- Syrnola laterariae* BÁLDI, 1966
holotype, M.65.1101
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 94, pl. 4, fig. 10
- Tectura tauroconica gradiformis* NOSZKY, 1940
holotype, M.59.5107
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 59 (9), pl. 1, fig. 8
Paratype under same inventory number
- Terebra simplex* TELELDI ROTH, 1914
syntypes, M.75.202
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 33 (28) (pl. 3, figs. 16-17)
- Theodoxus (Vittoclython) pilisensis* BÁLDI, 1973
holotype, M.65.793
Egerian, Törökbálint Sandstone Formation
borehole Leányfalu-1, Leányfalu
Bál73, p. 244, pl. 24, fig. 4
Paratype under same inventory number
- Theodoxus (Vittoclython) supraoligocaenicus*
BÁLDI, 1973
holotype, M.65.805
Egerian, Törökbálint Sandstone Formation
borehole Leányfalu-1, Leányfalu
Bál73, p. 245, pl. 26, fig. 1
- Theodoxus (Vittoclython) supraoligocaenicus*
BÁLDI, 1973
paratypes, M.65.805
Egerian, Törökbálint Sandstone Formation
borehole Leányfalu-1, Leányfalu
Bál73, p. 245, pl. 26, fig. 2
- Tinostoma egerensis* BÁLDI, 1966
holotype, M.65.1082
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 86, pl. 2, figs. 5a-b
Teinostoma egerensis, Bál73
- Triton euthiraformis* NOSZKY, 1940
holotype, M.59.5129
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (25), pl. 1, fig. 24
- Triton tudiclaeformis* NOSZKY, 1940
holotype, M.59.5128
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 63 (24), pl. 1, fig. 23
- Trochus (Solaria) oligoobscura* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 102 (56), pl. 6, fig. 5
Type destroyed in 1956
- Turbo (Bolma) muricata laevis* NOSZKY, 1940
syntypes, M.59.5100
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 5

Turbo (Senectus) ranellaformis NOSZKY, 1940
holotype, M.59.5101
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 57 (5), pl. 1, fig. 4
Paratypes under same inventory number

Turbo (Senectus) ranellaformis granulata
NOSZKY, 1940
holotype, M.59.5103
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 58 (6), pl. 1, fig. 6
Paratypes under same inventory number

Turbo (Senectus) ranellaformis peralta
NOSZKY, 1940
holotype, M.59.5102
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 58 (6), pl. 1, fig. 5

Turbo (Senectus) ranellaformis retifera
NOSZKY, 1940
syntypes, M.59.5104
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 6

Turbo (Senectus) ranellaformis simplex
NOSZKY, 1940
syntypes, M.59.5105
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 6

Turricula ilonae BÁLDI, 1966
holotype, M.65.1092
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 90, pl. 3, figs. 2–3

Turricula ilonae BÁLDI, 1966
paratypes, M.83.52
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 90

Turricula (Knefastia) legányii BÁLDI, 1966
holotype, M.65.1093
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 91, pl. 3, fig. 19

Turricula (Knefastia) legányii BÁLDI, 1966
paratypes, M.69.210
Egerian, Eger Formation
Wind brickyard, Eger
Bál66, p. 91, pl. 3, fig. 18

Turritella beyrichi percarinata TELELDI ROTH,
1914
syntypes, M.75.208
Egerian, Eger Formation
Wind brickyard, Eger
Tel14a (Tel14b), p. 45 (38) (pl. 3, fig. 20, pl. 4,
figs. 18–20)

Turritella sandbergeri margarethae GAÁL, 1938
syntypes, M.63.6459
Egerian, Törökbálint Sandstone Formation
Fehér-hegy, Balassagyarmat
Gaá38a, p. 51 (5)

Turritella (Haustator) venus margarethae, Bál61

Turritella (Haustator) strangulata alexandri
NOSZKY, 1940
holotype, M.60.4360
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 59 (13), pl. 1, fig. 2

Turritella (Haustator) strangulata bisulcata
NOSZKY, 1940
holotype, M.59.5108
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 59 (13), pl. 1, fig. 3

Umbrella mikszáthi NOSZKY, 1940
holotype, M.60.4356
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 72 (49), pl. 2, fig. 30

- Umbrella sancti-petri* NOSZKY, 1940
holotype, M.59.5175
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 72 (50), pl. 2, fig. 31
- Vaginella tricuspidata* ZORN & JANSSEN, 1993
paratypes, M.95.11
Egerian, Eger Formation
borehole Múcsony-136, Múcsony
Zor93, p. 63, pl. 1, figs. 1–7 (pl. 2, figs. 1–5,
pl. 3, figs. 1–4, pl. 4, figs. 1–5)
- Voluta (Lyria) gárdonyii* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 106 (72), pl. 5, fig. 15
Type destroyed in 1956
- Voluta suturalis undulatocollaris* NOSZKY, 1940
holotype, M.59.5146
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 34
- Volutilithes (Neoathleta) affinis effusa* NOSZKY,
1940
holotype, M.59.5155
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 38
- Volutilithes (Neoathleta) affinis wagneri* NOSZKY,
1940
holotype, M.59.5156
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 69 (38), pl. 2, fig. 20
Paratypes under same inventory number
- Volutilithes apenninica irregularis* NOSZKY, 1940
holotype, M.59.5149
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 68 (35), pl. 2, fig. 15
Paratypes under same inventory number
- Volutilithes (Neoathleta) consanguinea humilis*
NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 106 (72), pl. 5, fig. 11
Spelling of genus name and gender agreement
of species name incorrect in original
description. Type destroyed in 1956
- Volutilithes (Athleta) consanguinea székessyi*
NOSZKY, 1940
holotype
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 69 (37), pl. 2, fig. 19
Type destroyed in 1956
- Volutilithes elevata laevis* NOSZKY, 1940
holotype, M.59.5152
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 36
- Volutilithes (Athleta) ficalina bistriata* NOSZKY,
1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 106 (72), pl. 5, fig. 7
Athleta ficalina, Bál73
Genus name spelled incorrectly as *Volutilithes*
in original description. Type destroyed in 1956
- Volutilithes intercostaeciliata* NOSZKY, 1940
holotype, M.59.5153
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 69 (36), pl. 2, fig. 18
Paratypes under same inventory number
- Volutilithes multicostata krepuskai* NOSZKY, 1940
holotype, M.59.5151
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 69 (36), pl. 2, fig. 17
Paratypes under same inventory number

- Volutilithes (Neoathleta) obliqua altispirata* NOSZKY, 1940
holotype, M.59.5158
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 70 (39), pl. 2, fig. 21
- Volutilithes (Neoathleta) obliqua gracilior* NOSZKY, 1940
syntypes, M.59.5157
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 39
- Volutilithes (Athleta) pygmaea tenerinodosa* NOSZKY, 1940
holotype, M.59.5154
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 37
- Volutilithes rotaridesi* NOSZKY, 1940
holotype, M.59.5150
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Nos40, p. 68 (36), pl. 2, fig. 16
- Volutilithes rotaridesi* NOSZKY, 1940
paratypes, M.59.5150
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Nos40, p. 68 (36)
- Volutilithes telegyi* GAÁL, 1938
syntypes, M.63.6482
Egerian, Törökbálint Sandstone Formation
Fehér-hegy, Balassagyarmat
Athleta rarispina, Bál73
Gaá38a, p. 54 (9), text-fig. 3
Genus name spelled incorrectly as *Voluthilites* in original description
- Zebinella decussata curvicostata* NOSZKY, 1936
holotype
Egerian, Eger Formation
Wind brickyard, Eger
Nos36, p. 102 (56), pl. 5, fig. 21
Type destroyed in 1956

7.5. Oligocene Cephalopoda

- Nectoteuthis hungarica* KRETZOLI, 1942
holotype
Kiscellian, Kiscell Clay Formation
Csillaghegy, Budapest
Kre42a, p. 126, pl. 18
Type probably lost
- Sepia harmati* SZÖRÉNYI, 1933
lectotype, M.60.4371
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö33, p. 184, pl. 6, fig. 5 (pl. 6, figs. 4, 6)
Lectotype fixed and figured in WAGNER 1938: p. 180, text-fig. 1
- Sepia harmati* SZÖRÉNYI, 1933
paralectotypes, M.59.4655, M.59.4657,
M.59.4658, M.59.4660
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö33, p. 184
- Sepia harmati* SZÖRÉNYI, 1933
paralectotypes, M.59.4656
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö33, p. 184, pl. 6, fig. 10
- Sepia harmati* SZÖRÉNYI, 1933
paralectotype, M.59.4659
Kiscellian, Kiscell Clay Formation
Szépvölgy, Budapest
Szö33, p. 184
- Sepia kiscellensis* WAGNER, 1938
holotype, M.60.4372
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Wag38, p. 194 (183), text-fig. 2 (left side)
- Sepia kiscellensis* WAGNER, 1938
paratypes, M.59.4654
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Wag38, p. 194 (183)

7.6. Oligocene Cirripedia

- Scalpellum hungaricum* SZÖRÉNYI, 1934
syntype, M.59.5179
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö34, p. 273, pl. 17, fig. 9 (pl. 17, figs. 7, 8,
11, 13)
Carina
- Scalpellum hungaricum* SZÖRÉNYI, 1934
syntype, M.59.5179
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö34, p. 273, pl. 17, fig. 10
Tergum
- Scalpellum hungaricum* SZÖRÉNYI, 1934
syntype, M.59.5179
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö34, p. 273, pl. 17, fig. 12
Scutum
- Scalpellum lóczyi* SZÖRÉNYI, 1934
syntype, M.59.4699
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö34, p. 273, pl. 17, figs. 3a–b (pl. 17,
figs. 1, 2, 4, 5, 6, 14, 15)
- Scalpellum lóczyi* SZÖRÉNYI, 1934
syntype, M.59.4849
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Szö34, p. 273, pl. 17, figs. 16a–b

7.7. Oligocene Decapoda

- Calappa tridentata* BEURLEN, 1939
syntypes, M.59.4676, M.59.4679
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 150, text-fig. 5 (pl. 7, fig. 8)
- Calappa tridentata* BEURLEN, 1939
syntypes, M.59.4681
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 150, text-fig. 5, pl. 7, fig. 9

- Calappa tridentata* BEURLEN, 1939
syntype, M.60.758
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 150, text-fig. 5, pl. 7, fig. 10
- Callianassa brevimanus* BEURLEN, 1939
syntypes, M.59.4683
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 142, text-fig. 2, pl. 7, figs. 5, 6
- Callianassa brevimanus* BEURLEN, 1939
syntypes, M.59.4684, M.59.4685
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 142, text-fig. 2
- Callianassa nuda* BEURLEN, 1939
syntypes, M.59.4682, M.59.4686, M.59.4689,
M.59.4690, M.59.4691
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 144 (text-fig. 3, pl. 7, figs. 3, 4)
- Lyreidus hungaricus* BEURLEN, 1939
syntype, M.59.4687
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 146, text-fig. 4
- Lyreidus hungaricus* BEURLEN, 1939
syntype, M.59.4688
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 146, text-fig. 4, pl. 7, fig. 7
- Plagiolophus sulcatus* BEURLEN, 1939
holotype, M.59.4692
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 155, pl. 7, fig. 11
- Thaumastocheles rupeliensis* BEURLEN, 1939
syntypes, M.59.4693, M.59.4694, M.59.4697,
M.59.4700, M.59.4703, M.59.4704, M.59.4705,
M.59.4706, M.59.4707, M.59.4708, M.59.4709,
M.59.4712, M.66.961
Kiscellian, Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Beu39, p. 137, text-fig. 1

Thaumastocheles rupeliensis BEURLEN, 1939
 syntype, M.59.4696
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Beu39, p. 137, text-fig. 1, pl. 7, fig. 2

Thaumastocheles rupeliensis BEURLEN, 1939
 syntypes, M.59.4701
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Beu39, p. 137, text-fig. 1, pl. 7, fig. 1

7.8. Oligocene Ophiuroidea

Pseudaspidura hungarica KOLOSVÁRY, 1941
 holotype, M.59.4731
 Kiscellian, Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Kol41b, p. 307, text-figs. 1-3

Cristellaria pseudo-spinulosa FRANZENAU, 1894
 syntypes
 Badenian, Sámsonháza Formation
 Letkés
 Fra94a, p. 14, pl. 1, figs. 9a-b
 Types probably destroyed in 1956

Cymbaloporella tortonica FRANZENAU & MÁJZON,
 1956
 syntypes
 Badenian, Sámsonháza Formation
 Kemence
 Fra56, p. 217, pl. 1, figs. 4-6
 Types probably destroyed in 1956

Discorbis kerékhegyensis FRANZENAU, 1956
 holotype, M.61.4318
 Badenian, Sámsonháza Formation
 Kerék-hegy, Szob
 Fra56, p. 217, pl. 1, figs. 12-14

Fissurina szobensis FRANZENAU, 1956
 holotype, M.61.4319
 Badenian, Sámsonháza Formation
 Kerék-hegy, Szob
 Fra56, p. 217, pl. 1, figs. 21, 22

Frondicularia formosa FRANZENAU, 1894
 syntypes
 Badenian, Sámsonháza Formation
 Letkés
 Fra94a, p. 12, pl. 1, figs. 6a-b
 Types probably destroyed in 1956

Gaudryina kerékhegyensis FRANZENAU, 1956
 holotype, M.61.4317
 Badenian, Sámsonháza Formation
 Kerék-hegy, Szob
 Fra56, p. 217, pl. 1, fig. 38

Glandulina hantkeni FRANZENAU, 1894
 syntypes
 Pannonian, "reworked into compact sand
 (Congeria-beds) from older Miocene strata"
 Markuševac, Croatia
 Fra94b, p. 258
 Types probably destroyed in 1956, figured
 in FRANZENAU & MÁJZON 1956: p. 211, pl. 1,
 fig. 18

8. Miocene types

8.1. Miocene Foraminiferida

Bulimina scalena FRANZENAU, 1956
 syntypes
 Badenian, "calcareous clay"
 Zsupanek, Orşova (Nagyzsupány), Romania
 Fra56, p. 139 (108), pl. 1, figs. 10, 11
 Types probably destroyed in 1956

Cibicides kismartonensis FRANZENAU, 1956
 syntypes
 Badenian, Hartl Formation
 Hartelberg, Eisenstadt (Kismarton), Austria
 Fra56, p. 218, pl. 1, figs. 7-9
 Types probably destroyed in 1956

Cristellaria dicampyla FRANZENAU, 1894
 syntypes
 Badenian, Sámsonháza Formation
 Letkés
 Fra94a, p. 13, pl. 1, figs. 8a-b
 Types probably destroyed in 1956

- Glandulina hosiusii* FRANZENAU, 1894
 syntypes
 Pannonian, "reworked into compact sand (Congeria-beds) from older Miocene strata"
 Markuševac, Croatia
 Fra94b, p. 258
 Types probably destroyed in 1956, figured in FRANZENAU & MAJZON 1956: p. 212, pl. 1, fig. 19, name emended as *G. hosiusi*
- Glandulina schlichti* FRANZENAU, 1894
 syntypes
 Pannonian, "reworked into compact sand (Congeria-beds) from older Miocene strata"
 Markuševac, Croatia
 Fra94b, p. 259
 Types probably destroyed in 1956, figured in FRANZENAU & MAJZON 1956: p. 212, pl. 1, fig. 20
- Gyroidina laciniata* FRANZENAU, 1956
 syntypes
 Badenian, Hartl Formation
 Hartelberg, Eisenstadt (Kismarton), Austria
 Fra56, p. 219, pl. 1, figs. 46–48
 Types probably destroyed in 1956
- Hopkinsina franzenaui* MAJZON, 1956
 holotype
 Pannonian, "reworked into compact sand (Congeria-beds) from older Miocene strata"
 Markuševac, Croatia
 Fra56, p. 213, pl. 1, figs. 34, 35
 Type probably destroyed in 1956. Type excluded from *Uvigerina venusta* FRANZENAU 1894: p. 284, pl. 6, fig. 61
- Lagena pseudoacuticosta* FRANZENAU & MAJZON, 1956
 syntypes
 Badenian, Sámonháza Formation
 Kemence
 Fra56, p. 216, pl. 1, fig. 24
 Types probably destroyed in 1956
- Nodophthalmidium asperum* GÖRÖG, 1992
 holotype, 2004.54.1.
 Sarmatian, Kozárd Formation
 borehole Perbál-5, Perbál
 Gör92, p. 61, pl. 1, fig. 5
- Nodophthalmidium rugosum* GÖRÖG, 1992
 holotype, 2004.55.1.
 Sarmatian, Kozárd Formation
 borehole M-17, Mány
 Gör92, p. 62, pl. 1, fig. 3 (pl. 1, fig. 4)
- Nodosaria binominata* FRANZENAU, 1894
 syntypes
 Badenian, Sámonháza Formation
 Letkés
 Fra94a, p. 10, pl. 1, figs. 3a–b
Siphonodosaria scharbergana, Fra56
 Types probably destroyed in 1956
- Nodosaria letkésiensis* FRANZENAU, 1894
 syntypes
 Badenian, Sámonháza Formation
 Letkés
 Fra94a, p. 10, pl. 1, figs. 4a–b
 Types probably destroyed in 1956
- Nodosaria pertenuis* FRANZENAU, 1894
 syntypes
 Badenian, Sámonháza Formation
 Letkés
 Fra94a, p. 11, pl. 1, figs. 5a–b
 Types probably destroyed in 1956
- Nodosaria pseudo-scharbergana* FRANZENAU, 1894
 syntypes
 Pannonian, "reworked into compact sand (Congeria-beds) from older Miocene strata"
 Markuševac, Croatia
 Fra94b, p. 265
Siphonodosaria scharbergana, Fra56
 Types probably destroyed in 1956
- Plectina perturbata* FRANZENAU, 1956
 syntypes
 borehole Városliget, Budapest
 Fra56, p. 219, pl. 1, figs. 26, 26a
 Types probably destroyed in 1956

Quinqueloculina ermani trigonostomea

FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 98 (45), pl. 3, figs. 10–12

Types probably destroyed in 1956

Quinqueloculina krenneri FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 98 (46), pl. 3, figs. 13–18

Types probably destroyed in 1956

Quinqueloculina peregrina edentula FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 97 (45), pl. 3, figs. 4–6

Types probably destroyed in 1956

Quinqueloculina rákosiensis FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 98 (45), pl. 3, figs. 7–9

Types probably destroyed in 1956

Reophax incerta FRANZENAU, 1894

holotype

Badenian, Sámonháza Formation

Letkés

Fra94a, p. 7, pl. 1, figs. 2a–b

Type probably destroyed in 1956

Siphonaptera longidentata GÖRÖG, 1992

holotype, 2004.56.1.

Sarmatian, Kozárd Formation

borehole M-17, Mánya

Gör92, p. 66, pl. 2, fig. 8 (pl. 2, figs. 5–7)

Triloculina divaricata FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 91 (39), pl. 3, figs. 1–3

Types probably destroyed in 1956

Truncatulina letkésiensis FRANZENAU, 1894

syntypes

Badenian, Sámonháza Formation

Letkés

Fra94a, p. 16, pl. 1, figs. 11a–c

Types probably destroyed in 1956

Uvigerina venusta FRANZENAU, 1894

syntypes

Pannonian, “reworked into compact sand (Congeria-beds) from older Miocene strata”

Markuševac, Croatia

Fra94b, p. 284, pl. 6, figs. 60a–b, 61a–b

Types probably destroyed in 1956, figured in FRANZENAU & MAJZON 1956: p. 213, pl. 1, figs. 36, 37. Specimen of fig. 61 subsequently regarded as *Hopkinsina franzenaui* MAJZON (in FRANZENAU & MAJZON 1956: p. 213, pl. 1, figs. 34, 35)

Vertebralina foveolata FRANZENAU, 1881

syntypes

Badenian, Rákos Limestone Formation

railway cut, Rákos, Budapest

Fra81, p. 101 (49), pl. 3, figs. 19–21

Types probably destroyed in 1956

8.2. Miocene Polyplacophora

Cryptoplax margitae DULAI, 2001

holotype, M.99.113

Badenian, Sámonháza Formation?

borehole Szokolya-2, Szokolya

Dul01, p. 45, pl. 4, figs. 1–6

8.3. Miocene Bivalvia

Aequipecten opercularis hevesensis SCHRÉTER, 1929

lectotype, M.59.2843

Ottangian-Karpatian, Salgótarján Formation
mining colony, Egercsehi

ScZ29, p. 381

Chlamys opercularis hevesensis, C-M60

Right valve. Lectotype designated and
figured in CSEPREGHY-MEZNERICS 1960: p. 26,
pl. 17, fig. 11

- Aequipecten opercularis hevesensis* SCHRÉTER, 1929
lectotype, M.59.2843
Ottrangian–Karpatian, Salgótarján Formation
mining colony, Egercsehi
ScZ29, p. 381
Chlamys opercularis hevesensis, C-M60
Left valve. Lectotype designated and
figured in CSEPREGHY-MEZNERICS 1960: p. 26,
pl. 17, fig. 13
- Aequipecten opercularis hevesensis* SCHRÉTER, 1929
paralectotype, M.63.758
Ottrangian–Karpatian, Salgótarján Formation
mining colony, Egercsehi
ScZ29, p. 381
Chlamys opercularis hevesensis, C-M60
Type figured in CSEPREGHY-MEZNERICS 1960:
p. 26, pl. 17, fig. 12
- Aloidis carinata rostrata* CSEPREGHY-MEZNERICS, 1950
holotype
Badenian, Hidas Lignite Formation
Hidas
C-M50, p. 114 (87), pl. 6, figs. 6–7
Type destroyed in 1956
- Arca borsodensis* CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2113
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 102, pl. 7, fig. 15
- Arca (Anadara) hidasensis* STRAUSZ, 1950
holotype, M.62.546
Badenian, Hidas Lignite Formation
Hidas
Str50, p. 245 (239), text-figs. 1–2
- Arca (Barbatia) pseudobarbata* SZALAI, 1926
syntypes, M.60.5016
Badenian, Pusztamiske Formation
open pit mine, Várpalota
SzL26a, p. 341, text-figs. 1a–c
- Astarte solidula bukkense* CSEPREGHY-MEZNERICS, 1969
syntype, M.59.3416
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 106, pl. 7, fig. 17
- Astarte (Goodallia) triangularis* nigradensis
CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4356
Badenian, Sámonháza Formation?
Nógrádszakál
C-M54, p. 145 (78), pl. 11, fig. 1
Inventory number in original publication:
M. 52/1578
- Beguina (Carditamera) striatellata* cserhátensis
CSEPREGHY-MEZNERICS, 1954
syntypes, M.61.4357
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
C-M54, p. 145 (82), pl. 11, figs. 8–10
- Cardita (Cardiocardita) nigradensis* Cs.
MEZNERICS, 1950
syntype, M.61.4370
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
CsM50, p. 400, pl. 2, fig. 5
- Cardita (Cardiocardita) nigradensis* Cs.
MEZNERICS, 1950
syntypes, M.61.8576
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
CsM50, p. 400
- Cardium manyense* KÓKAY, 1967
holotype, M.66.963
Badenian, Pusztamiske Formation?
borehole M-6, Mány
Kók67, p. 88 (83), pl. 7, fig. 1
Type missing
- Chama bukkensis* CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2116
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 111, pl. 8, fig. 13
Paratypes under same inventory number

Chama bukkensis CSEPREGHY-MEZNERICS, 1969
 paratype, 2007.15.1.
 Badenian, Sámonháza Formation?
 Urasági szőlők, Borsodbóta
 C-M69, p. 111, pl. 8, fig. 12

Chama bukkensis CSEPREGHY-MEZNERICS, 1969
 paratypes, 2007.15.2-3.
 Badenian, Sámonháza Formation?
 Urasági szőlők, Borsodbóta
 C-M69, p. 111

Chama gryphina umbonata CSEPREGHY-
 MEZNERICS, 1969
 syntype, 2007.14.1.
 Badenian, Sámonháza Formation?
 Rendek-völgyi quarry, Borsodbóta
 C-M69, p. 110, pl. 8, figs. 18, 20

Chama gryphoides borsodensis CSEPREGHY-
 MEZNERICS, 1969
 holotype, 2007.16.1.
 Badenian, Sámonháza Formation?
 Urasági szőlők, Borsodbóta
 C-M69, p. 111, pl. 8, figs. 15, 19

Chlamys biaense CSEPREGHY-MEZNERICS, 1960
 holotype, M.59.2844
 Badenian, Rákos Limestone Formation
 large quarry, Bia, Biatorbágy
 C-M60, p. 28, pl. 19, fig. 3
 Right valve

Chlamys biaense CSEPREGHY-MEZNERICS, 1960
 holotype, M.59.2844
 Badenian, Rákos Limestone Formation
 large quarry, Bia, Biatorbágy
 C-M60, p. 28, pl. 19, fig. 5
 Left valve

Chlamys biaense CSEPREGHY-MEZNERICS, 1960
 paratype, M.63.748
 Badenian, Rákos Limestone Formation
 Bia, Biatorbágy
 C-M60, p. 28, pl. 19, fig. 6

Chlamys biaense CSEPREGHY-MEZNERICS, 1960
 paratype, M.63.749
 Badenian, Rákos Limestone Formation
 Bia, Biatorbágy
 C-M60, p. 28, pl. 19, fig. 4

Chlamys darnóensis CSEPREGHY-MEZNERICS,
 1960
 holotype, M.59.2847
 Eggenburgian, Pétervására Sandstone
 Formation
 Darnó-hegy, Recsk
 C-M60, p. 30, pl. 20, fig. 6
 Right valve

Chlamys darnóensis CSEPREGHY-MEZNERICS,
 1960
 holotype, M.59.2847
 Eggenburgian, Pétervására Sandstone
 Formation
 Darnó-hegy, Recsk
 C-M60, p. 30, pl. 20, fig. 10
 Left valve

Chlamys darnóensis CSEPREGHY-MEZNERICS,
 1960
 paratype, M.62.5435
 Eggenburgian, Pétervására Sandstone
 Formation
 Darnó-hegy, Recsk
 C-M60, p. 30

Chlamys palmata bipartita CSEPREGHY-
 MEZNERICS, 1960
 holotype, M.63.757
 Eggenburgian, Budafok Sand Formation
 Pacsirta-hegy, Budafok, Budapest
 C-M60, p. 36, pl. 33, figs. 8, 9 (pl. 33, fig. 10)
 Inventory number given erroneously in
 original publication as M.59.2849

Chlamys palmata bipartita CSEPREGHY-
 MEZNERICS, 1960
 paratype, M.59.2849
 Eggenburgian, Filakovo Formation
 Rapovce (Rapp), Slovakia
 C-M60, p. 36
 Original publication erroneously quotes this
 inventory number for holotype

Chlamys rákosense CSEPREGHY-MEZNERICS, 1960
 holotype, M.59.2845
 Badenian, Rákos Limestone Formation
 large quarry, Bia, Biatorbágy
 C-M60, p. 29, pl. 19, fig. 9
 Right valve

- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2845
Badenian, Rákos Limestone Formation
Rákos, Budapest
C-M60, p. 29, pl. 19, fig. 14
Left valve
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.744
Badenian, Rákos Limestone Formation
Bia, Biatorbágy
C-M60, p. 29, pl. 19, fig. 13
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.745
Badenian, Rákos Limestone Formation
Bia, Biatorbágy
C-M60, p. 29, pl. 19, fig. 12
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.750
Badenian, Rákos Limestone Formation
large quarry, Bia, Biatorbágy
C-M60, p. 29, pl. 19, fig. 15
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.753
Badenian, Rákos Limestone Formation
canal works, west side of Rákosfalva, Budapest
C-M60, p. 29, pl. 19, fig. 11
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.754
Badenian, Rákos Limestone Formation
Kerepesi út, Rákosfalva, Budapest
C-M60, p. 29, pl. 19, fig. 10
- Chlamys rákosense* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.3020
Badenian, Rákos Limestone Formation
large quarry, Fertőrákos
C-M60, p. 29, pl. 34, fig. 12
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2842
Karpatian, Fót Formation
Somló-hegy, Fót
C-M60, p. 21, pl. 13, fig. 2
Right valve
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratypes, M.62.5245, M.62.5250
Karpatian, Fót Formation
Szadai-szurdok, Fót
C-M60, p. 21
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratype, M.62.5246
Karpatian, Fót Formation?
Imreházapuszta, Dunakeszi
C-M60, p. 21
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratypes, M.62.5248
Karpatian, Egyházasgerge Formation
Öreghegy-tető, Csomád
C-M60, p. 21
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.747
Karpatian, Fót Formation
Somló-hegy, Fót
C-M60, p. 21, pl. 13, fig. 7
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.751
Karpatian, Fót Formation
Szadai-szurdok, Fót
C-M60, p. 21, pl. 13, fig. 3
- Chlamys scabrella hungarica* CSEPREGHY-MEZNERICS, 1960
paratype, M.63.755
Karpatian, Fót Formation
Szadai-szurdok, Fót
C-M60, p. 21, pl. 12, fig. 23

Chlamys scabrella hungarica CSEPREGHY-MEZNERICS, 1960
 paratype, M.63.756
 Karpatian, Egyházasgerge Formation
 Öreghegy-tető, Csomád
 C-M60, p. 21, pl. 13, fig. 6

Chlamys scabrella hungarica CSEPREGHY-MEZNERICS, 1960
 paratypes, M.63.2462, M.63.2478
 Karpatian, Fót Formation
 Somló-hegy, Fót
 C-M60, p. 21

Chlamys scabrella hungarica CSEPREGHY-MEZNERICS, 1960
 paratype, M.63.2488
 Karpatian, Egyházasgerge Formation
 Öreghegy-tető, Csomád
 C-M60, p. 21, pl. 13, fig. 8

Chlamys scabrella hungarica CSEPREGHY-MEZNERICS, 1960
 paratype, 2007.9.1.
 Karpatian, Egyházasgerge Formation
 Csomád
 C-M60, p. 21

Congeria soproniensis VITÁLIS, 1934
 syntype, M.57.872
 Pannonian, Szák Formation
 Hacker brickyard, Sopron
 Vit34, p. 518 (511) (pl. 1, figs. 1-5)

Congeria sümeghyi STRAUSZ, 1942
 syntypes, M.57.1740
 Pannonian, Somló Formation
 Nyárád
 Str42, p. 75, pl. 5, figs. 1-8

Gastrana tellinoides CSEPREGHY-MEZNERICS, 1969
 holotype, M.68.2117
 Badenian, Sámsonháza Formation?
 Rendek-völgyi quarry, Borsodbóta
 C-M69, p. 114, pl. 9, fig. 2

Limnocardium pappi STRAUSZ, 1940
 syntype, M.57.2007
 Pannonian, Algyő Formation
 borehole No. 1, Magyarszentmiklós
 Str40, p. 82 (86)

Limnocardium pappi STRAUSZ, 1940
 syntype, M.57.2008
 Pannonian, Algyő Formation
 borehole No. 1, Magyarszentmiklós
 Str40, p. 82 (86), text-fig. 1

Limnocardium praedesertum STRAUSZ, 1940
 syntype, M.57.2005
 Pannonian, Endrőd Formation
 borehole No. 2, Magyarszentmiklós
 Str40, p. 83 (86)

Limnocardium praedesertum STRAUSZ, 1940
 syntype, M.57.2006
 Pannonian, Endrőd Formation
 borehole No. 2, Magyarszentmiklós
 Str40, p. 83 (86), text-fig. 2

Mactra nögrádensis CSEPREGHY-MEZNERICS, 1951
 holotype
 Badenian, Sámsonháza Formation?
 Homokterenyé, Mátraterenyé
 C-M51, p. 317, pl. 13, fig. 9
 Type probably destroyed in 1956

Mactra nögrádensis CSEPREGHY-MEZNERICS, 1951
 paratype
 Badenian, Sámsonháza Formation?
 Homokterenyé, Mátraterenyé
 C-M51, p. 317, pl. 13, fig. 10
 Type probably destroyed in 1956

Miltha (Eomiltha) borsodensis CSEPREGHY-MEZNERICS, 1969
 holotype, M.68.2114
 Badenian, Sámsonháza Formation?
 Urasági szőlők, Borsodbóta
 C-M69, p. 109, pl. 8, fig. 2

- Modiolus excellens* CSEPREGHY-MEZNERICS, 1951
holotype, M.61.4354
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
C-M51, p. 315
Type figured in CSEPREGHY-MEZNERICS 1954:
p. 144 (67), pl. 8, fig. 14
- Modiolus excellens* CSEPREGHY-MEZNERICS, 1951
paratypes, M.61.4355
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
C-M51, p. 315
- Ostrea lamellosa exogyroides* CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4358
Badenian, Sámonháza Formation
Meszes-tető, Mátraverebély
C-M54, p. 144 (78), pl. 16, fig. 1
- Paphia waldmanni cserhátensis* Cs. MEZNERICS, 1950
syntype, M.61.4371
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
CsM50, p. 401, pl. 2, fig. 6
- Paphia waldmanni cserhátensis* Cs. MEZNERICS, 1950
syntypes, M.61.4372
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
CsM50, p. 401
- Pecten fótensis* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2839
Karpatian, Fót Formation
Somlyó-hegy, Fót
C-M60, p. 13, pl. 5, fig. 5 (pl. 5, fig. 4)
Right valve
- Pecten fótensis* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2839
Karpatian, Fót Formation
Somlyó-hegy, Fót
C-M60, p. 13, pl. 5, fig. 6
Left valve
- Pecten promontorensis* CSEPREGHY-MEZNERICS, 1960
holotype, M.59.2840
Eggenburgian, Budafok Sand Formation
Nagy-árok, Budafok, Budapest
C-M60, p. 13, pl. 6, fig. 3
- Pitaria lamarcki orientalis* CSEPREGHY-MEZNERICS, 1969
syntype, 2007.17.1.
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 112, pl. 9, fig. 13
- Solenocurtus antiquatus vindobonensis* CSEPREGHY-MEZNERICS, 1954
syntype, M.62.3025
Badenian, Sámonháza Formation
Márkháza
C-M54, p. 146 (100), pl. 13, fig. 12
- Sphenia minima* CSEPREGHY-MEZNERICS, 1950
holotype
Badenian, Hidas Lignite Formation
Hidas
C-M50, p. 112 (89), pl. 6, figs. 8–9
Type destroyed in 1956
- Venus chamaeformis* CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2115
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 113, pl. 9, figs. 10, 11
- Venus chamaeformis* CSEPREGHY-MEZNERICS, 1969
paratype, M.65.1032
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 113
- Venus chamaeformis* CSEPREGHY-MEZNERICS, 1969
paratype, 2007.18.1.
Badenian, Sámonháza Formation?
Borsodbóta
C-M69, p. 113

8.4. Miocene Gastropoda

Acirsa plana CSEPREGHY-MEZNERICS, 1950

holotype, M.63.7792

Badenian, Hidas Lignite Formation

MÁSZ drill core, Hidas

C-M50, p. 111 (33), pl. 2, fig. 6

Adeorbis ammonoides CSEPREGHY-MEZNERICS, 1950

holotype

Badenian, Hidas Lignite Formation

Hidas

C-M50, p. 111 (23), pl. 1, fig. 8

Type destroyed in 1956

Adeorbis várpalotensis SZALAI, 1926

syntypes, M.60.4682

Badenian, Pusztamiske Formation

Szabó sand pit, Várpalota

Szl26a, p. 344, text-figs. 5a-c

Cyclostrema varpalotensis, Str54

Alvania montagui trochiformis CSEPREGHY-MEZNERICS, 1956

holotype, M.61.4302

Badenian, Sámsonháza Formation

"large outcrop", Szob

C-M56, p. 443 (379), pl. 2, figs. 7-10

Inventory number in original publication:
M. 52/34

Alvania montagui trochiformis CSEPREGHY-MEZNERICS, 1956

paratypes, M.61.4304

Badenian, Sámsonháza Formation

"large outcrop", Szob

C-M56, p. 443 (379)

Bittium botense CSEPREGHY-MEZNERICS, 1969

holotype, M.68.2105

Badenian, Sámsonháza Formation?

Rendek-völgyi quarry, Borsodbóta

C-M69, p. 71, pl. 1, fig. 20

Bittium botense CSEPREGHY-MEZNERICS, 1969

paratypes, 2007.10.1-2.

Badenian, Sámsonháza Formation?

Rendek-völgyi quarry, Borsodbóta

C-M69, p. 71

Bittium botense CSEPREGHY-MEZNERICS, 1969

paratype, 2007.11.1.

Badenian, Sámsonháza Formation?

Mogyorós, Bükkmogyorós

C-M69, p. 71

Bittium leganyii CSEPREGHY-MEZNERICS, 1969

holotype, M.68.2104

Badenian, Sámsonháza Formation?

Királdi út, roadside, Borsodbóta

C-M69, p. 72, pl. 1, fig. 19

Locality on specimen label: Rendek-völgyi
quarry

Calyptraea kutassyi CSEPREGHY-MEZNERICS,
1969

holotype, M.68.2106

Badenian, Sámsonháza Formation?

Királdi út, roadside, Borsodbóta

C-M69, p. 76, pl. 2, figs. 19, 20

Clavatula (Surcula) consobrina badensis

CSEPREGHY-MEZNERICS, 1953

syntype, M.61.4296

Badenian, Sámsonháza Formation

Szob

C-M53, p. 11, pl. 1, figs. 7-8

Clavatula (Surcula) inermiformis CSEPREGHY-MEZNERICS, 1969

holotype, M.68.2112

Badenian, Sámsonháza Formation?

Urasági szőlők, Borsodbóta

C-M69, p. 96, pl. 6, fig. 7

Locality on specimen label: Gyükerfő, Balaton

Clavatula (Surcula) inermiformis CSEPREGHY-MEZNERICS, 1969

paratypes, M.68.2111

Badenian, Sámsonháza Formation?

Gyükerfői-dombo, Balaton

C-M69, p. 96, pl. 6, fig. 6

Locality on specimen label: Urasági szőlők,
Borsodbóta

Clavatula (Surcula) krenneri CSEPREGHY-

MEZNERICS, 1953

holotype, M.61.4299

Badenian, Sámsonháza Formation

"large outcrop", Szob

C-M53, p. 13, pl. 2, figs. 17-18

- Clavatula letkésensis* CSEPREGHY-MEZNERICS, 1953
holotype, M.61.4373
Badenian, Sámonháza Formation
Letkés
C-M53, p. 13, pl. 2, figs. 13–14
- Clavatula (Surcula) nodosa borsodensis* CSEPREGHY-MEZNERICS, 1969
holotype, M.70.619
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 96, pl. 6, fig. 5
- Clavatula nigradensis* CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4366
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
C-M54, p. 141 (51), pl. 7, fig. 13
- Clavatula orientoromana* BÁLDI, 1960
holotype, M.60.145
Badenian, Sámonháza Formation?
Szokolya
Bál60, p. 82, pl. 3, figs. 6a–b
- Clavatula orientoromana* BÁLDI, 1960
paratypes, M.60.146
Badenian, Sámonháza Formation?
Bíró-völgy, Szokolya
Bál60, p. 82
- Clavatula orientoromana* BÁLDI, 1960
paratype, M.60.147
Badenian, Sámonháza Formation?
Hagyigácsó, Szokolya
Bál60, p. 82
- Clavatula sublaevigata* BÁLDI, 1960
holotype, M.60.139
Badenian, Sámonháza Formation?
Bíró-völgy, Szokolya
Bál60, p. 80, pl. 3, figs. 1a–c
- Clavatula (Perrona) vindobonensis nodosa* CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4367
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
C-M54, p. 141 (52), pl. 7, figs. 8–9, 11–12
Inventory number in original publication:
M. 52/1435
- Clavatula (Perrona) vindobonensis nodosa* CSEPREGHY-MEZNERICS, 1954
paratypes, M.61.4368
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
C-M54, p. 141 (52)
- Clavus (Brachytoma) hungaricus* BÁLDI, 1960
holotype, M.60.156
Badenian, Sámonháza Formation?
Szokolya
Bál60, p. 84, pl. 3, figs. 4a–b
- Clavus (Brachytoma) hungaricus* BÁLDI, 1960
paratype, M.60.157
Badenian, Sámonháza Formation?
Hegypő, Szokolya
Bál60, p. 84
- Cythara (Mangelia) hontensis* CSEPREGHY-MEZNERICS, 1953
holotype, M.61.4298
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M53, p. 17, pl. 3, figs. 1–2
- Cythara (Mangelia) vulpecula polonica* CSEPREGHY-MEZNERICS, 1953
holotype, M.61.4297
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M53, p. 16, pl. 3, figs. 8–9
- Drillia granaria szobensis* CSEPREGHY-MEZNERICS, 1953
holotype, M.61.4295
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M53, p. 8, pl. 1, figs. 15–16
- Drillia noszkyi* Cs. MEZNERICS, 1950
syntype, M.61.4365
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
CsM50, p. 400, pl. 1, figs. 9–10
- Drillia noszkyi* Cs. MEZNERICS, 1950
syntypes, M.66.900
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
CsM50, p. 400

Erato laevis punctata CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4308
Badenian, Sámonháza Formation
Kerék-hegy, Szob
C-M56, p. 445 (395), pl. 4, figs. 4–5

Euthria subnodosa fortstriata CSEPREGHY-MEZNERICS, 1969
holotype, M.70.522
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 86, pl. 4, figs. 11, 13
Locality on specimen label: Gyükerfői-dombok,
Balaton

Galeodes (Volema) cornuta hungarica CSEPREGHY-MEZNERICS, 1950
holotype
Badenian, Hidas Lignite Formation
Hidas
C-M50, p. 113 (51), pl. 3, fig. 3
Type destroyed in 1956

Genota (Pseudotoma) bonellii botensis CSEPREGHY-MEZNERICS, 1969
syntype, M.70.642
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 98, pl. 6, fig. 23

Genota (Pseudotoma) bonellii botensis CSEPREGHY-MEZNERICS, 1969
syntypes, M.70.643
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 98

Genota (Pseudotoma) bonellii lapugensis CSEPREGHY-MEZNERICS, 1969
syntype, M.70.641
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 98, pl. 6, fig. 27

Genota (Pseudotoma) bonellii lapugensis CSEPREGHY-MEZNERICS, 1969
syntype, 2007.19.1.
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 98, pl. 6, fig. 24

Genota ramosa pupoides CSEPREGHY-MEZNERICS, 1969
holotype, M.70.639
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 99, pl. 6, fig. 19

Gibbula (Colliculus) biangula obtusata CSEPREGHY-MEZNERICS, 1969
syntype, M.70.374
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 67, pl. 1, figs. 1, 5, 8

Hastula hungarica CSEPREGHY-MEZNERICS, 1954
holotype, M.62.2971
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
C-M54, p. 142 (56), pl. 8, fig. 4
Inventory number in original publication:
M. 52/1442

Lathyrus crassus vindobonensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4315
Badenian, Sámonháza Formation
Letkés
C-M56, p. 447 (407), pl. 7, figs. 1–2

Lathyrus (Dolicolathyrus) hontensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4312
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M56, p. 448 (407), pl. 8, figs. 7–8
Inventory number in original publication:
M. 52/309

Lymnaea pachygaster nögrádensis CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4350
Karpatian, “freshwater limestone”
Krétabánya-völgy, Litke
C-M54, p. 143 (60), pl. 8, fig. 1
Inventory number in original publication:
M. 52/1432; originally quoted figure number
(fig. 7) is likely in error

Lymnaea pachygaster nögrádensis CSEPREGHY-MEZNERICS, 1954
paratypes, M.61.4351
Karpatian, “freshwater limestone”
Krétabánya-völgy, Litke
C-M54, p. 143 (60), pl. 8, fig. 7

Lymnaea pachygaster nogrädensis CSEPREGHY-MEZNERICS, 1954
 paratypes, M.61.8923, M.61.8926, M.61.8929,
 M.61.8930
 Karpatian, "freshwater limestone"
 Krétabánya-völgy, Litke
 C-M54, p. 143 (60)

Macularia lartetii grundensis CSEPREGHY-MEZNERICS, 1954
 syntype, M.61.4352
 Badenian, Sámonháza Formation
 Buda-hegy, Sámonháza
 C-M54, p. 143 (61), pl. 8, fig. 12

Macularia lartetii grundensis CSEPREGHY-MEZNERICS, 1954
 syntype, M.61.4353
 Badenian, Sámonháza Formation
 Buda-hegy, Sámonháza
 C-M54, p. 143 (61), pl. 8, fig. 13

Melania (Balanocochlis) incerta CSEPREGHY-MEZNERICS, 1969
 holotype, M.68.2103
 Badenian, Sámonháza Formation?
 Urasagi szőlők, Borsodbóta
 C-M69, p. 70, pl. 1, figs. 17, 18

Melanopsis vitálisi STRAUSZ, 1942
 holotype, M.57.1717
 Pannonian, Somló Formation
 Nyárád
 Str42, p. 92, pl. 5, figs. 14, 20, 29

Micromelania zalaensis STRAUSZ, 1940
 holotype, M.57.1994
 Pannonian, Endrőd Formation
 borehole No. 2, Magyarszentmiklós
 Str40, p. 84 (86), text-fig. 3

Mitra goniophora austriaca CSEPREGHY-MEZNERICS, 1950
 syntype
 Badenian, Hidas Lignite Formation
 Hidas
 C-M50, p. 56
 Type destroyed in 1956

Mitra goniophora transsylvaniaica CSEPREGHY-MEZNERICS, 1954
 syntype
 Badenian, Sámonháza Formation
 Buda-hegy, Sámonháza
 C-M54, p. 140 (47), pl. 6, figs. 1, 15
 Type missing, not found in inventory

Mitra goniophora transsylvaniaica CSEPREGHY-MEZNERICS, 1954
 syntype, M.61.4364
 Badenian, Sámonháza Formation
 Buda-hegy, Sámonháza
 C-M54, p. 140 (47), pl. 6, figs. 2, 16

Moniliopsis (Bathytoma) cataphracta orientalis CSEPREGHY-MEZNERICS, 1953
 syntype, M.61.4313
 Badenian, Sámonháza Formation
 Letkés
 C-M53, p. 16, pl. 3, figs. 19–20

Moniliopsis (Bathytoma) cataphracta orientalis CSEPREGHY-MEZNERICS, 1953
 syntypes, M.61.4314
 Badenian, Sámonháza Formation
 Letkés
 C-M53, p. 16

Murex (Muricantha) sedgwicki vindobonensis CSEPREGHY-MEZNERICS, 1950
 holotype
 Badenian, Hidas Lignite Formation
 Hidas
 C-M50, p. 112 (46), pl. 2, fig. 17
 Type destroyed in 1956

Myxas hidensis CSEPREGHY-MEZNERICS, 1950
 holotype
 Badenian, Hidas Lignite Formation
 Hidas
 C-M50, p. 113 (64), pl. 4, fig. 2
 Type destroyed in 1956

Nassa (?Tritia) borsodensis CSEPREGHY-MEZNERICS, 1969
 holotype, M.68.2109
 Badenian, Sámonháza Formation?
 Rendek-völgyi quarry, Borsodbóta
 C-M69, p. 89, pl. 5, figs. 13, 14

Nassa (Hima) franzenaui CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4311
Badenian, Sámsonháza Formation
“large outcrop”, Szob
C-M56, p. 447 (405), pl. 7, figs. 32–33
Inventory number in original publication:
M. 52/304

Nassa (Uzita) hontensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4310
Badenian, Sámsonháza Formation
“large outcrop”, Szob
C-M56, p. 446 (405), pl. 7, figs. 24–25
Inventory number in original publication:
M. 52/303

Nassa (?Leiodomus) sturi spiralostrigata CSEPREGHY-MEZNERICS, 1969
syntype, M.59.7778
Badenian, Sámsonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 89, pl. 5, figs. 11, 12

Nassa (?Leiodomus) sturi spiralostrigata CSEPREGHY-MEZNERICS, 1969
syntype, M.59.7909
Badenian, Sámsonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 89

Nassa (Hima) turbinella semilaevis CSEPREGHY-MEZNERICS, 1969
syntype, M.70.538
Badenian, Sámsonháza Formation?
Királdi út, roadside, Borsodbóta
C-M69, p. 87, pl. 5, fig. 7

Nassa (Hima) turbinella semilaevis CSEPREGHY-MEZNERICS, 1969
syntype
Badenian, Sámsonháza Formation?
Királdi út, roadside, Borsodbóta
C-M69, p. 87, pl. 5, fig. 6
Type missing, not found in inventory

Odontostomia hontensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4305
Badenian, Sámsonháza Formation
Szob
C-M56, p. 444 (393), pl. 3, figs. 33–34
Inventory number in original publication:
M. 52/164

Odontostomia hontensis CSEPREGHY-MEZNERICS, 1956
paratype, M.61.4306
Badenian, Sámsonháza Formation
Szob
C-M56, p. 444 (393)

Odontostomia szobensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4307
Badenian, Sámsonháza Formation
“large outcrop”, Szob
C-M56, p. 445 (393), pl. 3, figs. 38–41
Inventory number in original publication:
M. 52/448

Olivella (Lamprodoma) clavula vindobonensis CSEPREGHY-MEZNERICS, 1954
syntype, M.61.4360
Badenian, Sámsonháza Formation
Márkháza
C-M54, p. 139 (44), pl. 6, figs. 3, 9

Philbertia hungarica CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4369
Badenian, Sámsonháza Formation
Buda-hegy, Sámsonháza
C-M54, p. 142 (54), pl. 7, figs. 16–17, 21–22
Inventory number in original publication:
M. 52/1441

Pirenella picta pseudogamlitzenis STRAUSZ, 1955
holotype
Badenian, Hidas Lignite Formation
Hidas
Str55, p. 167 (46), text-fig. 7, pl. 4, fig. 32 (pl. 4,
fig. 31)
Type probably destroyed in 1956

Pisanianura gaboraroni CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2108
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 85, pl. 4, figs. 17, 18

Pithocerithium pseudobliquistoma SZALAI, 1926
syntypes, M.60.4789
Badenian, Pusztamiske Formation
Szabó sand pit, Várpalota
Szl26a, p. 343, text-figs. 2-4
Cerithium (Vulgocerithium) pseudobliquistoma,
Str54

Plesiothyreus leganyii CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2102
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 68, pl. 1, figs. 12, 16

Pusionella pseudofusus palatinus STRAUSZ, 1954
syntypes, M.60.4948, M.60.4949, M.60.4950,
M.60.4951, M.60.4952
Badenian, Pusztamiske Formation
sand pit, Várpalota
Str54, p. 110 (32), pl. 4, figs. 75a-d

Pyrene (Anachis) bellardii grussbachensis
CSEPREGHY-MEZNERICS, 1969
syntypes, M.70.503
Badenian, Sámonháza Formation?
Gyükerfői-dombok, Balaton
C-M69, p. 83

Pyrene (Anachis) bellardii grussbachensis
CSEPREGHY-MEZNERICS, 1969
syntype, M.70.504
Badenian, Sámonháza Formation?
Gyükerfő, Balaton
C-M69, p. 83, pl. 4, figs. 1, 5

Pyrene (Anachis) guembeli buekkense CSEPREGHY-MEZNERICS, 1969
syntype, M.59.7911
Badenian, Sámonháza Formation?
Királdi út, road cut, Borsodbóta
C-M69, p. 82, pl. 4, fig. 24

Pyrene (Anachis) guembeli buekkense CSEPREGHY-MEZNERICS, 1969
syntype, M.70.506
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 82, pl. 4, fig. 19

Rissoa hidensis CSEPREGHY-MEZNERICS, 1950
holotype
Badenian, Hidas Lignite Formation
Hidas
C-M50, p. 110 (22), pl. 1, fig. 2
Type destroyed in 1956

Rissoina (Zebinella) nágrádensis CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4359
Badenian, Sámonháza Formation
Halastóhegy, Sámonháza
C-M54, p. 139 (15), pl. 1, fig. 20

Roualtia lapugensis badensis CSEPREGHY-MEZNERICS, 1969
syntypes, M.70.627
Badenian, Sámonháza Formation?
Gyükerfői-dombok, Balaton
C-M69, p. 99

Roualtia lapugensis badensis CSEPREGHY-MEZNERICS, 1969
syntype, 2007.12.1.
Badenian, Sámonháza Formation?
Gyükerfő, Balaton
C-M69, p. 99, pl. 6, fig. 13

Roualtia lapugensis badensis CSEPREGHY-MEZNERICS, 1969
syntype, 2007.13.1.
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 99

Sabatia (Mnestia) pannonica CSEPREGHY-MEZNERICS, 1952
holotype, M.61.4301
Badenian, Sámonháza Formation
"large outcrop", Szob
C-M52, p. 229, pl. 14, figs. 15-17

Sigaretus borsodensis CSEPREGHY-MEZNERICS, 1969
holotype, M.68.2107
Badenian, Sámonháza Formation?
Urasági szőlők, Borsodbóta
C-M69, p. 77, pl. 2, figs. 17, 18

Taurasia szobensis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4309
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M56, p. 445 (402), pl. 6, figs. 1–4
Inventory number in original publication:
M. 52/253

Theodoxus (Calvertia?) grateloupianus dentatus
KÓKAY, 1967
holotype, M.66.964
Badenian, Pusztamiske Formation?
bentonite quarry, Bánd
Kók67, p. 89 (83), pl. 8, fig. 1
Type missing

Trochocerithium hungaricum CSEPREGHY-MEZNERICS, 1952
holotype, M.61.4300
Badenian, Sámonháza Formation
“large outcrop”, Szob
C-M52, p. 227, pl. 14, figs. 1, 2, 5, 7, 8, 11

Trophon varicosissimus spiniferus CSEPREGHY-MEZNERICS, 1969
syntype, M.70.488
Badenian, Sámonháza Formation?
Uszófői-dombok, Csernely
C-M69, p. 81, pl. 4, fig. 3
Locality on specimen label: Csermely

Trophon varicosissimus spiniferus CSEPREGHY-MEZNERICS, 1969
syntype, M.70.489
Badenian, Sámonháza Formation?
Rendek-völgyi quarry, Borsodbóta
C-M69, p. 81, pl. 4, figs. 2, 4
Locality on specimen label: Uszófői-dombok,
Csermely

Turritella incisaformis CSEPREGHY-MEZNERICS, 1956
holotype, M.61.4303
Badenian, Sámonháza Formation
beside road to Ipolydamásd, Szob
C-M56, p. 444 (384), pl. 2, fig. 36
Inventory number in original publication:
M. 52/66

Turritella incisaformis CSEPREGHY-MEZNERICS, 1956
paratypes, M.61.4494
Badenian, Sámonháza Formation
Szob
C-M56, p. 444 (384)

Valvata hidensis KÓKAY, 1967
holotype, M.66.965
Badenian, Hidas Lignite Formation
Hidas
Kók67, p. 90 (84), pl. 8, fig. 5
Type missing

Vexillum (Costellaria) harmati CSEPREGHY-MEZNERICS, 1954
holotype, M.61.4361
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
C-M54, p. 140 (45), pl. 5, figs. 29, 31–33

Vexillum (Costellaria) harmati CSEPREGHY-MEZNERICS, 1954
paratypes, M.61.4362
Badenian, Sámonháza Formation
Buda-hegy, Sámonháza
C-M54, p. 140 (45)

8.5. Miocene Annelida

Ditrupa transsilvanica MEZNERICS, 1944
holotype, M.60.399
Badenian, Dej Formation?
Lăpușiu (Lapugy), Romania
Mez44b, p. 45, pl. 2, fig. 8

Ditrupa transsilvanica MEZNERICS, 1944
paratype, M.60.400
Badenian, Dej Formation?
Coștei (Kostej), Romania
Mez44b, p. 45, pl. 2, fig. 7

Ditrupa transsilvanica MEZNERICS, 1944
paratype
Badenian, Dej Formation?
Lăpușiu (Lapugy), Romania
Mez44b, p. 45, pl. 2, fig. 9
Type probably lost

Ditrupa transsilvanica MEZNERICS, 1944
 paratype
 Badenian, Dej Formation?
 Coștei (Kostej), Romania
 Mez44b, p. 45, pl. 2, fig. 6
 Type probably lost

Balanus amphitrite archi-inexpectatus
 KOLOSVÁRY, 1948
 syntypes
 Badenian, Pusztaiske Formation
 Szabó sand pit, Várpalota
 Kol48, p. 111 (108), text-fig. 7
 Types probably destroyed in 1956

8.6. Miocene Brachiopoda

Hemithiris acuta MEZNERICS, 1944
 holotype, M.62.957
 Karpatian, Fót Formation?
 Csömör
 Mez44a, p. 22, pl. 3, figs. 9, 10
Aphelesia acuta, Bit04

Balanus amphitrite helenae KOLOSVÁRY, 1949
 syntypes, M.57.3463
 Eggenburgian, Budafok Sand Formation
 Pacsirta-hegy, Budafok, Budapest
 Kol49a, p. 117, text-fig. 6

Hemithiris acuta MEZNERICS, 1944
 paratypes, M.62.956
 Karpatian, Fót Formation?
 Csömör
 Mez44a, p. 22
Aphelesia acuta, Bit04

Balanus amphitrite litoralis KOLOSVÁRY, 1948
 syntypes
 Badenian, Pusztaiske Formation
 Szabó sand pit, Várpalota
 Kol48, p. 110 (106), text-fig. 5
 Types probably destroyed in 1956

Hemithiris acuta MEZNERICS, 1944
 paratype, M.61.4375
 Karpatian, Fót Formation?
 Csömör
 Mez44a, p. 22, pl. 3, figs. 7, 8
Aphelesia acuta, Bit04

Balanus hungaricus KOLOSVÁRY, 1941
 syntypes, M.62.6046
 Eggenburgian, Budafok Sand Formation
 Pacsirta-hegy, Budafok, Budapest
 Kol41a, p. 282, text-figs. 1-7

Terebratulina plana MEZNERICS, 1944
 holotype, M.62.1006
 Badenian, Dej Formation
 Aghireșu (Egeres), Romania
 Mez44a, p. 35, pl. 2, figs. 10, 11

Balanus improvisus fossilis KOLOSVÁRY, 1942
 syntypes, M.62.6042
 Eggenburgian, Budafok Sand Formation
 Pacsirta-hegy, Budafok, Budapest
 Kol42, p. 204, text-figs. 1-2

8.7. Miocene Cirripedia

Andromacheia noszkyi KOLOSVÁRY, 1949
 holotype
 Sarmatian, Kozárd Formation?
 Magyarszék
 Kol49a, p. 114, text-fig. 4
 Type probably destroyed in 1956

Balanus legányii KOLOSVÁRY, 1950
 syntypes, M.57.3402
 Karpatian, Egyhásasgerge Formation
 Bő-völgy, Felsőtárkány
 Kol50, p. 275 (272), text-fig. 2

Balanus amphitrite abundantus KOLOSVÁRY, 1948
 syntypes
 Badenian, Pusztaiske Formation
 Szabó sand pit, Várpalota
 Kol48, p. 111 (106), text-fig. 6
 Types probably destroyed in 1956

Balanus legányii KOLOSVÁRY, 1950
 syntypes, M.63.5819
 Karpatian, Egyhásasgerge Formation
 Felsőtárkány
 Kol50, p. 275 (272), text-fig. 2

Balanus pannonicus KOLOSVÁRY, 1952
 syntypes
 Badenian, Sámsonháza Formation
 Kálvária-hegy, Kemence
 Kol52, p. 234, text-figs. 1a-b
 Types probably destroyed in 1956

Balanus tintinnabulum honti KOLOSVÁRY, 1950
holotype, M.57.3508
Badenian, Sámonháza Formation
Honti-szakadék, Hont
Kol50, p. 275 (271), text-fig. 1

Balanus tintinnabulum vadászi KOLOSVÁRY, 1948
syntypes
Badenian, Pusztamiske Formation
Szabó sand pit, Várpalota
Kol48, p. 110 (104), text-fig. 1
Types probably destroyed in 1956

Balanus transsylvanicus KOLOSVÁRY, 1950
holotype, M.57.3445
Badenian, Dej Formation?
Coștei (Kostej), Romania
Kol50, p. 276 (273), text-fig. 4
Type missing

Creusia spinulosa cladangiae KOLOSVÁRY, 1949
syntypes
Badenian, Sámonháza Formation
Halastó, Sámonháza
Kol49a, p. 112, text-fig. 1
Types probably destroyed in 1956

Creusia spinulosa praespinulosa KOLOSVÁRY, 1949
syntypes, M.57.3378, M.57.3382, M.57.3407,
M.57.3408, M.57.3409
Badenian, Rákos Limestone Formation?
Pécsbudafa, near cemetery, Mánfa
Kol49a, p. 113, text-figs. 2, 3, 5

8.8. Miocene Decapoda

Achaeus magnus MÜLLER, 1978
holotype, M.86.81
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 288 (279), pl. 15, figs. 1-2
Author's inventory number: MOE-5-1;
paratypes under same inventory number
(pl. 15, figs. 3-4)

Actaea turcocampestris MÜLLER, 1984
holotype, M.86.340
Badenian, Rákos Limestone Formation
borehole FRK-21, Fertőrákos
Mül84, p. 87, pl. 74, figs. 1-2
Author's inventory number: MFR-1-1

Actaea turcocampestris MÜLLER, 1984
paratypes, M.86.118
Badenian, Sámonháza Formation
Törökmező, Nagymaros
Mül84, p. 87, pl. 74, figs. 3-5
Author's inventory number: MTZ-10

Anapagurus marginatus MÜLLER, 1978
holotype, M.86.70
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 286 (278), pl. 4, fig. 1
Author's inventory number: MOE-3-1;
paratypes under same inventory number
(pl. 4, figs. 2-3)

Anapagurus miocenicus MÜLLER, 1978
holotype, M.86.252
Badenian, Rákos Limestone Formation
Gyakorló út, Budapest
Mül78, p. 286 (277), pl. 6, figs. 2-5
Author's inventory number: MGY-1-1;
paratypes under same inventory number

Asthenognathus rakosensis MÜLLER, 2006
holotype, 2005.100.1.
Badenian, Rákos Limestone Formation
Rákos, Budapest
Mül06, p. 43, pl. 1, fig. 6
Author's inventory number: MRF-15

Brachynotus februarius MÜLLER, 1974
holotype, 2004.157.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 127 (123), pl. 5, fig. 2
Author's inventory number: MB-6-1

Brachynotus februarius MÜLLER, 1974
paratype, M.86.37
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 127 (123)
Author's inventory number: MKF-5

Brachynotus februarius MÜLLER, 1974
paratype, M.86.41
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 127 (123)
Author's inventory number: MM-4

- Brachynotus februarius* MÜLLER, 1974
 paratypes, M.86.523
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül74a, p. 127 (123), pl. 5, figs. 3–4
 Author's inventory number: MB-6
- "*Callianassa*" *kerepeziensis* MÜLLER, 1976
 holotype, M.86.412
 Badenian, Rákos Limestone Formation
 Kerepezi út, Budapest
 Mül76, p. 154 (149), pl. 1, figs. 2–4
 Author's inventory number: MKC-2-1;
 paratypes under same inventory number
 (pl. 1, figs. 1, 5)
- "*Callianassa*" *kerepeziensis* MÜLLER, 1976
 paratypes, M.86.368
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül76, p. 154 (149)
 Author's inventory number: MA-2
- ?*Callianassa szobensis* MÜLLER, 1984
 holotype, 2004.158.1.
 Badenian, Sámonháza Formation
 Damásdi-patak, Szob
 Mül84, p. 53, pl. 7, figs. 3–4
 Author's inventory number: MSZ-1-1
- "*Carcinus*" *lörentheyi* MÜLLER, 1974
 holotype, 2006.85.1.
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül74a, p. 126 (122), pl. 2, figs. 4–5
Pirimela lorentheyi, Mül84
 Author's inventory number: MB-4-1
- Charybdis (Goniosupradens) mathiasi* MÜLLER,
 1984
 holotype, M.86.498
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 81, pl. 63, figs. 1, 4
 Author's inventory number: MRZ-5-1;
 paratypes under same inventory number
 (pl. 63, figs. 2–3, pl. 64, figs. 1–5, pl. 67, figs. 3–5)
- Charybdis (Goniosupradens) mathiasi* MÜLLER,
 1984
 paratype, M.86.349
 Badenian, Sámonháza Formation
 Fekete-hegy, Visegrád
 Mül84, p. 81
 Author's inventory number: MV-7
- Charybdis (Goniosupradens) mathiasi* MÜLLER,
 1984
 paratypes, 2007.98.1–3., 2007.99.1.
 Badenian, Rákos Limestone Formation
 sand pit, Diósd
 Mül84, p. 81
 Author's inventory number: MDZ-6
- Chlorodiella juglans* MÜLLER, 1984
 holotype, M.89.63
 Badenian, Rákos Limestone Formation
 sand pit, Diósd
 Mül84, p. 89, pl. 78, figs. 5–6
 Author's inventory number: MDZ-11-1;
 paratypes under same inventory number
- Chlorodiella loczyi* MÜLLER, 1984
 holotype, M.86.204
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 89, pl. 78, figs. 1–3
 Author's inventory number: MRZ-7-1;
 paratypes under same inventory number
 (pl. 78, fig. 4)
- Chlorodiella mediterranea tetenyensis* MÜLLER,
 1984
 holotype, M.86.205
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 88, pl. 77, fig. 5
 Author's inventory number: MRZ-19-1;
 paratypes under same inventory number
 (pl. 77, figs. 6–7)
- Chlorodiella mediterranea tetenyensis* MÜLLER,
 1984
 paratypes, M.86.333
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül84, p. 88
 Author's inventory number: MKK-7
- Chlorodiella mediterranea tetenyensis* MÜLLER,
 1984
 paratypes, M.86.352, M.86.353
 Badenian, Sámonháza Formation
 Fekete-hegy, Visegrád
 Mül84, p. 88
 Author's inventory number: MV-13

Crossotonotus diosdensis MÜLLER, 1984
holotype, M.89.60
Badenian, Rákócs Limestone Formation
sand pit, Diósdi
Mül84, p. 99, text-fig. 11, pl. 97, figs. 1–3
Author's inventory number: MDZ-15-1;
paratype under same inventory number
(pl. 97, figs. 4–5)

Diogenes matrensis MÜLLER, 1984
holotype, M.61.2633
Badenian, Sámonháza Formation
Szentkút, Mátraverebély
Mül84, p. 57, pl. 16, figs. 1–3

Dorippe ornatissima MÜLLER, 2006
holotype, 2005.95.1.
Badenian, Rákócs Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 41, pl. 1, fig. 1
Author's inventory number: MOLI 5.1

Dromia neogenica MÜLLER, 1978
holotype, M.86.419
Badenian, Rákócs Limestone Formation
Kerepesi út, Budapest
Mül78, p. 287 (278), pl. 8, fig. 1
Author's inventory number: MKC-4-1

Dromilites eotvoesi MÜLLER, 1975
holotype, 2007.97.1.
Badenian, Rákócs Limestone Formation
Tétényi-fennsík, Budapest
Mül75a, p. 512 (508), pl. 1, fig. 5, pl. 2, figs. 1, 4
Dromia eotvoesi, Mül84
Author's inventory number: MF-1-1 (MFA-1-1)

Dromilites eotvoesi MÜLLER, 1975
paratype, M.86.418
Badenian, Rákócs Limestone Formation
Kerepesi út, Budapest
Mül75a, p. 512 (508)
Dromia eotvoesi, Mül84
Author's inventory number: MKC-12

Dromilites eotvoesi MÜLLER, 1975
paratype, M.86.369
Badenian, Rákócs Limestone Formation
Tétényi-fennsík, Budapest
Mül75a, p. 512 (508)
Dromia eotvoesi, Mül84
Author's inventory number: MA-4

Ebalia globulosa MÜLLER, 1975
holotype, M.86.372
Badenian, Rákócs Limestone Formation
Tétényi-fennsík, Budapest
Mül75a, p. 512 (509), pl. 2, figs. 2, 3, 5
“*Palaeomyra*” *globulosa*, Mül84
Author's inventory number: MA-1-1

Ebalia hungarica MÜLLER, 1974
holotype, 2004.162.1.
Badenian, Rákócs Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 126 (121), pl. 1, figs. 6–7
Author's inventory number: MB-3-1

Ebalia hungarica MÜLLER, 1974
paratypes, M.86.511
Badenian, Rákócs Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 126 (121), pl. 1, fig. 5
Author's inventory number: MB-3

Ebalia oersi MÜLLER, 1978
holotype, M.86.78
Badenian, Rákócs Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 288 (279), pl. 13, figs. 1–3
Author's inventory number: MOE-4-1;
paratypes under same inventory number

Ethusa octospinosa MÜLLER, 2006
holotype, 2005.96.1.
Badenian, Rákócs Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 42, pl. 1, fig. 2
Author's inventory number: MOLI-4

Ethusa octospinosa MÜLLER, 2006
paratype, 2005.97.1.
Badenian, Rákócs Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 42, pl. 1, fig. 3
Author's inventory number: MOE-4-1

Haydnella steiningeri MÜLLER, 1984
paratype, M.86.111
Badenian, Sámonháza Formation
Törökmező, Nagymaros
Mül84, p. 90 (pl. 81, figs. 1–3)
Author's inventory number: MTZ-16

- Haydnella steiningeri* MÜLLER, 1984
paratypes, M.86.355
Badenian, Sámonháza Formation
Fekete-hegy, Visegrád
Mül84, p. 90
Author's inventory number: MV-15
- Haydnella steiningeri* MÜLLER, 1984
paratypes, M.86.449
Badenian, Leitha Limestone Formation
Grosshöflein (Nagyhöflány), Austria
Mül84, p. 90, pl. 80, figs. 3–5
Author's inventory number: MNH-9
- Haydnella steiningeri* MÜLLER, 1984
paratypes, M.86.450
Badenian, Leitha Limestone Formation
Grosshöflein (Nagyhöflány), Austria
Mül84, p. 90, pl. 81, fig. 4
Author's inventory number: MNH-9
- Kerepesia viai* MÜLLER, 1976
holotype, M.86.420
Badenian, Rákos Limestone Formation
Kerepesi út, Budapest
Mül76, p. 155 (150), pl. 3, figs. 1–3
Author's inventory number: MKC-3-1;
paratypes under same inventory number
(pl. 2, fig. 4)
- Metopograpsus badenis* MÜLLER, 2006
holotype, 2005.104.1.
Badenian, Rákos Limestone Formation
Rákos, Budapest
Mül06, p. 44, pl. 2, fig. 5
Author's inventory number: MRZ-21.1
- Metopograpsus badenis* MÜLLER, 2006
paratype, 2005.105.1.
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 44, pl. 2, fig. 6
Author's inventory number: MOX-19. Author
expressed doubt in identifying with holotype
- Micromithrax? grippi* MÜLLER, 1974
holotype, 2006.81.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 284 (279), pl. 2, figs. 1–2
Trachypirimela grippi, Mül84
Author's inventory number: MG-3-1
(MDG-15-1)
- Mioranina asymmetrica* MÜLLER, 1978
holotype, M.86.254
Badenian, Rákos Limestone Formation
Gyakorló út, Budapest
Mül78, p. 287 (278), pl. 9, fig. 2
Albunea asymmetrica, Mül84
Author's inventory number: MGY-2-1;
paratypes under same inventory number
(pl. 9, figs. 1, 3, pl. 10, figs. 1–3)
- Mioxaiava psammophila* MÜLLER, 1978
holotype, M.86.266
Badenian, Rákos Limestone Formation
Gyakorló út, Budapest
Mül78, p. 289 (281), pl. 16, figs. 1–4
Author's inventory number: MGY-5-1
- Pachygrapsus hungaricus* MÜLLER, 1974
holotype, 2004.163.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 126 (123), pl. 4, figs. 1–2
Author's inventory number: MB-5-1
- Pachygrapsus hungaricus* MÜLLER, 1974
paratypes, M.86.42
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74a, p. 126 (123), pl. 4, fig. 3, pl. 5, fig. 1
Author's inventory number: MB-5
- Paguristes cserhatensis* MÜLLER, 1984
holotype, M.61.3907
Badenian, Sámonháza Formation
Halastópuszta, Sámonháza
Mül84, p. 59, pl. 19, figs. 1–4
- Pagurus albus* MÜLLER, 1978
holotype, M.86.67
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 286 (277), pl. 4, figs. 4–5
Author's inventory number: MOE-2-1;
paratypes under same inventory number
- Pagurus concavus* MÜLLER, 1978
holotype, M.86.72
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 286 (277), pl. 7, figs. 1–2
Author's inventory number: MOE-1-1;
paratypes under same inventory number
(pl. 7, figs. 3–5)

Pagurus rakosensis MÜLLER, 1978
holotype, M.86.310
Badenian, Rákos Limestone Formation
Rákos, Budapest
Mül78, p. 286 (277), pl. 5, fig. 3, pl. 6, fig. 1
Author's inventory number: MR-1-1

Pagurus rakosensis MÜLLER, 1978
paratypes, M.86.68
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül78, p. 286 (277)
Author's inventory number: MOE-6

Pagurus rakosensis MÜLLER, 1978
paratypes, M.86.251
Badenian, Rákos Limestone Formation
Gyakorló út, Budapest
Mül78, p. 286 (277)
Author's inventory number: MGY-11

?*Pagurus turcus* MÜLLER, 1984
holotype, 2004.164.1.
Badenian, Sámonháza Formation
Törökmező, Nagymaros
Mül84, p. 56, pl. 13, figs. 1–3
Author's inventory number: MTZ-17-1;
paratypes under same inventory number
(pl. 12, fig. 6, pl. 13, fig. 4). Types missing

Palicus hungaricus MÜLLER, 2006
holotype, 2005.106.1.
Badenian, Rákos Limestone Formation
sand pit, Diósd
Mül06, p. 45, pl. 2, fig. 7
Author's inventory number: MDN-16.1

Palicus hungaricus MÜLLER, 2006
paratype, 2005.107.1.
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 45, pl. 2, fig. 8
Author's inventory number: MOE-32.2

Panopeus wronai MÜLLER, 1984
paratypes, M.86.117
Badenian, Sámonháza Formation
Törökmező, Nagymaros
Mül84, p. 91 (pl. 82, figs. 1–2)
Author's inventory number: MTZ-9

Panopeus wronai MÜLLER, 1984
paratypes, M.86.452
Badenian, Leitha Limestone Formation
Grosshöflein (Nagyhöflány), Austria
Mül84, p. 91, pl. 81, figs. 5–6, pl. 82, figs. 3–4,
pl. 83, figs. 1–4
Author's inventory number: M NH-10

Panopeus wronai MÜLLER, 1984
paratype, 2007.100.1.
Badenian, Rákos Limestone Formation
sand pit, Diósd
Mül84, p. 91
Author's inventory number: MDZ-2

Parthenope loczyi MÜLLER, 1974
holotype, 2006.84.1.
Badenian, Rákos Limestone Formation
Kerepesi út, Budapest
Mül74b, p. 283 (277), pl. 1, fig. 7
Mursia lienharti, Mül84
Author's inventory number: MKC-1-1

Parthenope szaboi MÜLLER, 1974
holotype, 2004.165.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 283 (277), pl. 1, fig. 8
Author's inventory number: MG-2-1
(MDG-9-1)

Parthenope szaboi MÜLLER, 1974
paratypes, M.86.169
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 283 (277), pl. 1, fig. 9
Author's inventory number: MDG-9

Parthenope tetenyensis MÜLLER, 1984
holotype, M.86.86
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül84, p. 74, pl. 55, figs. 1–2
Author's inventory number: MOE-16-1;
paratypes under same inventory number
(pl. 55, figs. 3–7, pl. 56, figs. 1, 3–7).
Paratype MOS-3-1 (pl. 56, fig. 2) not found
in inventory

Parthenope tetenyensis MÜLLER, 1984
 paratype, M.86.26
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül84, p. 74
 Author's inventory number: MBH-11

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.47
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül84, p. 74
 Author's inventory number: MUO-2

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.130
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül84, p. 74
 Author's inventory number: MJL-6

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.168
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül84, p. 74
 Author's inventory number: MDG-8

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.237
 Badenian, Rákos Limestone Formation
 Keresztúri út, Budapest
 Mül84, p. 74
 Author's inventory number: MEG-11

Parthenope tetenyensis MÜLLER, 1984
 paratype, M.86.261
 Badenian, Rákos Limestone Formation
 Gyakorló út, Budapest
 Mül84, p. 74
 Author's inventory number: MGY-18

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.283
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 74
 Author's inventory number: MRF-9

Parthenope tetenyensis MÜLLER, 1984
 paratypes, M.86.316
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 74
 Author's inventory number: MR-17

Parthenope tetenyensis MÜLLER, 1984
 paratype, M.86.405
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 74
 Author's inventory number: MR8-14

Petrolisthes haydni MÜLLER, 1984
 paratypes, M.86.106
 Badenian, Sámonháza Formation
 Törökmező, Nagymaros
 Mül84, p. 61 (pl. 26, fig. 1)
 Author's inventory number: MTZ-16

Petrolisthes haydni MÜLLER, 1984
 paratype, M.86.339
 Badenian, Rákos Limestone Formation
 borehole FRK-21, Fertőrákos
 Mül84, p. 61

Petrolisthes haydni MÜLLER, 1984
 paratypes, M.86.438
 Badenian, Leitha Limestone Formation
 Grosshöflein (Nagyhöflány), Austria
 Mül84, p. 61, pl. 26, figs. 2, 3, 6, 7, 8
 Author's inventory number: MNH-1

Petrolisthes haydni MÜLLER, 1984
 replica of paratype, 2007.102.1.
 Badenian, Leitha Limestone Formation
 Grosshöflein (Nagyhöflány), Austria
 Mül84, p. 61, pl. 26, figs. 4, 5
 Author's inventory number: MZZ-1

Petrolisthes magnus MÜLLER, 1984
 holotype, M.86.199
 Badenian, Rákos Limestone Formation
 Rákos, Budapest
 Mül84, p. 60, pl. 23, figs. 1-2
 Author's inventory number: MRZ-2-1;
 paratypes under same inventory number
 (pl. 23, figs. 3-4, pl. 24, figs. 1-4, pl. 25,
 figs. 4-5)

Petrolisthes magnus MÜLLER, 1984

paratypes, M.86.332

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül84, p. 60

Author's inventory number: MKK-2

Pilumnopaeus paratethiensis MÜLLER, 1984

holotype, M.86.203

Badenian, Rákos Limestone Formation

Rákos, Budapest

Mül84, p. 95, pl. 92, fig. 1

Author's inventory number: MRZ-10-1;

paratypes under same inventory number
(pl. 92, figs. 2-5)

"*Pilumnopaeus*" *tetenensis* MÜLLER, 1984

holotype, M.86.521

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül84, p. 95, pl. 91, figs. 4-5

Author's inventory number: MB-16-1;

paratype under same inventory number

"*Pilumnopaeus*" *tetenensis* MÜLLER, 1984

paratypes, M.86.301

Badenian, Rákos Limestone Formation

Rákos, Budapest

Mül84, p. 95

Author's inventory number: MRS-5

Pilumnus telegdi MÜLLER, 1974

holotype, 2004.166.1.

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül74b, p. 284 (280), pl. 2, figs. 7-8

Actumnus telegdi, Mül84

Author's inventory number: MG-4-1

(MDG-23-1)

Pilumnus telegdi MÜLLER, 1974

paratypes, M.86.175

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül74b, p. 284 (280)

Actumnus telegdi, Mül84

Author's inventory number: MG-4 (MDG-23)

Porcellana (Pisidia) kokayi MÜLLER, 1974

holotype, 2006.86.1.

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül74a, p. 126 (121), pl. 1, figs. 1-2

Pisidia kokayi, Mül84

Author's inventory number: MB-2-1

Porcellana (Pisidia) kokayi MÜLLER, 1974

paratype, M.86.34

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül74a, p. 126 (121)

Pisidia kokayi, Mül84

Author's inventory number: MKF-1

Porcellana (Pisidia) kokayi MÜLLER, 1974

paratypes, M.86.508

Badenian, Rákos Limestone Formation

Tétényi-fennsík, Budapest

Mül74a, p. 126 (121), pl. 1, figs. 3-4

Pisidia kokayi, Mül84

Author's inventory number: MB-2

Portunus (Monomia) miocaenicus MÜLLER, 1984

holotype, M.86.188

Badenian, Rákos Limestone Formation

Rákos, Budapest

Mül84, p. 80, pl. 62, fig. 5

Author's inventory number: MRO-10-1

Portunus neogenicus MÜLLER, 1978

holotype, M.86.262

Badenian, Rákos Limestone Formation

Gyakorló út, Budapest

Mül78, p. 288 (280), pl. 19

Author's inventory number: MGY-3-1

"*Pylopagurus*" *leganyii* MÜLLER, 1984

holotype, M.62.3253

Badenian, Sámonháza Formation

foot of Csúcs-hegy, Sámonháza

Mül84, p. 56, pl. 14, figs. 1-3

Rakosia carupoides MÜLLER, 1984

paratypes, M.86.202

Badenian, Rákos Limestone Formation

Rákos, Budapest

Mül84, p. 82, pl. 68, figs. 3-7 (pl. 68, figs. 1-2)

Author's inventory number: MRZ-14

Rakosia carupoides MÜLLER, 1984

paratype, M.86.444

Badenian, Leitha Limestone Formation

Grosshöflein (Nagyhöflány), Austria

Mül84, p. 82

Author's inventory number: MHN-6.

External mould of holotype carapace

(deposited in Naturhistorisches Museum, Vienna) under same inventory number

- Rakosia carupoides* MÜLLER, 1984
paratype, M.86.445
Badenian, Leitha Limestone Formation
Grosshöflein (Nagyhöflány), Austria
Mül84, p. 82
Author's inventory number: MNH-6
- Rakosia carupoides* MÜLLER, 1984
paratype, 2007.101.1.
Badenian, Rákos Limestone Formation
sand pit, Diósd
Mül84, p. 82
Author's inventory number: MDZ-7
- Schizophrys visegradensis* MÜLLER, 1984
holotype, M.86.223
Badenian, Sámonháza Formation
Törökmező, Nagymaros
Mül84, p. 72, pl. 49, figs. 4–5, pl. 50, fig. 1
Author's inventory number: MTZ-20-1;
paratypes under same inventory number
(pl. 50, figs. 2–3)
- Schizophrys visegradensis* MÜLLER, 1984
paratypes, M.86.348
Badenian, Sámonháza Formation
Fekete-hegy, Visegrád
Mül84, p. 72
Author's inventory number: MV-8
- Thalamita fragilis* MÜLLER, 1978
holotype, M.86.263
Badenian, Rákos Limestone Formation
Gyakorló út, Budapest
Mül78, p. 289 (281), pl. 17, fig. 1
Author's inventory number: MGY-4-1;
paratypes under same inventory number
(pl. 17, figs. 2–4)
- Thalamita fragilis* MÜLLER, 1978
paratypes, M.86.374
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül78, p. 289 (281)
Author's inventory number: MA-5
- Thalamita fragilis* MÜLLER, 1978
paratypes, M.86.430
Badenian, Rákos Limestone Formation
Kerepesi út, Budapest
Mül78, p. 289 (281)
Author's inventory number: MKC-15
- Thia szoereniae* MÜLLER, 1974
holotype, 2006.83.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 283 (278), pl. 2, figs. 3–4
"Lissocarcinus" szoeraeniae, Mül84
Author's inventory number: MD-1-1
(MDG-18-1)
- Trachypirimela radula* MÜLLER, 1974
holotype, 2006.82.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 284 (278), pl. 2, figs. 5–6
Trachypirimela grippi, Mül84
Author's inventory number: MA-1-1
(MDG-15-2)
- Trapezia glaessneri* MÜLLER, 1975
holotype, M.86.356
Badenian, Sámonháza Formation
Fekete-hegy, Visegrád
Mül75b, p. 521 (517), pl. 2, fig. 1
Author's inventory number: MV-1-1;
paratypes under same inventory number
(pl. 1, fig. 3, pl. 2, figs. 2–3)
- Tritodynamia miocaenica* MÜLLER, 2006
holotype, 2005.101.1.
Badenian, Rákos Limestone Formation
Rákos, Budapest
Mül06, p. 44, pl. 2, figs. 1, 2
Author's inventory number: MRW-1
- Tritodynamia miocaenica* MÜLLER, 2006
paratype, 2005.102.1.
Badenian, Rákos Limestone Formation
Örs vezér tere, Budapest
Mül06, p. 44, pl. 2, fig. 3
Author's inventory number: MOFH11
- Upogebia scabra* MÜLLER, 1974
holotype, 2004.167.1.
Badenian, Rákos Limestone Formation
Tétényi-fennsík, Budapest
Mül74b, p. 283 (276), pl. 1, fig. 2
Author's inventory number: MG-1-1
(MDG-3-1). Type redescribed as *Upogebia* sp.
in MÜLLER 1984

Upogebia scabra MÜLLER, 1974
 paratypes, M.86.157
 Badenian, Rákos Limestone Formation
 Tétényi-fennsík, Budapest
 Mül74b, p. 283 (276)
 Author's inventory number: MDG-3. Types
 redescribed as *Upogebia* sp. in MÜLLER 1984

"*Xaiva*" *bachmayeri* MÜLLER, 1984
 holotype, M.86.91
 Badenian, Rákos Limestone Formation
 Örs vezér tere, Budapest
 Mül84, p. 85, pl. 72, figs. 5-7
 Author's inventory number: MOE-26-1;
 paratype under same inventory number

8.9. Miocene Crinoidea

Actinometra mátraverebelyensis SZALAI, 1926
 holotype, M.62.3175
 Badenian, Sámonháza Formation
 Meszes-tető, Mátraverebély
 Szl26b, p. 173, pl. 1, figs. 6a-c

Antedon bölcseyensis SZALAI, 1926
 holotype
 Badenian, Sámonháza Formation
 Meszes-tető, Mátraverebély
 Szl26b, p. 171, pl. 1, figs. 1a-c
 Type probably destroyed in 1956

Antedon neograidiensis SZALAI, 1926
 holotype
 Badenian, Sámonháza Formation
 Meszes-tető, Mátraverebély
 Szl26b, p. 172, pl. 1, figs. 2a-c
 Type probably destroyed in 1956

Antedon quinquepetallus SZALAI, 1926
 holotype, M.62.3170
 Badenian, Sámonháza Formation
 Meszes-tető, Mátraverebély
 Szl26b, p. 172, pl. 1, figs. 4a-c
 Paratypes under same inventory number

Isocrinus stellata SZALAI, 1926
 holotype, M.62.3176
 Badenian, Sámonháza Formation
 Meszes-tető, Mátraverebély
 Szl26b, p. 173, pl. 1, figs. 5a-c

8.10. Miocene Echinoidea

Clypeaster epianthus MEZNERICS, 1941
 holotype, M.62.9614
 Badenian, Sámonháza Formation
 Mátraszólós
 Mez41, p. 87, pl. 2, fig. 1, pl. 3, fig. 2

Clypeaster kemencensis MEZNERICS, 1941
 holotype, M.62.9607
 Badenian, Sámonháza Formation
 Kemence
 Mez41, p. 88, pl. 2, fig. 3, pl. 3, fig. 6

Echinocardium intermedium LÓCZY, 1877
 syntype, M.60.9201
 Badenian, Dej Formation?
 Minișu de Sus (Felménés), Romania
 Lóc77, p. 64 (41), pl. 5, figs. 2a-h

Echinocardium intermedium LÓCZY, 1877
 syntypes, M.61.5750
 Badenian, Rákos Limestone Formation
 Bia, Biatorbágy
 Lóc77, p. 64 (41), pl. 5, figs. 1a-e

Echinolampas percassus MEZNERICS, 1941
 holotype, M.62.9612
 Badenian, Sámonháza Formation
 Kemence
 Mez41, p. 90, pl. 2, fig. 4, pl. 3, fig. 4

Prionechinus felmenensis LAMBERT & THIÉRY, 1911
 holotype, M.60.9212
 Badenian, Dej Formation?
 Minișu de Sus (Felménés), Romania
 Lam11, p. 230
 Species erected by indication with reference to
Echinus cf. *dux* LAUBE in LÓCZY 1877: p. 63 (40),
 pl. 5, figs. 3a-c. Species name emended to
felménensis by VADÁSZ (1914, 1915)

Prionechinus loczyi LAMBERT & THIÉRY, 1911
 holotype
 Badenian, Dej Formation?
 Minișu de Sus (Felménés), Romania
 Lam11, p. 230
 Species erected by indication with reference to
Psammechinus cf. *monilis* DESMAREST in LÓCZY
 1877: p. 63 (40), pl. 5, figs. 4a-d. Type lost
 according to VADÁSZ (1914, 1915)

9. Pliocene types

9.1. Pliocene Gastropoda

Bythinia (?) cholnokyi SCHLOSSER, 1906

syntypes

San-tao-kou Valley, China

ScM06, p. 391, pl. 10, figs. 32–39

Types probably lost

Melanopsis doboi rugosa SCHRÉTER, 1975

paratype, M.65.393

Early Pleistocene, Dunaalmás Travertine

Formation

Budakalász

ScZ75, p. 11 (pl. 3, figs. 15–18)

Melanopsis doboi rugosa SCHRÉTER, 1975

paratype, M.72.79

Early Pleistocene, Dunaalmás Travertine

Formation

railway cut by Zárkány bastion, Eger Castle,

Eger

ScZ75, p. 11

10. Pleistocene types

10.1. Pleistocene Gastropoda

Helicigona goemoeensis SOÓS, 1935

syntypes, M.57.1951

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög),

Slovakia

Tas35, p. 19

Melanopsis doboi SCHRÉTER, 1975

paratypes, M.65.392, M.72.81

Early Pleistocene, Dunaalmás Travertine

Formation

railway cut by Zárkány bastion, Eger Castle,
Eger

ScZ75, p. 7 (text-fig. 1, pl. 1, figs. 5–6)

Melanopsis doboi bicarinata SCHRÉTER, 1975

paratypes, M.65.394, M.72.78

Early Pleistocene, Dunaalmás Travertine

Formation

railway cut by Zárkány bastion, Eger Castle,
Eger

ScZ75, p. 8 (text-fig. 2, pl. 2, figs. 7–10)

Melanopsis doboi multifilosa SCHRÉTER, 1975

paratypes, M.72.77

Early Pleistocene, Dunaalmás Travertine

Formation

railway cut by Zárkány bastion, Eger Castle,
Eger

ScZ75, p. 10 (pl. 2, figs. 11–12, pl. 3, fig. 13)

Catalogue of vertebrate type specimens

11. Pisces

Belone harmati WEILER, 1933

holotype, V.61.229

skeleton

Kiscellian, Oligocene; Tard Clay Formation
Eged-hegy, Eger

Wei33, p. 18, text-fig. 7

Belone harmati WEILER, 1933

paratype, V.61.230

skeleton

Kiscellian, Oligocene; Tard Clay Formation
Eged-hegy, Eger

Wei33, p. 18

Belone harmati WEILER, 1933

paratype

skeleton

Kiscellian, Oligocene; Tard Clay Formation
Eged-hegy, Eger

Wei33, p. 18, text-fig. 8

Type missing, not found in inventory

Caranx bökhi GORJANOVIC-KRAMBERGER, 1902

holotype, V.69.244

skeleton

Badenian, Miocene; Leitha-Kalk Formation

Sankt Margarethen im Burgenland
(Szentmargitbánya), Austria

Gor02b (Gor02a), p. 9 (9), pl. 2, fig. 2

Carcharodon elegans NEUGEBOREN, 1850

syntypes, V.69.959

two teeth

Priabonian, Eocene; Valea Nișului Formation
Grohotiș, Turnu Roșu (Porcsesd), Romania
Neu50, p. 28 (pl. 1, figs. 28a-c)

Carcharodon humilis KOCH, 1904

syntype, V.69.895

one tooth

Eggenburgian, Miocene; Lipovany Sandstone
Formation

Pôtor (Nógrádszentpéter), Slovakia
Koc04a, p. 265 (194), pl. 1, figs. 4a-c

Carcharodon humilis KOCH, 1904

syntypes, V.69.1090

two teeth

Eggenburgian, Miocene; Lipovany Sandstone
Formation

Horné Strháre (Felsőesztergály), Slovakia
Koc04a, p. 265 (194), pl. 1, figs. 3a-c, 5a-c

Cybium lóczyi WEILER, 1938

syntype, V.61.669

skeletal fragments

Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest

Wei38, p. 22, pl. 5, fig. 3

Inventory number in original publication:
Nr. 369

Cybium lóczyi WEILER, 1938

syntype, V.61.716

skeletal fragments

Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest

Wei38, p. 22, pl. 4, fig. 1

Inventory number in original publication:
Nr. 370/b

Cybium lóczyi WEILER, 1938

syntype, V.61.728, V.61.748

skeletal fragments

Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest

Wei38, p. 22, pl. 4, fig. 2

Inventory number in original publication:
Nr. 370/a. Inventory items represent fragments
of same individual

Cybium lóczyi WEILER, 1938

syntype, V.61.771

skeletal fragments

Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest

Wei38, p. 22, pl. 1, fig. 3

Inventory number in original publication:
Nr. 486

Cybium lóczyi WEILER, 1938
 syntypes, V.61.792
 skeletal fragments
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
Wei38, p. 22, pl. 1, fig. 7
 Inventory number in original publication:
 Nr. 371

Gadus (Merlangus) pannonicus KOCH, 1904
 syntypes, V.69.235
 skeletal fragments
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4), pl. 1, fig. 1 (pl. 1, figs. 2–4,
 pl. 2, figs. 2–4, 6, 10–12, pl. 4, fig. 2, pl. 5, fig. 3)

Gadus (Merlangus) pannonicus KOCH, 1904
 syntypes, V.69.1123, V.69.1130, V.69.1141,
 V.69.1156, V.69.1214
 skeletal fragments
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4)

Gadus (Merlangus) pannonicus KOCH, 1904
 syntype, V.69.1153
 fragment of preoperculum
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4)

Gadus (Merlangus) pannonicus KOCH, 1904
 syntypes, V.69.1191
 premaxillae and mandible fragments
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4), pl. 1, figs. 7–8

Gadus (Merlangus) pannonicus KOCH, 1904
 syntype, V.69.1205
 frontale fragment
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4), pl. 2, fig. 1

Gadus (Merlangus) pannonicus KOCH, 1904
 syntypes, V.69.1206
 vertebrae and ribs
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
Koc04b, p. 37 (4)

Gymnosarda zsiivnyi WEILER, 1938
 syntype, M.62.2517
 skeleton
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
Wei38, p. 19, text-fig. 2, pl. 3, fig. 1
 Inventory number in original publication:
 Nr. 201

Gymnosarda zsiivnyi WEILER, 1938
 syntype, V.61.725, V.61.871
 skeletal fragments
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
Wei38, p. 19, pl. 1, fig. 9
 Inventory number in original publication:
 Nr. 350/b. Inventory items represent part and
 counterpart

Gymnosarda zsiivnyi WEILER, 1938
 syntype, V.61.731, V.61.864
 skeletal fragments
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
Wei38, p. 19, pl. 3, fig. 2
 Inventory number in original publication:
 Nr. 349. Inventory items represent fragments
 of same individual

Gymnosarda zsiivnyi WEILER, 1938
 syntype
 skeletal fragments
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
Wei38, p. 19, pl. 1, fig. 8
 Inventory number in original publication:
 Nr. 350/a. Type missing, not found in current
 inventory

Lamna ackneri NEUGEBOREN, 1851
 syntypes, V.69.1030
 eight teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotiș, Turnu Roșu (Porcsesd), Romania
Neu51, p. 200, pl. 5, figs. 63, 64a–b, 65a–b,
 66, 67

Lamna (Odontaspis) alveata NEUGEBOREN, 1851
 syntypes, V.69.1069
 two teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotiș, Turnu Roșu (Porcsesd), Romania
Neu51, p. 197, pl. 5, figs. 54a–c, 55a–c

Lamna cavidens NEUGEBOREN, 1851
 syntypes, V.69.1077
 eight teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 200, pl. 5, figs. 58a-c, 59a-b, 60-62

Lamna depressa NEUGEBOREN, 1851
 syntypes, V.69.998
 five teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 175, pl. 4, figs. 12a-c, 13a-c, 14-16

Lamna (Odontaspis) elongata NEUGEBOREN, 1851
 holotype, V.2008.35.1.
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 198, pl. 5, figs. 56a-c

Lamna (Odontaspis) ferox fossilis NEUGEBOREN,
 1851
 syntypes, V.69.1036
 three teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 202, pl. 5, figs. 72a-c, 73a-b, 74a-b

Lamna haueri NEUGEBOREN, 1851
 syntypes, V.69.902
 seven teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 201, pl. 5, figs. 68a-b, 69a-b, 70, 71a-c

Lamna minima NEUGEBOREN, 1851
 holotype, V.69.904
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 205, pl. 5, figs. 78a-b

Lamna minuta NEUGEBOREN, 1851
 holotype, V.69.1087
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 205, pl. 5, figs. 77a-b

Lamna (Odontaspis) serrata NEUGEBOREN, 1851
 syntypes, V.69.1094
 two teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 203, pl. 5, figs. 75a-c, 76

Lamna speciosa NEUGEBOREN, 1851
 syntypes, V.69.1032
 three teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 178, pl. 4, figs. 29a-b, 30a-c

Lamna xyphodon NEUGEBOREN, 1851
 holotype, V.69.1081
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 199, pl. 5, figs. 57a-d

Lates pliocaenus KOCH, 1904
 syntype, V.69.1155
 incomplete skeleton
 Pannonian, Miocene; "cement marl"
 cement quarry, Beočin (Beocsin), Serbia
 Koc04b, p. 52 (17), pl. 6, figs. 14a-b

Lepidopus hungaricus BÖHM, 1941
 syntypes, V.69.220
 tooth impressions
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 193 (pl. 2, fig. 8b, pl. 3, figs. 1a, 2,
 3a-b, pl. 4, fig. 1)

Lepidopus hungaricus BÖHM, 1941
 syntype, V.69.241
 impression of partial skeleton
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 193

Lepidopus hungaricus BÖHM, 1941
 syntypes, V.69.254
 four teeth
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 193, pl. 2, fig. 8a, pl. 3, fig. 1b

Lepidopus hungaricus BÖHM, 1941
 syntypes, V.69.1111
 one tooth, one vertebra, one fragmentary fin
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 193

Lepidopus hungaricus BÖHM, 1941
 syntypes, V.69.1162
 two partial skeletons and their impressions
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 193, pl. 2, figs. 6, 7

Leuciscus carpathicus BÖHM, 1941
 holotype, V.69.218
 skeleton without skull
 Kiscellian, Oligocene; Menilite Formation
 Covasna (Kovászna), Romania
 Böh41, p. 191, pl. 1, fig. 6

Nemopteryx kubacskei WEILER, 1935
 holotype, M.62.2518, V.61.898
 skeleton
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Wei35, p. 28, text-fig. 1
 Inventory items represent part and counterpart

Otodus ambiguus NEUGEBOREN, 1851
 holotype, V.69.892
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 158, pl. 3, figs. 9a-c

Otodus arcuato-decrescens NEUGEBOREN, 1851
 holotype, V.69.994
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 159, pl. 3, figs. 10a-b

Oxyrhina haueri NEUGEBOREN, 1851
 syntypes, V.69.995
 five teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 169, pl. 3, figs. 34a-c, 35, 36a-d,
 37a-b

Oxyrhina heckeliana NEUGEBOREN, 1851
 holotype, V.69.1049
 one tooth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 168, pl. 3, figs. 33a-c

Oxyrhina lata NEUGEBOREN, 1851
 syntypes, V.69.999
 two teeth
 Priabonian, Eocene; Valea Nișului Formation
 Grohotișu, Turnu Roșu (Porcsesd), Romania
 Neu51, p. 171, pl. 3, figs. 38, 39

Pycnodus scrobiculatus REUSS, 1844
 syntype, V.63.644
 one tooth
 Priabonian, Eocene
 Koštice (Kosstitz), Czech Republic
 Res44, p. 211
 Type redescribed and figured in REUSS 1846:
 p. 10, pl. 4, fig. 15

Serranus transsylvaniaicus BÖHM, 1942
 holotype, V.69.1178
 complete skeleton
 Kiscellian, Oligocene; Menilite Formation
 Malnaš (Málnás), Romania
 Böh42, p. 18, text-fig. 2, pl. 2, fig. 3

Smerdis hungaricus WEILER, 1938
 holotype, V.61.819
 skeleton
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Wei38, p. 18, pl. 1, figs. 4, 13, pl. 5, fig. 1
 Inventory number in original publication:
 Nr. 343

Smerdis hungaricus WEILER, 1938
 paratype, V.61.767
 skeleton
 Kiscellian, Oligocene; Kiscell Clay Formation
 Újlaki brickyard, Kiscell, Budapest
 Wei38, p. 18, pl. 1, fig. 24, pl. 5, fig. 2
 Inventory number in original publication:
 Nr. 330

Sphyraena pannonica WEILER, 1938
holotype, V.61.724, V.61.870
skeletal fragments
Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Wei38, p. 24, pl. 1, fig. 22, pl. 3, fig. 3, pl. 6, fig. 3
Inventory number in original publication:
Nr. 348. Inventory items represent part and
counterpart

Sphyraena pannonica WEILER, 1938
paratype, V.61.741, V.61.747
skeletal fragments
Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Wei38, p. 24, pl. 1, fig. 25
Inventory number in original publication:
Nr. 345. Inventory items represent part and
counterpart

Sphyraena pannonica WEILER, 1938
paratype
skeletal fragments
Kiscellian, Oligocene; Kiscell Clay Formation
Újlaki brickyard, Kiscell, Budapest
Wei38, p. 24, pl. 1, fig. 14
Inventory number in original publication:
Nr. 545/a. Type missing, not found in current
inventory

Clemmys hungarica SZALAI, 1934
holotype
left ilium
Pannonian, Miocene
cave deposit, Esterházy-barlang, Csákvár
Szl34, p. 106, pl. 4, fig. 1
Type lost

Emys strandi SZALAI, 1934
paralectotype, V.61.1155
plastron fragment
Kiscellian, Oligocene; Gruia Sandstone
Formation
Cluj-Napoca (Kolozsvár), Romania
Szl34, p. 108, pl. 1, figs. 3–4 (pl. 1, figs. 1–2, 5–6)
Chinemys strandi, Mly66
Type missing

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.1.
left premaxilla
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 2a–d
Inventory number of holotype skeleton in
original publication: Gyn/404.

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.2.
right premaxilla with one tooth
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 2a–d

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.3.
vomer
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 4b

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.4.
right postorbital, jugal and quadratojugal
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 3a–b

12. Reptilia

Bakonydraco galaczi ŐSI, WEISHAMPEL & JIANU, 2005
holotype, 2007.110.1.
nearly complete mandible
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05b, p. 778, text-figs. 2a–f
Inventory number in original publication:
Gyn/3.

Bakonydraco galaczi ŐSI, WEISHAMPEL & JIANU, 2005
paratypes, 2007.111.1.
21 symphyseal fragments of dentary
Santonian, Late Cretaceous; Csehbánya
Formation
Iharkút, Bakonyjákó
Ősi05b, p. 778, text-figs. 3a–b
Inventory number in original publication:
Gyn/4.

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.5.
left prefrontal
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 2g-h

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.6.
left lacrimal
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.8.
left frontal
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 2e-f

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.9.
pterygoid
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 4a

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.10.
right quadrate
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 5b, 5d

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.11.
part of left quadrate
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 5a

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.12.
occipital condyle
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 4c

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.13.
21 isolated teeth
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 4e-g

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.14.
hyoid bone
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 4d

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.15.
anterior part of dentary, surangular, angular,
small posterior part of dentary
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 6a-e

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.16.
three cervical vertebrae
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 7a-i

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.17.
six dorsal vertebrae
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 8a-q

- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.18.
ten caudal vertebrae
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 10a–z, 10a'–d'
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.19.
three cervical ribs
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.20.
13 dorsal ribs
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.21.
five chevrons
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.22.
some tendon fragments
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.23.
complete left scapulocoracoid
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 11b
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.24.
right scapula and anterior part of right coracoid
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 11a
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.25.
fused sacral vertebrae (described originally as right radius)
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 11c–d
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.26.
four fragmentary sacral ribs (described originally as metacarpals)
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 11f–i
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.27.
preacetabular part of left ilium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.28.
preacetabular part of right ilium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
- Hungarosaurus tormai* ŐSI, 2005
holotype, 2007.26.29.
left ischium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.30.
right femur
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 12e-f

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.31.
right fibula
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 12g-i

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.32.
dorsal osteoderms
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 13b-g

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.26.33.
cervical osteoderms
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 13a

Hungarosaurus tormai ŐSI, 2005
holotype, 2007.89.1.
ungual phalanx and middle phalanx
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 11j-m

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.22.1.
one anterior dorsal vertebra, cervical and dorsal ribs, one half-ring osteoderm, some small circular osteoderms
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
Inventory number in original publication: Gyn/405.

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.23.1.
sacral, fused osteoderm
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-fig. 13h
Inventory number in original publication: Gyn/407.

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.23.2.
left ischium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 12c-d
Inventory number in original publication: Gyn/407.

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.23.3.
right ischium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
Inventory number in original publication: Gyn/407.

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.23.4.
left ilium
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371
Inventory number in original publication: Gyn/407.

Hungarosaurus tormai ŐSI, 2005
paratype, 2007.90.1.
sacrum with sacral rod
Santonian, Late Cretaceous; Csehbánya Formation
Iharkút, Bakonyjákó
Ősi05a, p. 371, text-figs. 9a-b
Inventory number in original publication: Gyn/407.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.90.2.
 left fragmentary ilium
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371, text-figs. 12a–b
 Inventory number in original publication:
 Gyn/407.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.24.1.
 distal end of femur?
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371
 Inventory number in original publication:
 Gyn/406.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.24.2.
 fragmentary ulna
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371, text-fig. 11e
 Inventory number in original publication:
 Gyn/406.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.24.5.
 osteoderms
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371
 Inventory number in original publication:
 Gyn/406.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.24.6.
 fragmentary ribs
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371
 Inventory number in original publication:
 Gyn/406.

Hungarosaurus tormai ŐSI, 2005
 paratype, 2007.24.10.
 metapodium
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi05a, p. 371
 Inventory number in original publication:
 Gyn/406.

Iharkutosuchus makadii ŐSI, CLARK &
 WEISHAMPEL, 2007
 holotype, 2006.52.1.
 nearly complete skull
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170, text-figs. 1a–f, 2i–j

Iharkutosuchus makadii ŐSI, CLARK &
 WEISHAMPEL, 2007
 paratype, 2006.53.1.
 nearly complete skull
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170, text-figs. 2a–b, 3a

Iharkutosuchus makadii ŐSI, CLARK &
 WEISHAMPEL, 2007
 paratypes, 2006.54.1., 2006.55.1., 2006.56.1.
 fragmentary skulls
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK &
 WEISHAMPEL, 2007
 paratype, 2006.57.1.
 skull fragment
 Santonian, Late Cretaceous; Csehbánya
 Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170, text-figs. 3b–d

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.58.1.
 fragmentary mandible
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170, text-figs. 2c-h

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratypes, 2006.59.1., 2006.60.1., 2006.61.1., 2006.62.1., 2006.63.1., 2006.64.1., 2006.71.1., 2006.72.1., 2006.74.1., 2006.75.1., 2006.76.1.
 fragmentary mandibles
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratypes, 2006.65.1., 2006.66.1.
 skull fragments
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.67.1.
 parietale
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.68.1.
 frontale
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.69.1.
 supraoccipitale
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.70.1.
 right angulare
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.73.1.
 left premaxilla
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.77.1.
 left squamosum
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.78.1.
 right surangulare
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratype, 2006.79.1.
 frontale
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

Iharkutosuchus makadii ŐSI, CLARK & WEISHAMPEL, 2007
 paratypes, 2006.80.1.
 148 isolated teeth
 Santonian, Late Cretaceous; Csehbánya Formation
 Iharkút, Bakonyjákó
 Ősi07, p. 170

- Magyarosaurus transylvanicus* HUENE, 1932
syntype, V.60.1709
left fibula
Maastrichtian, Late Cretaceous; Sânpetru Formation
Vălioara (Valiora), Romania
Hue32, p. 266, pl. 46, fig. 11 (pl. 46, figs. 1–10, 12)
Magyarosaurus dacus, LeL93
This specimen also serves as syntype of *Titanosaurus dacus* NOPCSA, 1915
- Mosasaurus carentonensis* COQUAND, 1860
holotype, V.69.1167
tooth
Coniacian, Late Cretaceous
Faubourg Saint-Jacques, Cognac, France
Coq60, p. 70
- Testudo fejérváryi* SZALAI, 1930
lectotype, V.61.1142
plastron fragment
Ottnangian, Miocene; Salgótarján Formation
Kotyháza, Salgótarján
Szl30, p. 359, pl. 24, figs. 1–3 (pls. 21–22, pl. 23, fig. 1)
Ptychogaster fejervaryi, Mly66
Lectotype designated in MŁYNARSKI 1966, erroneously quoting inventory number V.61.1139/1
- Testudo fejérváryi* SZALAI, 1930
paralectotypes, V.61.1139
plastron fragments
Ottnangian, Miocene; Salgótarján Formation
Kotyháza, Salgótarján
Szl30, p. 359, pl. 23, figs. 2–4, pl. 24, figs. 4–6
Ptychogaster fejervaryi, Mly66
- Testudo hungarica* SZALAI, 1934
syntype, V.61.1144
left ilium
Pannonian, Miocene
cave deposit, Polgárdi
Szl34, p. 123, pl. 4, figs. 9a–b
Type redescribed as Emydinae gen. et sp. indet. in MŁYNARSKI 1966
- Testudo hungarica* SZALAI, 1934
syntype, V.61.1145
right tibia
Pannonian, Miocene
cave deposit, Polgárdi
Szl34, p. 123, pl. 4, figs. 15a–b
Type redescribed as Emydinae gen. et sp. indet. in MŁYNARSKI 1966
- Testudo lambrechti* SZALAI, 1934
syntypes
plastron fragments and right humerus
Middle Pliocene
quarry, Beremend
Szl34, p. 128, pl. 4, fig. 19a
Types lost
- Testudo lambrechti* SZALAI, 1934
syntype
right femur
Middle Pleistocene
Nagyharsány-hegy, Villány
Szl34, p. 128, pl. 4, fig. 19b
Type lost
- Testudo lambrechti* SZALAI, 1934
syntypes
left and right humerus, left femur, bone and plastron fragments
Middle Pliocene
Csarnóta
Szl34, p. 128, pl. 4, fig. 19c
Types lost
- Testudo rácmeckeensis* SZALAI, 1934
lectotype, V.61.1160
epiplastron and xiphiplastron fragments
Karpatian, Miocene; Budafa Sandstone Formation
Rácmecke, Erdősmecske
Szl34, p. 118
Lectotype fixed by inference of holotype in MŁYNARSKI 1966; paralectotypes (carapace and plastron fragments) under same inventory number

Testudo süttöensis SZALAI, 1934

lectotype, V.61.1224

right coracoid

Late Pleistocene; fissure fill in Dunaalmás

Travertine Formation

Süttő

Szl34, p. 131, pl. 4, fig. 20

Lectotype fixed by inference of holotype in MŁYNARSKI 1966

Testudo süttöensis SZALAI, 1934

paralectotype, V.61.1223

femur

Late Pleistocene; fissure fill in Dunaalmás

Travertine Formation

Süttő

Szl34, p. 131

Testudo süttöensis SZALAI, 1934

paralectotype, V.61.1225

humerus

Late Pleistocene; fissure fill in Dunaalmás

Travertine Formation

Süttő

Szl34, p. 131

Testudo süttöensis SZALAI, 1934

paralectotype, V.61.1227

plastron fragment

Late Pleistocene; fissure fill in Dunaalmás

Travertine Formation

Süttő

Szl34, p. 131

Titanosaurus dacus NOPCSA, 1915

syntype, V.60.1709

left fibula

Maastrichtian, Late Cretaceous; Sânpetru Formation

Vălioara (Valiora), Romania

Nop15b (Nop15a), p. 14 (14) (pl. 3, figs. 4, 5)

Magyarosaurus dacus, LeL93

This specimen also serves as syntype of

Magyarosaurus transylvanicus HUENE, 1932

Trionyx nopscai SZALAI, 1934

lectotype

mandible

Pannonian, Miocene

Brusturi (Tataros), Romania

Szl34, p. 134, pl. 4, fig. 22

Lectotype fixed by inference of holotype in

MŁYNARSKI 1966. Type in GIH collection as Ob.3980 but quoted in original publication as in HNHM

Trionyx nopscai SZALAI, 1934

paralectotype

carapace fragment

Pannonian, Miocene

Brusturi (Tataros), Romania

Szl34, p. 134

Type in GIH collection as Ob.3146 but quoted in original publication as in HNHM

13. Aves

Anas crecca percrecca JÁNOSSY, 1992

holotype, V.91.150

ulna

Middle Pliocene

quarry, locality 16, Beremend

Ján92, p. 17, text-fig. 4.8.

Apus submelba JÁNOSSY, 1972

holotype, V.64.435

ulna

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján72c, p. 56 (pl. 2, figs. 12–17)

Ciconia stehlini JÁNOSSY, 1992

holotype, V.91.151

proximal and distal fragments of tarsometatarsus

Middle Pliocene

quarry, locality 15, Beremend

Ján92, p. 16, text-figs. 4.1–3.

Ciconia stehlini JÁNOSSY, 1992

paratype, V.94.22

distal fragment of tibiotarsus

Middle Pliocene

quarry, locality 15, Beremend

Ján92, p. 16, text-fig. 4.4.

Otis khosatzkii beremendensis JÁNOSSY, 1991

holotype, V.90.11

tarsometatarsus

Middle Pliocene

quarry, locality 15, Beremend

Ján91, p. 24

Otis khosatzkii beremendensis JÁNOSSY, 1991

paratype, V.90.11

phalanx

Middle Pliocene

quarry, locality 15, Beremend

Ján91, p. 24

- Porzana estramosi* JÁNOSSY, 1979
holotype, V.78.120
broken distal fragment of left tarsometatarsus
Middle Pliocene
Osztramos, locality 9, Tornaszentandrás
Ján79, p. 20, text-fig. 4.8.
- Strix intermedia* JÁNOSSY, 1972
holotype, V.64.746
coracoid
Middle Pleistocene
Tar-kő, rock-shelter, Felsőtárkány
Ján72c, p. 53 (pl. 2, fig. 6)
- Tetrastes praebonasia* JÁNOSSY, 1974
holotype, V.64.789
right tarsometatarsus
Middle Pleistocene
Tar-kő, rock-shelter, Felsőtárkány
Ján74b, p. 218
- Tetrastes praebonasia* JÁNOSSY, 1974
paratype, V.64.617
dorsal end of right coracoid
Middle Pleistocene
Tar-kő, rock-shelter, Felsőtárkány
Ján74b, p. 218
- Turdoidea borealis* JÁNOSSY, 1979
holotype, V.78.119
proximal fragment of left humerus
Middle Pliocene
Osztramos, locality 1, Tornaszentandrás
Ján79, p. 27, text-fig. 4.7.
- 14. Mammalia**
- Agriarctos gaáli* KRETZOI, 1942
holotype, V.60.1751
P₃-M₂ and tip of canine of same individual
Pannonian, Miocene
brickyard, Strázsa-hegy, Hatvan
Kre42d, p. 351 (258), text-figs. 1.3-4.
- Agriarctos gaáli* KRETZOI, 1942
paratypes, V.60.1752
distal end of humerus and proximal part of
ulna
Pannonian, Miocene
brickyard, Strázsa-hegy, Hatvan
Kre42d, p. 351 (258)
- Alces brevirostris* KRETZOI in JÁNOSSY, 1969
holotype, V.82.14
incomplete skeleton
Middle Pleistocene
Ördöglyuk-barlang, Solymár
Ján69b, p. 620, pl. 2, figs. 1-3
Type redescribed and figured in VÖRÖS 1985:
p. 59, text-fig. 1, pl. 1, figs. 1-10, pl. 2, figs. 1-11.
Left side of antler (figured in JÁNOSSY 1969b:
pl. 2, fig. 3) destroyed in 1956
- Allocricetus bursae* SCHAUB, 1930
syntypes, V.61.2159
mandibles
Middle Pleistocene
Templom-hegy, Villány
ScS30, p. 33 (text-figs. 11, 16, 17, pl. 1, figs. 7, 9)
- Allocricetus bursae* SCHAUB, 1930
syntypes, V.61.2180
mandibles
Middle Pliocene
quarry, Beremend
ScS30, p. 33
- Allocricetus bursae* SCHAUB, 1930
syntypes, V.61.2204
mandibles
Pleistocene
Magyarkő, Brașov (Brassó), Romania
ScS30, p. 33
- Allocricetus bursae* SCHAUB, 1930
syntype, V.61.2205
left mandible
Pleistocene
Magyarkő, Brașov (Brassó), Romania
ScS30, p. 33
- Allocricetus bursae* SCHAUB, 1930
syntypes, V 2008.1.1.
two left and two right mandible fragments
Middle Pliocene
quarry, Beremend
ScS30, p. 33
- Allocricetus bursae* SCHAUB, 1930
syntype, V 2008.1.2.
right maxilla fragment
Middle Pliocene
quarry, Beremend
ScS30, p. 33

Allocricetus éhiki SCHAUB, 1930
 syntypes, V.61.2167
 mandibles
 Middle Pleistocene
 Templom-hegy, Villány
 ScS30, p. 34 (text-fig. 18, pl. 1, fig. 10)

Allocricetus éhiki SCHAUB, 1930
 syntype, V 2008.2.1.
 left mandible fragment
 Middle Pliocene
 quarry, Beremend
 ScS30, p. 34

Allocricetus éhiki SCHAUB, 1930
 syntypes, V 2008.3.1.
 five right mandible fragments
 Middle Pliocene
 quarry, Beremend
 ScS30, p. 34

Allophaiomys laguroides KORMOS, 1932
 holotype, V.61.1494
 skull fragment
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 331, text-fig. 4
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys laguroides KORMOS, 1932
 holotype, V.61.1494
 left mandible
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 331, text-fig. 5
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys laguroides KORMOS, 1932
 paratypes, V.61.1490
 left and right mandibles
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 331
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys pliocaenicus KORMOS, 1932
 holotype, V.61.1491
 skull fragment
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 326, text-fig. 1
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys pliocaenicus KORMOS, 1932
 holotype, V.61.1491
 left mandible
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 326, text-fig. 2
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys pliocaenicus KORMOS, 1932
 paratype, V.61.1491
 right mandible fragment
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 326, text-fig. 3
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys pliocaenicus KORMOS, 1932
 paratypes, V.61.1492
 left and right mandibles
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 326
Microtus (Allophaiomys) pliocaenicus, Rab81

Allophaiomys pliocaenicus KORMOS, 1932
 paratypes, V.61.1493
 left and right M³
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 326
Microtus (Allophaiomys) pliocaenicus, Rab81

Alopex praeglacialis KORMOS, 1932
 paratype, V.61.2164
 right P⁴
 Middle Pleistocene
 Templom-hegy, Villány
 Krm32b, p. 168, pl. 5, fig. 5 (pl. 5, figs. 1–4)
Vulpes praeglacialis, Rab76

- Alopex praeglacialis* KORMOS, 1932
 paratypes, V.61.2164
 mandible fragment and isolated canine
 Middle Pleistocene
 Templom-hegy, Villány
 Krm32b, p. 168
Vulpes praeglacialis, Rab76
- Amblycoptus oligodon* KORMOS, 1926
 syntype, V.58.1759
 left mandible
 Pannonian, Miocene
 cave deposit, Somlyó-hegy, Polgárdi
 Krm26, p. 372 (354), pl. 3, figs. 3a-d
 (pl. 3, figs. 1a-b, 2, 5a-b)
- Amblycoptus oligodon* KORMOS, 1926
 syntype, V.58.1759
 left upper incisor
 Pannonian, Miocene
 cave deposit, Somlyó-hegy, Polgárdi
 Krm26, p. 372 (354), pl. 3, figs. 4a-b
- Amblycoptus topali* JÁNOSSY, 1972
 holotype, V.71.149
 right mandible
 Middle Pliocene
 Osztramos, quarry No. 11, locality 1,
 Tornaszentandrás
 Ján72a, p. 38, pl. 3, figs. 8-10
Kordosia topali, Més97
 Inventory number quoted erroneously
 in original publication: V.71.49
- Amblycoptus topali* JÁNOSSY, 1972
 paratype, V.73.18
 skull fragment
 Middle Pliocene
 Osztramos, quarry No. 11, locality 1,
 Tornaszentandrás
 Ján72a, p. 38, pl. 3, fig. 7
Kordosia topali, Més97
- Amphicynops platyodon* KRETZOI, 1941
 holotype, V.60.1723
 left mandible
 Oligocene
 Badlands, Dakota, USA
 Kre41c, p. 174, text-fig. 2
- Amynodon hungaricus* KRETZOI, 1940
 holotype, V.60.149
 right mandible fragment with M_2-M_3
 Eocene?
 Tápiószele
 Kre40, p. 91, pl. 3, figs. 10, 11
- Anomalospalax tardosi* KORDOS, 1985
 holotype, V.84.135
 adult right mandible fragment with M_1-M_3
 Turolian, Miocene
 fissure fill in limestone quarry, Tardosbánya,
 Tardos
 Krd85, p. 27, pl. 1, fig. 1 (pl. 1, fig. 3, pl. 2,
 figs. 1-3, pl. 3, figs. 2-4, pl. 4, figs. 1-2, pl. 5,
 figs. 1-9, pl. 6, figs. 1-9)
- Anomalospalax tardosi* KORDOS, 1985
 paratype, V.84.136
 bilateral maxilla fragment with M^1-M^3 and
 M^2-M^3
 Turolian, Miocene
 fissure fill in limestone quarry, Tardosbánya,
 Tardos
 Krd85, p. 27, pl. 3, fig. 1
- Anomalospalax tardosi* KORDOS, 1985
 paratype, V.84.137
 bilateral maxilla fragment with M^1-M^3 and
 M^1-M^2
 Turolian, Miocene
 fissure fill in limestone quarry, Tardosbánya,
 Tardos
 Krd85, p. 27, pl. 4, figs. 3-4
- Anomalospalax tardosi* KORDOS, 1985
 paratype, V.84.138
 left mandible fragment with M_1-M_3
 Turolian, Miocene
 fissure fill in limestone quarry, Tardosbánya,
 Tardos
 Krd85, p. 27, pl. 1, fig. 2

Anomalopalax tardosi KORDOS, 1985
 paratype, V.84.139
 right mandible fragment
 Turolian, Miocene
 fissure fill in limestone quarry, Tardosbánya,
 Tardos
 Krd85, p. 27, pl. 1, fig. 4, pl. 2, fig. 4

Archidiskodon meridionalis ürömensis VÖRÖS,
 1979
 holotype, V.72.116
 left M^3
 Middle Pleistocene
 Üröm-hegy, Budapest
 Völ79, p. 5, pl. 1

Archidiskodon meridionalis ürömensis VÖRÖS,
 1979
 paratype, V.59.913
 aboral fragment of left M^2
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Völ79, p. 5

Baranogale helbingi KORMOS, 1934
 holotype, V.61.1380
 left mandible
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34b, p. 145, pl. 2, figs. 6a-b (pl. 2, figs.
 7a-b)
 Inventory number in original publication:
 Nr. 3914

Baranomys (Warthamys) kowalskii progressus
 JÁNOSSY, 1972
 holotype, V.71.151
 left M_3
 Middle Pliocene
 Osztramos, quarry No. 11, locality 1,
 Tornaszentandrás
 Ján72a, p. 44, pl. 1, fig. 11 (pl. 1, figs. 4-10)
 Inventory number quoted erroneously
 in original publication: V.71.51

Baranomys lóczyi KORMOS, 1933
 holotype, V.61.1527
 left mandible
 Middle Pliocene
 upper quarry, Csarnóta
 Krm33b, p. 48 (45), text-figs. 1-3
 Inventory number in original publication:
 Nr. 3808

Barbastella rostrata TOPÁL, 1970
 holotype, V.70.94
 left mandible with C- M_3 and alveoli of incisors
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Top70, p. 17 (6)

Barbastella rostrata TOPÁL, 1970
 paratype, V.70.95
 rostrum, right P^4-M^3 , left M^2-M^3
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Top70, p. 17 (6)

Bison bonasus hungarorum KRETZOI, 1946
 holotype
 skull fragment with one horn core
 Holocene
 locality unknown
 Kre46, p. 105, text-figs. 1-2
 Type missing, not found in inventory

Blarinella europaea REUMER, 1985
 paratypes, V.73.16
 left maxilla fragment with M^1-M^3 , two upper
 incisors, one mandible fragment
 Middle Pliocene
 Osztramos, locality 1, Tornaszentandrás
 Rem85, p. 69, pl. 21, fig. 5 (pl. 21, figs. 1-2, 4,
 6-7, pl. 22, figs. 1, 3)

Blarinella europaea REUMER, 1985
 paratypes, V.75.11
 right ascending ramus and isolated M^1
 Middle Pliocene
 Osztramos, locality 1, Tornaszentandrás
 Rem85, p. 69, pl. 22, figs. 2a-c

Blarinella europaea REUMER, 1985
 paratype, V 2008.11.1.
 maxilla fragment
 Late Pliocene
 Osztramos, locality 7, Tornaszentandrás
 Rem85, p. 69, pl. 21, figs. 3a-b

Blarinella europaea REUMER, 1985
 paratypes, V 2008.11.2.
 three mandible fragments
 Late Pliocene
 Osztramos, locality 7, Tornaszentandrás
 Rem85, p. 69, pl. 21, figs. 9a-b

- Blarinella europaea* REUMER, 1985
paratypes, V 2008.11.3.
one upper and one lower incisor
Late Pliocene
Osztramos, locality 7, Tornaszentandrás
Rem85, p. 69, pl. 21, figs. 8a–b
- Cadurcopsis dakotana* KRETZOI, 1942
holotype
skull
Oligocene
Badlands, Dakota, USA
Kre42b, p. 140, text-fig. 2
Type probably destroyed in 1956
- Canis spelaeoides* KRETZOI, 1942
holotype, 2007.95.1.
left mandible fragment with P_4 and trigonid
of M_1
Early Pleistocene
fissure fill in quarry, Gombasek (Gombaszög),
Slovakia
Kre42f, p. 349
Replacement name for "*Canis*" *gigas* KRETZOI,
1938. Description in KRETZOI 1938: p. 128,
pl. 2, fig. 10. Referred to genus *Xenocyon* in
KRETZOI 1941a. Inventory number in original
publication: Fa. 19
- Capra (Capra) zimmermanni* KRETZOI, 1942
holotype
skull fragment with horn cores
Late Pleistocene
from riverbed of Tisza, exact locality unknown
Kre42e, p. 356 (262), text-fig. 1.1., pl. 24,
figs. 1–3
Type missing, not found in inventory
- Castor eberezkyi* KRENNER, 1867
syntype, 2007.304.1.
right M^3
Middle Pliocene
Hajnáčka (Ajnácskő), Slovakia
Krn67, p. 129, pl. 2, figs. 17–18
- Castor eberezkyi* KRENNER, 1867
syntype, 2007.304.2.
right M^2
Middle Pliocene
Hajnáčka (Ajnácskő), Slovakia
Krn67, p. 129, pl. 2, figs. 12–14
- Castor eberezkyi* KRENNER, 1867
syntype, 2007.304.3.
upper molar
Middle Pliocene
Hajnáčka (Ajnácskő), Slovakia
Krn67, p. 129, pl. 2, figs. 15–16
- Citellus primigenius* KORMOS, 1934
paratypes, V.67.249
skull and left mandible
Middle Pleistocene
Templom-hegy, Villány
Krm34a, p. 314 (text-fig. 45)
- Clethrionomys acrorhiza* KORMOS, 1933
holotype, V.61.1483
left mandible fragment with M_1 – M_2
Early Pleistocene
Dealul Sprenghi (Fortyogó-hegy,
Gesprengberg), Brașov (Brassó), Romania
Krm33a, p. 5, text-fig. 1
- Clethrionomys hintonianus* KRETZOI, 1958
paratype, V.61.1486
left mandible
Middle Pleistocene
Nagyharsány-hegy, Villány
Kre58, p. 55
Replacement name for *Clethrionomys hintoni*
KORMOS, 1934. Description in KORMOS 1934a:
p. 318, text-fig. 48
- Cricetinus beremendensis* HÍR, 1994
holotype, V.93.3
right mandible fragment with M_1 – M_3
Middle Pliocene
quarry, locality 15, Beremend
Hír94, p. 72, text-figs. 1, 3, 4c (text-figs. 4d,
5, 8, 10a–c, 13, 15–17)
- Cricetinus beremendensis* HÍR, 1994
paratype, V.93.4
left maxilla fragment with M^1 – M^3
Middle Pliocene
quarry, locality 15, Beremend
Hír94, p. 72, text-fig. 2
- Cricetinus janossyi* HÍR, 1996
holotype, 2007.91.1.
mandible fragment with M_1 – M_3
Late Pliocene
Osztramos, locality 7, Tornaszentandrás
Hír96, p. 80, text-figs. 4, 12 (text-figs. 5–11,
13–16)

Cricetusin janossyi HÍR, 1996

paratype, 2007.91.2.

maxilla fragment with M²-M³

Late Pliocene

Osztramos, locality 7, Tornaszentandrás
Hír96, p. 80, text-figs. 2, 3

Cricetus cricetus nanus SCHAUB, 1930

syntypes, V 2008.4.1.

three left and two right mandible fragments

Early Pleistocene

Somlyó-hegy, Püspökkürdő, Oradea
(Nagyvárad), Romania

ScS30, p. 30 (text-figs. 9, 10, 22, pl. 2, figs. 1, 3)

Cricetus cricetus nanus SCHAUB, 1930

syntypes, V 2008.4.2.

two skull fragments

Early Pleistocene

Somlyó-hegy, Püspökkürdő, Oradea
(Nagyvárad), Romania

ScS30, p. 30

Cricetus cricetus praeglacialis SCHAUB, 1930

syntypes, V.61.2142

maxillae and mandibles

Middle Pleistocene

Nagyharsány-hegy, Villány

ScS30, p. 28 (text-figs. 14, 15, 21, pl. 2, figs. 2, 4)

Cricetus cricetus praeglacialis SCHAUB, 1930

syntype, V.61.2202

left maxilla fragment

Pleistocene

Magyarkő, Brașov (Brassó), Romania

ScS30, p. 28

Cricetus runtonensis solymarensis HÍR, 1997

holotype, 2007.92.1.

viscerocranium

Middle Pleistocene

Ördöglyuk-barlang, Solymár

Hír97, p. 25 (text-figs. 16-17)

Cricetus runtonensis solymarensis HÍR, 1997

paratype, 2007.92.2.

right mandible

Middle Pleistocene

Ördöglyuk-barlang, Solymár

Hír97, p. 25

Crocidura obtusa KRETZOI, 1938

holotype

mandible

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög),
Slovakia

Kre38, p. 92, text-fig. 1a

Inventory number in original publication:
Fa. 16. Type missing, not found in current
inventory

Crossopus fissidens PETÉNYI in KUBINYI, 1856

syntypes, V.61.1585

six maxilla and 37 mandible fragments

Middle Pliocene

quarry, Beremend

Kub56, p. 75

Beremendia fissidens, Krm34a

Types figured and described in detail in
PETÉNYI 1864: p. 60, pl. 1, figs. 5a-o

Deinsdorffia janossyi REUMER, 1985

holotype, V.83.11

right horizontal ramus with complete
dentition

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Rem85, p. 45, pl. 12, figs. 5a-b (pl. 12, fig. 1)

Deinsdorffia janossyi REUMER, 1985

paratypes, V 2008.7.1.

seven mandible fragments

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Rem85, p. 45, pl. 12, figs. 4a-b, 6a-c

Deinsdorffia janossyi REUMER, 1985

paratypes, V 2008.7.2.

two lower incisors and three lower molars

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Rem85, p. 45

Deinsdorffia janossyi REUMER, 1985

paratypes, V 2008.7.3.

11 maxilla fragments

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Rem85, p. 45, pl. 12, figs. 2, 3a-b

- Deinsdorfia janossyi* REUMER, 1985
 paratypes, V 2008.7.4.
 four upper molars
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Rem85, p. 45
- Desmana kormosi* SCHREUDER, 1940
 holotype, V.61.2197
 left maxilla fragment with P⁴-M³
 Middle Pliocene
 quarry, Beremend
 ScA40, p. 314, text-figs. 8a-b, 19, 28
 (text-figs. 20, 40, 48a-b, 55a-d, 70, 75, 83,
 pl. 10, figs. 1-2, pl. 11, figs. 5, 7, 20)
- Desmana kormosi* SCHREUDER, 1940
 paratype, V.61.1576
 right mandible with M₂
 Middle Pliocene
 quarry, Beremend
 ScA40, p. 314, text-fig. 32a, pl. 10, fig. 3
- Desmana kormosi* SCHREUDER, 1940
 paratype, V.61.2197
 right humerus
 Middle Pliocene
 quarry, Beremend
 ScA40, p. 314, pl. 11, fig. 6
- Desmana kormosi* SCHREUDER, 1940
 paratypes, V.61.2197
 two left femurs
 Middle Pliocene
 quarry, Beremend
 ScA40, p. 314, pl. 11, figs. 15a-b
- Desmana kormosi* SCHREUDER, 1940
 paratypes, V.61.2199
 two mandible fragments and three limb bones
 Middle Pliocene
 quarry, Beremend
 ScA40, p. 314
- Dicerorhinus orientalis hungaricus* GAÁL, 1938
 holotype, V.60.1755
 left mandible
 Pannonian, Miocene
 brickyard, Strázsza-hegy, Hatvan
 Gaá38b, p. 130, text-figs. 2, 3, 4b (text-fig. 5)
Dicerorhinus hungaricus, Gaá43
 Other remains of same individual, listed in
 GAÁL 1943, not found
- Dolomys dalmatinus* KORMOS, 1931
 paratypes, V.61.1496
 left and right mandible fragments
 Early Pleistocene
 Podumci, Croatia
 Krm31, p. 130, text-fig. 6b (text-figs. 4b,
 5a-b, 7)
Dinaromys dalmatinus, Kre55
- Dolomys dalmatinus* KORMOS, 1931
 paratype, V.61.1538
 juvenile right M₁
 Early Pleistocene
 Podumci, Croatia
 Krm31, p. 130, text-fig. 6c
Dinaromys dalmatinus, Kre55
- Dolomys episcopalis bolkayi* KORMOS, 1931
 paratypes, V.61.1536
 two left M³
 Early Pleistocene
 Podumci, Croatia
 Krm31, p. 124, text-fig. 1a (text-fig. 3a)
Pliomys bolkayi, Rab83
- Dolomys episcopalis bolkayi* KORMOS, 1931
 paratypes, V.61.1545
 left and right mandible fragments
 Early Pleistocene
 Podumci, Croatia
 Krm31, p. 124
Pliomys bolkayi, Rab83
- Dolomys hungaricus* KORMOS, 1934
 holotype, V.61.1533
 left mandible
 Middle Pliocene
 upper quarry, Csarnóta
 Krm34a, p. 315, text-fig. 46
Propliomys hungaricus, Kre59
 Inventory number in original publication:
 Nr. 3799
- Dolomys hungaricus* KORMOS, 1934
 paratype, V.61.2189
 left mandible
 Middle Pliocene
 upper quarry, Csarnóta
 Krm34a, p. 315
Propliomys hungaricus, Kre59

Dolomys milleri NEHRING, 1898

syntypes, V.61.1520

maxilla and mandible fragments, isolated teeth

Middle Pliocene

quarry, Beremend

Neh98, p. 16 (text-figs. 1-3)

Dolomys milleri NEHRING, 1898

syntypes, V.61.1534

left and right mandible fragments

Middle Pliocene

quarry, Beremend

Neh98, p. 16

Dolomys nehringi KRETZOI, 1959

holotype, V.61.2193

right M₁

Middle Pliocene

quarry, locality 2, Csarnóta

Kre59, p. 242

Type figured in KRETZOI 1962: p. 310,
text-fig. 4a*Dystychoceras pannoniae* KRETZOI, 1941

holotype

both horn cores with fragment of skull

Sarmatian, Miocene; Kozárd Formation

Boór sand pit, Sopron

Kre41e, p. 336 (262), text-fig. 1, pl. 8

Miotragocerus pannoniae, Spa04

Type missing, not found in inventory

Eliomys quercinus helleri JÁNOSSY, 1962

holotype, V.64.483

right P₄

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 166, pl. 2, fig. 5

Eomellivora orlovi KRETZOI, 1965

holotype, V.64.1029

right maxilla fragments with P¹ and P^{3-P⁴}

of same individual

Pannonian, Miocene; Torony Lignite

Formation?

brickyard, Győrszentmárton, Pannonhalma

Kre65, p. 130, pl. 1, figs. 4-7

Episoriculus borsodensis JÁNOSSY, 1973

holotype, V.72.115

left maxilla fragment

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Ján73, p. 53, pl. 1, fig. 5 (pl. 1, figs. 6, 13)

Asoriculus gibberodon, Més99

Type refigured in REUMER 1985: p. 87,

pl. 27, fig. 6

Episoriculus borsodensis JÁNOSSY, 1973

paratype, V.72.123

left mandible fragment

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Ján73, p. 53, pl. 1, fig. 9

Asoriculus gibberodon, Més99

Type figured in JÁNOSSY 1972a: p. 51,

pl. 2, fig. 3

Episoriculus tornensis JÁNOSSY, 1973

holotype, V.72.114

viscerocranium with full dentition

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Ján73, p. 50, pl. 1, fig. 1

Asoriculus gibberodon, Més99

Type refigured in REUMER 1985: p. 87,

pl. 27, fig. 5

Episoriculus tornensis JÁNOSSY, 1973

paratype, V.72.122

left mandible with full dentition

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Ján73, p. 50, pl. 1, figs. 10-12

Asoriculus gibberodon, Més99*Erinaceus lechei* KORMOS, 1934

holotype, V.61.1558

right mandible

Middle Pliocene

quarry, Beremend

Krm34a, p. 296, text-fig. 31

Inventory number in original publication:

Nr. 3954

- Erinaceus ostramosi* JÁNOSSY, 1972
holotype, V.72.1
right mandible with P₄ and M₂
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Ján72b, p. 171
- Erinaceus ostramosi* JÁNOSSY, 1972
paratypes, V.73.78
maxilla fragment without teeth, right M²,
left M³, I₁ and isolated right M₁
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Ján72b, p. 171
- Estramomys simplex* JÁNOSSY, 1969
holotype, V.69.1227
right mandible with P₄-M₃
Early Pleistocene
Osztramos, locality 3, Tornaszentandrás
Ján69a, p. 36 (15)
Type figured in JÁNOSSY 1970: p. 113,
pl. 1, fig. 3
- Estramomys simplex* JÁNOSSY, 1969
paratype, V.69.1228
left maxilla fragment with P⁴-M³
Early Pleistocene
Osztramos, locality 3, Tornaszentandrás
Ján69a, p. 36 (15)
Type figured in JÁNOSSY 1970: p. 113,
pl. 1, fig. 1
- Foetorius palermineus* PETÉNYI, 1864
syntypes
right mandible fragment with P₄, M₁-M₂
and isolated right lower canine
Middle Pliocene
quarry, Beremend
Pet64, p. 50, pl. 1, figs. 3h-1
Mustela palerminea, Hel30; Rab76
Types missing, not found in inventory
- Glis antiquus* KORMOS, 1930
holotype, V.61.1561
maxilla fragment
Early Pleistocene
Somlyó-hegy, Püspökfürdő, Oradea
(Nagyvárad), Romania
Krm30, p. 242
- Hemilophodon osborni* KRETZOI, 1942
holotype
left mandible fragment with M₃
Burdigalian, Miocene
Sansan, France
Kre42b, p. 139, text-fig. 1
Type probably destroyed in 1956
- Hubacyon (Kanicyon) pannonicus* KRETZOI,
1985
holotype, V.84.209
left M₁
Pannonian, Miocene; Kálla Formation
sand pit, Danitz-puszta, Pécs
Kre85, p. 66, text-figs. 2, 4-6
- Hypsolophiodon csobánkanus* KRETZOI, 1940
holotype, V.60.162
right mandible fragment with P₂-P₄
Kiscellian, Oligocene; Hárshegy Sandstone
Formation
Csobánka
Kre40, p. 95, pl. 3, fig. 6
- Hystrix vinogradovi atavus* JÁNOSSY, 1972
holotype, V.72.2
left mandible with full dentition
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Ján72b, p. 173, pl. 1, figs. 1-2 (pl. 1,
figs. 5-9)
- Hystrix vinogradovi atavus* JÁNOSSY, 1972
paratype, V.73.69
left mandible with D₄-M₃
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Ján72b, p. 173, pl. 1, figs. 3-4
Type missing

Karstocricetus skofleki KORDOS, 1987

holotype, V.86.2

left mandible fragment with M₁–M₃

Turolian, Miocene

fissure fill in limestone quarry, Tardosbánya,

Tardos

Krd87, p. 75, pl. 1, fig. 1 (text-figs. 8, 9, pl. 1,

figs. 2–8, pl. 2, figs. 1–8)

Type missing

Karstocricetus skofleki KORDOS, 1987

paratype, V.86.3

right maxilla fragment with M¹–M³

Turolian, Miocene

fissure fill in limestone quarry, Tardosbánya,

Tardos

Krd87, p. 75

Type missing

Lagotona lázári KRETZOI, 1941

holotype, V.59.1063

mandible with P₃–M₃

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög),

Slovakia

Kre41a, p. 111, text-fig. 2i

Lagotona lázári KRETZOI, 1941

paratypes, V.59.1063

three upper teeth

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög),

Slovakia

Kre41a, p. 111

Lagurus pannonicus KORMOS, 1930

paratype, V.61.1445

juvenile M₁

Early Pleistocene

Somlyó-hegy, Püspökkfürdő, Oradea

(Nagyvárad), Romania

Krm30, p. 244

Prolagurus pannonicus, Rab81

Lagurus pannonicus KORMOS, 1930

paratypes, V 2008.6.1.

seven mandibles

Early Pleistocene

Somlyó-hegy, Püspökkfürdő, Oradea

(Nagyvárad), Romania

Krm30, p. 244

Prolagurus pannonicus, Rab81

Lagurus pannonicus KORMOS, 1930

paratype, V 2008.6.2.

maxilla fragment

Early Pleistocene

Somlyó-hegy, Püspökkfürdő, Oradea

(Nagyvárad), Romania

Krm30, p. 244

Prolagurus pannonicus, Rab81

Lagurus transiens JÁNOSSY, 1962

holotype, V.64.257

left M₁

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 168, text-fig. 2b (text-fig. 2d)

Lagurus transiens JÁNOSSY, 1962

paratypes, V.64.236

molars

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 168

Lagurus transiens JÁNOSSY, 1962

paratype, V.64.257

M₁

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 168, text-fig. 2c

Lagurus transiens JÁNOSSY, 1962

paratypes, V.64.286, V.64.448

M₁ and other molars

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 168

Lagurus transiens JÁNOSSY, 1962

paratype, V.64.529

M₁

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Ján62, p. 168

Leo gombaszögensis KRETZOI, 1938

holotype, 2007.94.1.

left P⁴

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög),

Slovakia

Kre38, p. 100, pl. 1, figs. 2–3

Panthera onca gombaszogensis, Hem01

Inventory number in original publication:

V. 991

- Leo gombaszögensis* KRETZOI, 1938
 paratype, V.59.1039
 left P_4
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100, pl. 1, fig. 5
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 W. 51
- Leo gombaszögensis* KRETZOI, 1938
 paratype, V.59.1041
 right P_3
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100, pl. 1, fig. 4
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 Fa. 24
- Leo gombaszögensis* KRETZOI, 1938
 paratype, V.59.1044
 left lower canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 Fa. 53
- Leo gombaszögensis* KRETZOI, 1938
 paratype, V.59.1084
 left M_1
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100, pl. 1, fig. 6
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication: W. 9
- Leo gombaszögensis* KRETZOI, 1938
 paratype, V.59.1085
 left M_1
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100, pl. 1, fig. 7
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 V. 915
- Leo gombaszögensis* KRETZOI, 1938
 paratypes, V.60.1776
 P_4 , P^3
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 Fa. 54
- Leo gombaszögensis* KRETZOI, 1938
 paratype
 right upper canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 100, pl. 1, fig. 1
Panthera onca gombaszoegensis, Hem01
 Inventory number in original publication:
 Fa. 23. Type missing, not found in current inventory
- Leptodontomys bodvanus* JÁNOSSY, 1972
 holotype, V.71.152
 right mandible
 Middle Pliocene
 Osztramos, locality 1, Tornaszentandrás Ján72a, p. 41, pl. 3, fig. 1 (pl. 3, figs. 2-5)
 Inventory number quoted erroneously in original publication: V.71.52
- Leptodontomys bodvanus* JÁNOSSY, 1972
 paratypes, V.73.22
 right M^1 and M^2
 Middle Pliocene
 Osztramos, locality 1, Tornaszentandrás Ján72a, p. 41
- Lepus praetimidus* KRETZOI in JÁNOSSY, 1969
 holotype, 2007.93.1.
 skull
 Middle Pleistocene
 Ördöglyuk-barlang, Solymár Ján69b, p. 610, pl. 6, figs. 1-2
Lepus timidus praetimidus, F-F06
 Type refigured in FOSTOWICZ-FRELIK & GASPARIK 2006: p. 153, figs. 1a-g

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.148

skull

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268) (text-fig. 1)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.149

right femur

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.150

left femur fragment

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.151

right tibia fragment

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Type lost

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.152

left tibia, calcaneus, tarsus

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.153

right humerus

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.154

left humerus

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.155

three thoracic vertebrae

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.156

left and right pelvis fragments

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.157

radius and ulna

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.158

atlas fragment

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

Mammonteus hungaricus KRETZOI, 1941

paratype, V.79.159

epistropheus

Late Pleistocene

Besenyői szőlők, Zalaegerszeg

Kre41f, p. 343 (268)

Mammuthus primigenius hungaricus, Völ81

- Mammonteus hungaricus* KRETZOI, 1941
 paratype, V.79.160
 two carpal
 Late Pleistocene
 Besenyői szőlők, Zalaegerszeg
 Kre41f, p. 343 (268)
Mammuthus primigenius hungaricus, VöI81
- Manis hungarica* KORMOS, 1934
 holotype
 ungual phalanx of right third digit
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34d, p. 91, text-figs. 1.1a-c
 Inventory number in original publication:
 Nr. 3932. Type missing, not found in current
 inventory
- Mastodon (Mammut) americanus praetypica*
 SCHLESINGER, 1922
 syntype, V.93.5
 left mandible with M₂-M₃
 Early Pliocene
 Százhalombatta-Érd
 ScG22, p. 115, 227, pl. 15, fig. 4 (pl. 15, fig. 1,
 pl. 19, fig. 2)
Mammut borsoni, Gas07
- Mastodon (Mammut) americanus praetypica*
 SCHLESINGER, 1922
 syntype, V.93.9
 left mandible fragment with M₃ and
 fragmentary M₂
 Middle Pliocene
 Hajnáčka (Ajnácskő), Slovakia
 ScG22, p. 115, 227, pl. 16, fig. 2
Mammut borsoni, Gas07
- Mastodon (Mammut) americanus praetypica*
 SCHLESINGER, 1922
 syntype
 right M²
 Late Pliocene
 Rákoskeresztúr, Budapest
 ScG22, p. 115, 227, pl. 14, fig. 5
Mammut borsoni, Gas07
 Type lost
- Mastodon (Mammut) americanus praetypica*
 SCHLESINGER, 1922
 syntypes
 right M²-M³ and right M₂-M₃ of same
 individual
 age unknown
 Subotica (Szabadka), Serbia
 ScG22, p. 115, 227, pl. 15, figs. 2, 3
Mammut borsoni, Gas07
 Types lost
- Mastodon (Mammut) americanus praetypica*
 SCHLESINGER, 1922
 syntype
 right mandible fragment with M₃
 Middle Pliocene
 Hajnáčka (Ajnácskő), Slovakia
 ScG22, p. 115, 227, pl. 16, fig. 1, pl. 17, fig. 1,
 pl. 18, fig. 5
Mammut borsoni, Gas07
 Type lost
- Mastodon (Bunolophodon) angustidens*
subtapiridea SCHLESINGER, 1917
 paratype, V.59.1134
 left M₃ fragment
 Sarmatian, Miocene
 Kőbánya, Budapest
 ScG17, p. 37 (pl. 3, fig. 2, pl. 4, figs. 1-2,
 pl. 5, figs. 1-2, pl. 6, figs. 1-2, pl. 7, figs. 1-3,
 pl. 8, figs. 1-2)
Gomphotherium angustidens, Gas07
 Type figured in SCHLESINGER 1922: pl. 7,
 fig. 9
- Mastodon (Bunolophodon) angustidens*
subtapiridea SCHLESINGER, 1917
 paratype, V.59.1145
 left M₂ fragment
 Sarmatian, Miocene
 Kőbánya, Budapest
 ScG17, p. 37
Gomphotherium angustidens, Gas07

Mastodon (Bunolophodon) grandincisivum
SCHLESINGER, 1917
syntype, V.79.34
one upper and two lower tusks, left and right
 M^3 , mandible fragments with left and right M_3 ,
postcranial skeletal parts, right scapula
fragment, humerus fragments, ulna and radius
fragments, left and right magnum, right
trapezoideum, right intermedium, right
unciforme, metapodium fragment, phalanx
fragment, left and right pelvis fragments,
femur fragments, right tibia, patella, calcaneus,
rib fragments, vertebrae
Pannonian, Miocene
Pestszentlőrinc, Budapest
ScG17, p. 119 (pl. 15, figs. 1-2, pl. 34, figs. 1-2)
"Mastodon" *grandincisivus*, Gas07
Type figured and described in detail in
SCHLESINGER 1922: p. 7, pl. 1, figs. 1-2, pl. 2,
figs. 1-3, pl. 3, figs. 1-7, pl. 4, figs. 1-3, pl. 5,
figs. 1-5, pl. 6, fig. 1

Mediocris commenticius KAZÁR, 2005
holotype, V.93.2
incomplete skeleton
Sarmatian, Miocene; Kozárd Formation
northwest of Herman Ottó-tó, Kovácszénája
Kaz05, p. 55, text-figs. 3-5, 7-10, 12
Sophianaecetus commenticius, Kaz06

Megaderma janossyi TOPÁL, 1974
holotype, V.74.57
left mandible with broken C and M_1
Middle Pliocene
Osztramos, locality 10, Tornaszentandrás
Top74, p. 96, text-figs. 1a-c

Mesocetus hungaricus KADIĆ, 1907
holotype, V.79.118
incomplete skeleton with skull
Badenian, Miocene; Baden Clay Formation
clay pit, Walbersdorf (Borbolya), Austria
Kad07b (Kad07a), p. 31 (28), text-figs. 3-70,
pl. 1, pl. 2, figs. 1-3, pl. 3, figs. 1-3
Some parts of original find lost

Microtus coronensis KORMOS, 1933
holotype, V.61.1426
left mandible
Early Pleistocene
Dealul Sprenghi (Fortyogó-hegy,
Gesprengberg), Brașov (Brassó), Romania
Krm33a, p. 11, text-fig. 2

Microtus coronensis KORMOS, 1933
paratypes, V.61.1426
three left mandibles
Early Pleistocene
Dealul Sprenghi (Fortyogó-hegy,
Gesprengberg), Brașov (Brassó), Romania
Krm33a, p. 11, text-fig. 3

Microtus coronensis KORMOS, 1933
paratype, V.61.1431
right maxilla
Early Pleistocene
Dealul Sprenghi (Fortyogó-hegy,
Gesprengberg), Brașov (Brassó), Romania
Krm33a, p. 11

Mimomys fejérváryi KORMOS, 1934
holotype, V.61.1514
right mandible
Middle Pleistocene
Nagyharsány-hegy, Villány
Krm34a, p. 317, text-fig. 47
Borsodia fejervaryi, Rab81
Inventory number in original publication:
Nr. 3804

Mimomys ostramosensis JÁNOSSY & VAN DER
MEULEN, 1975
holotype, V.73.2
viscerocranum and left mandible of same
individual
Late Pliocene
Osztramos, locality 3, Tornaszentandrás
Ján75, p. 382, pl. 1, figs. 1a-b

Mimomys ostramosensis JÁNOSSY & VAN DER
MEULEN, 1975
paratypes, V.73.50
cranial fragments, maxillae, mandibles and
isolated M_1
Late Pliocene
Osztramos, locality 3, Tornaszentandrás
Ján75, p. 382, pl. 1, figs. 2a-b, 3a-b

Mimomys petényii MÉHELY, 1914
lectotype, V.73.3
left mandible with full dentition
Middle Pliocene
quarry, Beremend
Méh14a (Méh14b), p. 191 (42), pl. 4, fig. 8
Borsodia petenyii, Rab81
Lectotype designated in JÁNOSSY & VAN DER
MEULEN 1975: p. 390

Mimomys petényii MÉHELY, 1914
 paralectotypes, V.61.1418
 two maxilla fragments, five mandible
 fragments, one upper molar, three lower
 molars
 Middle Pliocene
 quarry, Beremend
Méh14a (*Méh14b*), p. 191 (42), pl. 4, figs. 5–7
Borsodia petenyii, Rab81

Mimomys pitymyoides JÁNOSSY & VAN DER
 MEULEN, 1975
 holotype, V.73.4
 nearly complete skull
 Late Pliocene
 Osztramos, locality 3, Tornaszentandrás
Ján75, p. 386, pl. 2, figs. 8a–b, 9a–b

Mimomys pitymyoides JÁNOSSY & VAN DER
 MEULEN, 1975
 paratypes, V.73.45
 viscerocrania, cranial fragments, maxilla
 fragments, mandibles and isolated M_1
 Late Pliocene
 Osztramos, locality 3, Tornaszentandrás
Ján75, p. 386

Mimomys rex KORMOS, 1934
 holotype, V.61.1473
 right mandible
 Middle Pleistocene
 Templom-hegy, Villány
Krm34a, p. 319, text-fig. 49
 Inventory number in original publication:
 Nr. 3802/4

Mimomys silasensis JÁNOSSY, 1974
 holotype, V.74.4
 right mandible fragment with M_1
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
Ján74a, p. 22, pl. 1, figs. c–d

Mimomys tornensis JÁNOSSY & VAN DER
 MEULEN, 1975
 holotype, V.73.5
 viscerocranium fragment and left mandible
 of same individual
 Late Pliocene
 Osztramos, locality 3, Tornaszentandrás
Ján75, p. 385, pl. 1, figs. 4a–b

Mimomys tornensis JÁNOSSY & VAN DER
 MEULEN, 1975
 paratypes, V.73.54
 cranial fragments, maxilla fragments,
 mandibles and isolated M_1
 Late Pliocene
 Osztramos, locality 3, Tornaszentandrás
Ján75, p. 385, pl. 1, figs. 5–7

Muscardinus dacicus KORMOS, 1930
 holotype, V.61.1564
 left maxilla fragment with full dentition
 Early Pleistocene
 Somlyó-hegy, Püspökkürtő, Oradea
 (Nagyvárad), Romania
Krm30, p. 243

Muscardinus dacicus KORMOS, 1930
 paratype, V.61.1564
 right mandible
 Early Pleistocene
 Somlyó-hegy, Püspökkürtő, Oradea
 (Nagyvárad), Romania
Krm30, p. 243

Muscardinus dacicus KORMOS, 1930
 paratypes, V 2008.5.1.
 nine mandibles
 Early Pleistocene
 Somlyó-hegy, Püspökkürtő, Oradea
 (Nagyvárad), Romania
Krm30, p. 243

Muscardinus giganteus JÁNOSSY, 1974
 holotype, V.74.2
 right M^1
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
Ján74a, p. 20, pl. 1, fig. f

Mustela beremendensis PETÉNYI in KUBINYI,
 1856
 holotype
 right mandible fragment with P_4 , M_1 – M_2
 Middle Pliocene
 quarry, Beremend
Kub56, p. 72
Baranogale beremendensis, Kre42c
 Type figured and described in detail in
 PETÉNYI 1864: p. 48, pl. 1, figs. 2f–g. Type
 missing, not found in inventory

Mustela martelina PETÉNYI, 1864

holotype

right lower carnassial tooth

Middle Pliocene

quarry, Beremend

Pet64, p. 44, pl. 1, figs. 1a-c

Paratanuki martelinus, Kre56

Type missing, not found in inventory

Mustela praenivalis KORMOS, 1934

holotype, V.61.1384

right mandible fragment

Middle Pleistocene

Nagyharsány-hegy, Villány

Krm34b, p. 154, pl. 2, fig. 12

Inventory number in original publication:

Nr. 3930

Mustela praenivalis KORMOS, 1934

paratype, V.61.1383

right mandible fragment

Middle Pleistocene

Sackdilling, Auerbach, Germany

Krm34b, p. 154

Myotis baranensis KORMOS, 1934

holotype, V.61.1403

left mandible

Middle Pliocene

quarry, Beremend

Krm34a, p. 306, text-fig. 39

Inventory number in original publication:

Nr. 3851

Myotis bechsteini robustus TOPÁL, 1963

holotype, V.63.346

right mandible with C, P₂ and M₁-M₃

Middle Pleistocene

Kövesvárad-hegy, Répáshuta

Top63a, p. 146

Myotis bechsteini robustus TOPÁL, 1963

paratypes, V.63.333

maxilla fragments and upper teeth

Middle Pleistocene

Kövesvárad-hegy, Répáshuta

Top63a, p. 146

Myotis bechsteini robustus TOPÁL, 1963

paratypes, V.63.343

limb bones and other fragments

Middle Pleistocene

Kövesvárad-hegy, Répáshuta

Top63a, p. 146

Myotis bechsteini robustus TOPÁL, 1963

paratypes, V.63.344

mandible fragments and lower teeth

Middle Pleistocene

Kövesvárad-hegy, Répáshuta

Top63a, p. 146

Myotis estramosensis TOPÁL, 1983

holotype, V.82.173

injured left mandible with P₄-M₃

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Top83, p. 51, pl. 1, fig. 8

Myotis estramosensis TOPÁL, 1983

paratype, V.82.177

right maxilla fragment with M¹-M³

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Top83, p. 51

Myotis janossyi TOPÁL, 1983

holotype, V.82.169

well-preserved skull

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Top83, p. 46, pl. 1, figs. 4-6

Myotis janossyi TOPÁL, 1983

paratype, V.82.170

right mandible with full dentition

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Top83, p. 46, pl. 1, fig. 7

Myotis kretzoi TOPÁL, 1981

holotype, V.81.2

left mandible with full dentition except P₂

Middle Pleistocene

Ördöglyuk-barlang, Solymár

Top81, p. 59, pl. 1, figs. 4-5

Myotis kretzoii TOPÁL, 1981

paratype, V.81.3

skull with P⁴-M₃

Middle Pleistocene

Ördöglyuk-barlang, Solymár

Top81, p. 59, pl. 1, figs. 1-3

Myotis paradaubentoni TOPÁL, 1983

holotype, V.82.171

full rostrum with entire dentition

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Top83, p. 49, pl. 2, figs. 4-6

Myotis paradaubentoni TOPÁL, 1983

paratype, V.82.172

left mandible with full dentition

Middle Pliocene

Osztramos, locality 13, Tornaszentandrás

Top83, p. 49, pl. 2, fig. 7

Myotis schaubi KORMOS, 1934

holotype, V.61.1416

left mandible

Middle Pleistocene

Templom-hegy, Villány

Krm34a, p. 309, text-fig. 41

Inventory number in original publication:

Nr. 3849

Myotis steiningeri KORMOS, 1934

holotype, V.61.1414

left mandible

Middle Pleistocene

Templom-hegy, Villány

Krm34a, p. 308, text-fig. 40

Inventory number in original publication:

Nr. 3850

Myotis wüsti KORMOS, 1934

holotype, V.61.1412

right mandible

Middle Pleistocene

Nagyharsány-hegy, Villány

Krm34a, p. 310, text-fig. 42

Inventory number in original publication:

Nr. 3848. Locality on specimen label: Kalkberg
(Templom-hegy in current usage)

Neohyaenodon semseyi KRETZOI, 1941

holotype, V.60.1725

skull with mandible

Oligocene

Badlands, Dakota, USA

Kre41c, p. 172, text-fig. 1

Nesodon imbricatus maior KRETZOI, 1941

holotype

skull fragment

age unknown

Patagonia, Argentina

Kre41c, p. 170

Type probably destroyed in 1956

Pachycrocuta robusta progressa KRETZOI, 1938

holotype

left mandible with C-M₁

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög), Slovakia

Kre38, p. 121, pl. 1, figs. 13-14

Inventory number in original publication:
Fa. 18. Type missing, not found in current inventory

Pachycrocuta robusta progressa KRETZOI, 1938

paratype, V.59.931

left P₂

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög), Slovakia

Kre38, p. 121

Inventory number in original publication:
Fa. 46

Pachycrocuta robusta progressa KRETZOI, 1938

paratype, V.59.963

crown of left lower canine

Early Pleistocene

fissure fill in quarry, Gombasek (Gombaszög), Slovakia

Kre38, p. 121

Inventory number in original publication:
Fa. 49

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.971
 left P₄ fragment
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 44

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.971
 right P₄ fragment
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 45

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.986
 left I³
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 50

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.996
 crown of left upper canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 47. Type missing

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.996
 tip of crown of left upper canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 48

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.1020
 left P₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 Fa. 43

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.1038
 left P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121
 Inventory number in original publication:
 V. 810

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.59.1066
 right P⁴
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121, pl. 2, figs. 3–4
 Inventory number in original publication:
 Fa. 26

Pachycrocuta robusta progressa KRETZOI, 1938
 paratype, V.60.1775
 left mandible fragment with P₄–M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 121, pl. 2, figs. 1–2
 Inventory number in original publication:
 Fa. 25

Pachyura pannonica KORMOS, 1934
 holotype
 left mandible with full dentition
 Middle Pliocene
 quarry, Beremend
 Krm34a, p. 306, text-fig. 38
Paenelimnoecus pannonicus, Rem85
 Inventory number in original publication:
 Nr. 3960. Type missing, not found in current inventory

- Pachyura pannonica* KORMOS, 1934
 paratypes
 two mandible fragments
 Middle Pliocene
 quarry, Beremend
 Krm34a, p. 306
Paenelimnoecus pannonicus, Rem85
 Types missing, not found in inventory
- Pannonictis? janossyi* RABEDER, 1976
 holotype, V.61.1390
 skull fragment with left P²–P⁴, M¹ and right P³
 Middle Pleistocene
 Templom-hegy, locality 3 or 5, Villány
 Rab76, p. 71, text-fig. 30
 Other fragment of skull under same
 inventory number questionably referred
 to this species
- Pannonictis piligrimi* KORMOS, 1934
 paratypes, V.61.1363
 right mandible fragment, right upper and
 lower canine
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34b, p. 131, pl. 2, figs. 3a–b (pl. 2, fig. 1)
Pannonictis ardea, Rab76
 Inventory number in original publication:
 Nr. 3916
- Pannonictis piligrimi* KORMOS, 1934
 paratypes, V.61.1366
 skull fragments
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34b, p. 131
Pannonictis ardea, Rab76
- Pannonictis piligrimi* KORMOS, 1934
 paratypes, V.61.1368
 skull and both mandibles of same individual
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34b, p. 131
Pannonictis ardea, Rab76
- Pannonictis piligrimi* KORMOS, 1934
 paratype, V.61.1370
 right maxilla fragment
 Middle Pliocene
 quarry, Beremend
 Krm34b, p. 131
Pannonictis ardea, Rab76
- Pannonictis piligrimi* KORMOS, 1934
 paratypes, V.61.2169
 two upper canine teeth
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34b, p. 131
Pannonictis ardea, Rab76
- Pentaglis földvárii* KRETZOI, 1943
 holotype
 right M¹
 Karpatian, Miocene; Fót Formation
 borehole 2403, Lövölde tér, Budapest
 Kre43a, p. 271, text-fig. 1
 Type missing, not found in inventory
- Pentaglis földvárii* KRETZOI, 1943
 paratype
 second phalanx
 Karpatian, Miocene; Fót Formation
 borehole 2403, Lövölde tér, Budapest
 Kre43a, p. 271
 Type missing, not found in inventory
- Peromyscus gyulavarii* KRETZOI & VÖRÖS, 1989
 holotype, V.88.1
 left M₂ lacking alveolar half of tooth column
 Late Pliocene
 near Mashpaya (Shipibo Indian village),
 in upper reaches of River Pisqui, Peru
 Kre89, p. 112, text-fig. 17, pl. 1, figs. 1–4
 An undescribed and unfigured fragment of
 holotype specimen found in KRETZOI's bequest

Petényia hungarica KORMOS, 1934
 paratypes, V.61.1556
 maxilla fragment and left mandible
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34a, p. 301 (text-figs. 34–35)

Petényia neglecta KRETZOI, 1943
 holotype, V.61.1593
 left mandible
 Early Pleistocene
 Somlyó-hegy, Püspökkfürdő, Oradea
 (Nagyvárad), Romania
 Kre43b, p. 607, unnumbered text-figure
Petenyia hungarica, Rem85
 Type refigured in REUMER 1985: p. 65,
 pl. 19, figs. 2a–c

Petényia neglecta KRETZOI, 1943
 paratype, V.61.1593
 left maxilla fragment with P^4-M^2
 Early Pleistocene
 Somlyó-hegy, Püspökkfürdő, Oradea
 (Nagyvárad), Romania
 Kre43b, p. 607
Petenyia hungarica, Rem85
 Type figured in REUMER 1985: p. 59, pl. 16,
 fig. 5. Another paratype (left mandible) missing

Phoca holitschensis BRÜHL, 1860
 holotype, V.60.1734
 left foot
 Miocene
 Holíč (Holics), Slovakia
 Brü60, p. 1, pl. 1, fig. 2, pl. 2, fig. 2
Praepusa vindobonensis, Kry01

Pitymys hintoni KRETZOI, 1941
 holotype
 left mandible fragment with M_1-M_2
 Early Pleistocene
 Betfia (Betfia), Romania
 Kre41d, p. 319 (249), text-fig. 3.1.
Microtus (Allophaiomys) hintoni, Rab81
 Type missing, not found in inventory

Plecotus (Plecotus) plioacaenicus TOPÁL, 1989
 holotype, V.86.14
 right mandible fragment with I_1-I_3 , C, P_2-P_4 , M_1
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top89, p. 82, text-figs. 7–8

Plecotus (Plecotus) plioacaenicus TOPÁL, 1989
 paratype, V.86.15
 left maxilla fragment with C– P^2 , P^4 , M^1-M^3
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top89, p. 82, text-fig. 9

Pliolagus beremendensis KORMOS, 1934
 holotype, V.61.1376
 left mandible fragment
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34c, p. 74 (66), text-figs. 1s–t (text-figs.
 1p–r)
Hypolagus beremendensis, F-F07
 Inventory number in original publication:
 Nr. 3931

Pliolagus tóthi KRETZOI, 1941
 syntypes
 teeth
 Early Pleistocene
 Betfia (Betfia), Romania
 Kre41d, p. 322 (250), text-figs. 6.1–2.
Hypolagus beremendensis, F-F07
 Types missing, not found in inventory

Pliomys posterior JÁNOSSY, 1969
 holotype, V.65.29
 right mandible fragment with M_1-M_2
 Middle Pleistocene
 rock-shelter I., Uppony
 Ján69b, p. 607, pl. 7, figs. 6–7

"Pliomys" progressus KRETZOI, 1938
 holotype, V.59.1054
 mandible
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 96, text-fig. 2k
 Inventory number in original publication:
 Fa. 17. Type missing

"Pliomys" progressus KRETZOI, 1938
 paratypes
 two mandibles
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 96
 Types missing, not found in inventory

- Praepusa pannonica* KRETZOI, 1941
 holotype
 left mandible with M₁
 Sarmatian, Miocene; Kozárd Formation
 Postás-telep, Érd
Kre41g, p. 351 (275), text-figs. 1.1–2.
 Type missing, not found in inventory
- Praepusa pannonica* KRETZOI, 1941
 paratype
 proximal and distal ends of left tibia
 Sarmatian, Miocene; Kozárd Formation
 Postás-telep, Érd
Kre41g, p. 351 (275), text-fig. 2.1.
 Type missing, not found in inventory
- Praepusa pannonica* KRETZOI, 1941
 paratype
 distal end of left fibula
 Sarmatian, Miocene; Kozárd Formation
 Postás-telep, Érd
Kre41g, p. 351 (275)
 Type missing, not found in inventory
- Procapra (Protetraceros) collispannoni* KRETZOI,
 1965
 holotype, V.64.1030
 right horn core
 Pannonian, Miocene
 brickyard, Győrszentmárton, Pannonhalma
Kre65, p. 134, pl. 1, figs. 1–2
- Procapra (Protetraceros) collispannoni* KRETZOI,
 1965
 paratypes, V.72.99
 maxillae, mandibles, molars
 Pannonian, Miocene
 brickyard, Győrszentmárton, Pannonhalma
Kre65, p. 134
- Procapra (Protetraceros) collispannoni* KRETZOI,
 1965
 paratypes, V.72.100
 horn cores
 Pannonian, Miocene
 brickyard, Győrszentmárton, Pannonhalma
Kre65, p. 134, pl. 1, fig. 3
- Procapra (Protetraceros) collispannoni* KRETZOI,
 1965
 paratype, V.72.102
 distal end of humerus
 Pannonian, Miocene
 brickyard, Győrszentmárton, Pannonhalma
Kre65, p. 134
- Prodeinotherium petenyii* VÖRÖS, 1989
 holotype, V.88.5
 corpus of right mandible with P₃–M₂
 Ottangian, Miocene; Zagyvapálfalva
 Formation
 coal mine, Putnok
Völ89, p. 101, text-fig. 1a, pl. 1, figs. 1–3
 (pl. 2, figs. 1–3, pl. 3, figs. 1–3, pl. 4, figs. 1–3,
 pl. 5, figs. 1–2)
Prodeinotherium hungaricum, Gas93; Gas01
- Prodeinotherium petenyii* VÖRÖS, 1989
 paratype
 tusk fragment
 Ottangian, Miocene; Zagyvapálfalva
 Formation
 Zsigmond shaft, Királd
Völ89, p. 101
Prodeinotherium hungaricum, Gas93; Gas01
 Type lost, already at time of paratype
 designation
- Prodinotherium hungaricum* ÉHIK, 1930
 syntypes
 partial skeleton
 Ottangian, Miocene; Zagyvapálfalva
 Formation
 coal mine, Kotyháza, Salgótarján
Éhi30, p. 1, pl. 1, figs. 1–7, pl. 2, figs. 1–3,
 pl. 3, figs. 1–7, pl. 4, figs. 1–2
 Spelling of genus name emended to
Prodeinotherium, following HARRIS 1973.
 Types lost (see VÖRÖS 1989)
- Prodinotherium hungaricum* ÉHIK, 1930
 syntypes
 right P₃, right M₂ and left M₃
 Ottangian, Miocene; Zagyvapálfalva
 Formation
 Zsigmond shaft, Királd
Éhi30, p. 1, pl. 1, figs. 4–7
 Types lost (see VÖRÖS 1989)

Promimomys microdon JÁNOSSY, 1974

holotype, V.74.3

right fragmentary M₁

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Ján74a, p. 21, pl. 1, figs. a–b

Prospalax kretzoi JÁNOSSY, 1972

holotype, V.71.150

right mandible

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Ján72a, p. 43

Inventory number quoted erroneously
in original publication: V.71.50

Putorius stromeri KORMOS, 1934

holotype, V.61.1388

right mandible fragment

Middle Pliocene

quarry, Beremend

Krm34b, p. 148, pl. 2, fig. 8a (pl. 2, fig. 8b)

Inventory number in original publication:
Nr. 3921

Rhinolophus estramontis TOPÁL, 1979

holotype, V.79.191

left mandible with full dentition except I₁

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Top79, p. 75, pl. 8, figs. 4–5

Rhinolophus estramontis TOPÁL, 1979

paratypes, V.79.175

right maxilla fragments

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Top79, p. 75, pl. 3, figs. 1–4

Rhinolophus estramontis TOPÁL, 1979

paratype, 2007.114.1.

right maxilla fragment

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Top79, p. 75

Rhinolophus estramontis TOPÁL, 1979

paratypes, 2007.114.2.

upper teeth

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Top79, p. 75

Rhinolophus euryale praeglacialis KORMOS,

1934

holotype, V.61.1405

right mandible

Middle Pliocene

quarry, Beremend

Krm34a, p. 313, text-fig. 44

Inventory number in original publication:

Nr. 3959

Rhinolophus ferrumequinum tarkoensis TOPÁL,

1979

holotype, V.79.189

injured rostrum without premaxillae
and I₁–I₂

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Top79, p. 92, pl. 6, fig. 6

Rhinolophus ferrumequinum tarkoensis TOPÁL,

1979

paratype, V.79.187

left maxilla fragment

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Top79, p. 92, pl. 6, figs. 4–5

Rhinolophus ferrumequinum tarkoensis TOPÁL,

1979

paratype, V.79.188

right maxilla fragment

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Top79, p. 92, pl. 6, figs. 2–3

Rhinolophus ferrumequinum tarkoensis TOPÁL,

1979

paratype, V.79.190

left maxilla fragment

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Top79, p. 92, pl. 6, fig. 9

Rhinolophus ferrumequinum tarkoensis TOPÁL,

1979

paratype, V.79.203

right mandible

Middle Pleistocene

Tar-kő, rock-shelter, Felsőtárkány

Top79, p. 92, pl. 12, figs. 6–7

- Rhinolophus ferrumequinum tarkoensis* TOPÁL,
1979
paratypes, V.79.204
left and right mandible fragments
Middle Pleistocene
Tar-kő, rock-shelter, Felsőtárkány
Top79, p. 92, pl. 12, figs. 8, 10
- Rhinolophus ferrumequinum tarkoensis* TOPÁL,
1979
paratype, V.79.205
left mandible fragment
Middle Pleistocene
Tar-kő, rock-shelter, Felsőtárkány
Top79, p. 92, pl. 12, fig. 9
- Rhinolophus macrorhinus* TOPÁL, 1963
holotype, V.63.1551
skull
Middle Pliocene
quarry, Beremend
Top63b, p. 219, text-figs. 1-3
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
holotype, V.79.198
right mandible with full dentition except I₁
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81, pl. 11, figs. 3-4
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, V.63.331
mandibles and lower teeth
Middle Pleistocene
Kövesvárad-hegy, Répáshuta
Top79, p. 81, pl. 10, figs. 10-11
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, V.79.183
rostra
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81, pl. 4, figs. 2-4, pl. 5, figs. 1-2
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, V.79.184
left and right maxilla fragments
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81, pl. 4, figs. 1, 5-8, pl. 5, figs. 3-4
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, V.79.199
left mandibles
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81, pl. 11, figs. 1-2, 5-6
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratype, 2007.115.1.
rostrum
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, 2007.115.2.
maxilla fragments
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81
- Rhinolophus macrorhinus anomalidens* TOPÁL,
1979
paratypes, 2007.115.3.
mandible fragments
Early Pleistocene
Osztramos, locality 8, Tornaszentandrás
Top79, p. 81
- Rhinolophus postdelphinensis* TOPÁL, 1979
holotype, V.79.170
left maxilla fragment with C, P² and P⁴
Middle Pliocene
Osztramos, locality 1, Tornaszentandrás
Top79, p. 80, pl. 1, fig. 10
- Rhinolophus variabilis* TOPÁL, 1975
holotype, V.75.95
left mandible with full dentition except I₁ and P₃
Middle Pliocene
Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 1
- Rhinolophus variabilis* TOPÁL, 1975
paratype, V.75.87
right upper canine
Middle Pliocene
Osztramos, locality 1, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratypes, V.75.88

upper teeth

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.89

mandible

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 4

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.90

mandible

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 5

Rhinolophus variabilis TOPÁL, 1975

paratypes, V.75.91

mandibles

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratypes, V.75.92

distal parts of right humeri

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.93

left M¹

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratypes, V.75.94

left and right mandible fragments, isolated

left and right side teeth

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.96

left mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 6

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.97

left mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 4, fig. 3

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.98

right mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 9

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.99

right mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 10

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.100

left mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 3, fig. 2

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.101

left mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 4, fig. 2

Rhinolophus variabilis TOPÁL, 1975

paratype, V.75.102

left mandible fragment

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás
Top75, p. 5, pl. 4, fig. 4

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.103
 left mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 4, fig. 6

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.104
 left mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 3, fig. 8

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.105
 left mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 4, fig. 5

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.106
 left mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 4, fig. 1

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.107
 right mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 3, fig. 3

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.108
 left mandible fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 3, fig. 7

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.109
 maxillae and isolated upper teeth
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.110
 rostrum
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 2, fig. 4

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.111
 right maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 2, fig. 1

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.112
 right maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 10

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.113
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 8

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.114
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 13

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.115
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 12

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.116
 right maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 7

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.117
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 9

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.118
 right maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 11

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.119
 right maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 6

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.120
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 2, fig. 3

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.121
 left maxilla fragment
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 2, fig. 2

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.122
 premaxilla
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 1, fig. 1

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.123
 premaxillae
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.124
 proximal ends of humeri
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.125
 distal end of humerus
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 6, fig. 6

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.126
 distal end of humerus
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 6, figs. 4, 10

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.127
 distal end of humerus
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 6, figs. 2, 9

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.128
 distal end of humerus
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 6, figs. 5, 8

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.129
 distal end of humerus
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 6, fig. 3

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.130
 distal ends of humeri
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.131
 proximal fragments of radii
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.132
 distal fragment of radius
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.133
 claviculae
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.134
 cochlea
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratypes, V.75.135
 baculae
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.136
 bacula
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 7, figs. 1, 2, 5, 7

Rhinolophus variabilis TOPÁL, 1975
 paratype, V.75.137
 bacula
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
 Top75, p. 5, pl. 7, figs. 3, 4, 6, 8

Sciurus whitei hungaricus JÁNOSSY, 1962
 holotype, V.64.392
 right P⁴
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165, pl. 2, fig. 3

Sciurus whitei hungaricus JÁNOSSY, 1962
 paratype, V.64.392
 fragmentary M¹
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165

Sciurus whitei hungaricus JÁNOSSY, 1962
 paratypes, V.64.407
 two M₁
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165, pl. 2, fig. 4

Sciurus whitei hungaricus JÁNOSSY, 1962
 paratypes, V.64.486
 humerus, ulna, femur
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165, pl. 1, figs. 4, 5

Sciurus whitei hungaricus JÁNOSSY, 1962
 paratype, V.64.522
 right P⁴
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165

Sciurus whitei hungaricus JÁNOSSY, 1962
 paratype, V.67.168
 maxilla fragment
 Middle Pleistocene
 Tar-kő, rock-shelter, Felsőtárkány
 Ján62, p. 165

Sicista praeloriger KORMOS, 1930
 holotype, V.61.1569
 left mandible with full dentition
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm30, p. 241

Sicista praeloriger KORMOS, 1930

paratype, V.61.1567

maxilla fragment with M¹

Early Pleistocene

Somlyó-hegy, Püspökfürdő, Oradea
(Nagyvárad), Romania

Krm30, p. 241

Sicista praeloriger KORMOS, 1930

paratype, V.61.1568

left mandible fragment with M₁-M₃

Early Pleistocene

Somlyó-hegy, Püspökfürdő, Oradea
(Nagyvárad), Romania

Krm30, p. 241

Siphneus arvicolinus NEHRING, 1883

holotype

right mandible fragment

Pliocene

Kuei-te, Upper Hoang-ho River, Kansu, China

Neh83, p. 19, text-figs. a-c

Allosiphneus arvicolinus, Kre61

Type missing, not found in inventory;
old inventory number in KRETZOI 1961:

X. 540 / V. 26

Sirenavus hungaricus KRETZOI, 1941

holotype, V.60.1712

skull

Lutetian, Eocene; Szőc Limestone

Kálvária-hegy, Felsőgalla, Tatabánya

Kre41b, p. 147, text-fig. 1b

Type redescribed and photographed in
KORDOS 1981: p. 75, text-fig. 1, pl. 1, figs. 1-4

Sirenavus hungaricus KRETZOI, 1941

holotype, V.83.42

left mandible fragment with posterior part
of M₃

Lutetian, Eocene; Szőc Limestone

Kálvária-hegy, Felsőgalla, Tatabánya

Kre41b, p. 147

Type redescribed and photographed in
KORDOS 1981: p. 75, pl. 1, figs. 5-6. Previously
covered right mandible prepared from same
specimen (figs. 7-8)

Sorex araneus macrognathus JÁNOSSY, 1965

syntypes, V.64.902

28 mandibles

Middle Pleistocene

rock-shelter I., Uppony

Ján65, p. 64

Inventory number quoted for holotype
in original description but this lot contains
28 specimens, considered syntypes here,
whose measurements shown in JÁNOSSY 1965,
text-fig. 5

Sorex bor REUMER, 1985

holotype, V.83.12

right mandible with full dentition

Middle Pliocene

Osztramos, locality 9, Tornaszentandrás

Rem85, p. 32, pl. 7, figs. 4a-b (pl. 7, figs. 1, 3)

Sorex bor REUMER, 1985

paratype, V.73.19

right mandible fragment with I-M₂

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Rem85, p. 32, pl. 7, figs. 6a-b

Sorex bor REUMER, 1985

paratypes, V 2008.8.1.

four mandible fragments

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Rem85, p. 32

Sorex bor REUMER, 1985

paratype, V 2008.8.2.

maxilla fragment

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Rem85, p. 32

Sorex bor REUMER, 1985

paratypes, V 2008.8.3.

one incisor and three upper molars

Middle Pliocene

Osztramos, locality 1, Tornaszentandrás

Rem85, p. 32

- Sorex bor* REUMER, 1985
 paratypes, V 2008.9.1.
 12 mandible fragments
 Late Pliocene
 Osztramos, locality 7, Tornaszentandrás
Rem85, p. 32, pl. 7, figs. 5a–b, 7a–b, 8a–c
- Sorex bor* REUMER, 1985
 paratypes, V 2008.9.2.
 one upper incisor and two molars
 Late Pliocene
 Osztramos, locality 7, Tornaszentandrás
Rem85, p. 32
- Sorex bor* REUMER, 1985
 paratypes, V 2008.10.1.
 three mandible fragments
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
Rem85, p. 32
- Sorex bor* REUMER, 1985
 paratypes, V 2008.10.2.
 one upper incisor and one upper molar
 Middle Pliocene
 Osztramos, locality 9, Tornaszentandrás
Rem85, p. 32, pl. 7, fig. 2
- Sorex gracilis* PETÉNYI in KUBINYI, 1856
 syntypes
 right mandible fragment with one molar
 and left mandible fragment without teeth
 Middle Pliocene
 quarry, Beremend
Kub56, p. 76
- Sorex minutus*, *Rem85*
 Types figured and described in detail in
 PETÉNYI 1864: p. 70, pl. 1, figs. 6a–e.
 Types missing, not found in inventory
- Sorex margaritodon* KORMOS, 1930
 holotype, V.61.1589
 skull fragment with full dentition
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
Krm30, p. 240
- Drepanosorex margaritodon*, Kre56
- Sorex margaritodon* KORMOS, 1930
 paratypes, V.61.1589
 three mandible fragments
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
Krm30, p. 240
- Drepanosorex margaritodon*, Kre56
- Sorex savini austriacus* KORMOS, 1937
 lectotype, V.61.1590
 right mandible
 Middle Pleistocene
 Hundsheim, Austria
Krm37, p. 31, text-fig. 2
- Drepanosorex austriacus*, Rab72
 Lectotype designated in RABEDER 1972
- Sorex tasnádii* KRETZOI, 1941
 holotype, V.59.1045
 right mandible fragment
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
Kre41a, p. 109, text-fig. 1a
 Generic assignment in original description
 given as *Sorex* or *Drepanosorex*. Type missing
- Soriculus kubinyii* KORMOS, 1934
 paratype, V.61.1554
 left mandible
 Middle Pleistocene
 Templom-hegy, Villány
Krm34a, p. 303 (text-fig. 36)
Asoriculus gibberodon, Kre62; Més99
- Talpa vulgaris fossilis* PETÉNYI in KUBINYI, 1856
 syntypes, V 2008.34.1.
 two mandible fragments
 Middle Pliocene
 quarry, Beremend
Kub56, p. 73
- Talpa fossilis*, Kre38
 Types figured and described in detail in
 PETÉNYI 1864: p. 53, pl. 1, figs. 4a–e, 4g–l.
 Four additional mandibles, one sternum,
 eight ulnae, two radii and one tibia reported
 in PETÉNYI 1864; those not found in inventory
 and probably lost

- Talpa vulgaris fossilis* PETÉNYI in KUBINYI, 1856
 syntypes, V 2008.34.2.
 ten scapulae
 Middle Pliocene
 quarry, Beremend
 Kub56, p. 73
Talpa fossilis, Kre38
- Talpa vulgaris fossilis* PETÉNYI in KUBINYI, 1856
 syntypes, V 2008.34.3.
 15 humeri
 Middle Pliocene
 quarry, Beremend
 Kub56, p. 73
Talpa fossilis, Kre38
 Three additional humeri reported in PETÉNYI 1864
- Talpa vulgaris fossilis* PETÉNYI in KUBINYI, 1856
 syntypes, V 2008.34.4.
 six femurs
 Middle Pliocene
 quarry, Beremend
 Kub56, p. 73
Talpa fossilis, Kre38
 One additional femur reported in PETÉNYI 1864
- Trilophodon angustidens praetypica* TASNÁDI
 KUBACSKA, 1939
 syntype, 2007.96.1.
 maxilla fragment with left and right I²
 Ottnangian, Miocene; Salgótarján Formation
 coal mine, Zagyvapálfalva, Salgótarján
 Tas39, p. 154, text-fig. 1
Gomphotherium angustidens, Gas01
- Trilophodon angustidens praetypica* TASNÁDI
 KUBACSKA, 1939
 syntype, 2007.96.2.
 left mandible with I₂, P₄-M₂
 Ottnangian, Miocene; Salgótarján Formation
 coal mine, Zagyvapálfalva, Salgótarján
 Tas39, p. 154, pl. 4
Gomphotherium angustidens, Gas01
- Trilophodon angustidens praetypica* TASNÁDI
 KUBACSKA, 1939
 syntype, 2007.96.3.
 right mandible with I₂, P₄-M₂
 Ottnangian, Miocene; Salgótarján Formation
 coal mine, Zagyvapálfalva, Salgótarján
 Tas39, p. 154, pl. 4
Gomphotherium angustidens, Gas01
- Ungaromys nanus* KORMOS, 1932
 holotype, V.61.1528
 right M₁-M₃ toothrow
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 336, text-fig. 6 (text-fig. 7)
- Ungaromys nanus* KORMOS, 1932
 paratype, V.61.1528
 right M₂
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 336, text-fig. 8
- Ungaromys nanus* KORMOS, 1932
 paratype, V.61.1528
 incisor
 Early Pleistocene
 Somlyó-hegy, Püspökkürdő, Oradea
 (Nagyvárad), Romania
 Krm32a, p. 336
- Ursus etruscus gombaszögensis* KRETZOL, 1938
 holotype
 right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 21. Type not identified with certainty.
 Well-preserved, unfigured M₂ under inventory
 number V.59.1048 suspect as possible holotype
 but cannot be proven as no original inventory
 number marked on specimen. Type label found
 erroneously placed under figured paratype
 V.59.930
- Ursus etruscus gombaszögensis* KRETZOL, 1938
 paratype, V.59.930
 right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, figs. 15-16
 Inventory number in original publication:
 V. 883

- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.932
 right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 85
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.935
 right P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, figs. 9–10
 Inventory number in original publication:
 Fa. 33
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratypes, V.59.955
 canine fragments
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 798
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.961
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 872
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.964
 right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 987
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.964
 fragment of right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 92
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.969
 posterior half of left M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, fig. 12
 Inventory number in original publication:
 V. 806
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.972
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 920
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.972
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 39
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.985
 left P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 88

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1005
 right M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 W. 42

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1008
 right M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication: W. 3

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1009
 right M¹
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 90

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1010
 left M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, fig. 18
 Inventory number in original publication:
 V. 925

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1010
 left M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication: W. 4

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1010
 left M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 87

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1019
 left M¹
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 999

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1019
 left M¹
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 89

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1019
 left M¹
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, figs. 23–24
 Inventory number in original publication:
 V. 887

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1023
 right M₁ without talonid
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 W. 50

- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1023
 right M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 83
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1023
 anterior half of right M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, fig. 14
 Inventory number in original publication:
 V. 791
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1023
 anterior half of right M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 82
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1023
 posterior half of right M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, fig. 13
 Inventory number in original publication:
 V. 937
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1023
 posterior half of right M₁
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 81
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1043
 right mandible fragment with M₂–M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 37
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1048
 right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 38
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1048
 anterior fragment of right M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 86
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1049
 proximal fragment of penis bone
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, fig. 26
 Inventory number in original publication:
 Fa. 36
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, V.59.1061
 left P⁴
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög),
 Slovakia
 Kre38, p. 138, pl. 3, figs. 19–20
 Inventory number in original publication:
 Fa. 35

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1065
 left M²
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, fig. 25
 Inventory number in original publication:
 V. 926

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1065
 left M²
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 931

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1065
 left M²
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 93

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratypes, V.59.1080
 four lower incisors
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 W. 47

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratypes, V.59.1081
 four I³
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 893

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1089
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication: W. 5

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1099
 right M²
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 841

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1106
 right P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, figs. 7–8
 Inventory number in original publication:
 V. 955

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, V.59.1106
 posterior fragment of right P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, fig. 11
 Inventory number in original publication:
 Fa. 34

Ursus etruscus gombaszögensis KRETZOI, 1938
 paratype, 2007.105.1.
 right P₄
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, figs. 21–22
 Inventory number in original publication: W. 8

- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, 2007.106.1.
 right M₃
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138, pl. 3, fig. 17
 Inventory number in original publication:
 V. 975
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, 2007.107.1.
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 V. 996
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, 2007.108.1.
 right M²
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 91
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype, 2007.109.1.
 lower canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 95. Another lower canine under same original inventory number missing
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype
 upper canine
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 94. Type missing, not found in current inventory
- Ursus etruscus gombaszögensis* KRETZOI, 1938
 paratype
 left M₂
 Early Pleistocene
 fissure fill in quarry, Gombasek (Gombaszög), Slovakia
 Kre38, p. 138
 Inventory number in original publication:
 Fa. 84. Type missing, not found in current inventory
- Vespertilio majori* KORMOS, 1934
 holotype, V.61.1413
 left mandible
 Middle Pleistocene
 Templom-hegy, Villány
 Krm34a, p. 312, text-fig. 43
 Inventory number in original publication:
 Nr. 3847
- Vormela petenyii* KRETZOI, 1942
 holotype, V.61.1386
 right mandible fragment with P₄-M₁
 Middle Pleistocene
 Templom-hegy, Villány
 Kre42c, p. 331 (244)
 Inventory number in original publication:
 Nr. 3915. Type figured in KORMOS 1934b: pl. 2, fig. 5a
- Xenocyon lycaonoides* KRETZOI, 1938
 holotype, V.64.911
 right M¹
 Early Pleistocene
 fissure fill in quarry, locality 3, Gombasek (Gombaszög), Slovakia
 Kre38, p. 132, pl. 3, fig. 4
 Inventory number in original publication:
 Fa. 20

Index of species and subspecies names

abnormis	<i>Kahlerosphaera abnormis</i> KOZUR, MOIX & OZSVÁRT 22
abundantus	<i>Balanus amphitrite abundantus</i> KOLOSVÁRY 111
ackneri	<i>Lamna ackneri</i> NEUGEBOREN 124
acrorthiza	<i>Clethrionomys acrorthiza</i> KORMOS 139
acuminata	<i>Nodosaria acuminata</i> HANTKEN 69
acuta	<i>Cominella hungarica acuta</i> NOSZKY 83
acutareatus	<i>Hemithiris acuta</i> MEZNERICS 111
acuticaudata	<i>Sulcogladius collegnii acuta</i> NOSZKY 90
acutidorsatum	<i>Rhaetomegalodon acutareatus</i> VÉGH-NEUBRANDT 29
aqualis	<i>Lithoconus ineditus acuticaudata</i> NOSZKY 86
afer	<i>Haplophragmium acutidorsatum</i> HANTKEN 68
africana	<i>Trochosmilia aequalis</i> REUSS 49
agriense	<i>Turbo afer</i> COQUAND 34
agriensis	<i>Spiriferina africana</i> COQUAND 37
albus	<i>Bitium spina agriense</i> BÁLDI 81
alexandri	<i>Chlamys agriensis</i> CSEPREGHY-MEZNERICS 73
alternans	<i>Raphitoma roemerri agriensis</i> BÁLDI 89
altialata	<i>Sepia agriensis</i> WAGNER 60
altispirata	<i>Pagurus albus</i> MÜLLER 115
altus	<i>Turritella (Haustator) strangulata alexandri</i> NOSZKY 92
alveata	<i>Euthriofusus szontaghii alternans</i> NOSZKY 84
ambiguus	<i>Lucina (Megaxinus) elliptica altialata</i> NOSZKY 76
ammoniticeras	<i>Volutilithes (Neoathleta) obliqua altispirata</i> NOSZKY 94
ammonoides	<i>Caprocancer altus</i> MÜLLER & COLLINS 62
amplus	<i>Lamna (Odontaspis) alveata</i> NEUGEBOREN 124
ancestralis	<i>Otodus ambiguus</i> NEUGEBOREN 126
andreasii	<i>Heteramonites ammoniticeras</i> COQUAND 39
angusta	<i>Adeorbis ammonoides</i> CSEPREGHY-MEZNERICS 104
angustiannulatus	<i>Erycites telediroti amplius</i> GÉCZY 35
angustiformis	<i>Angulus (Peroniida) planatus ancestralis</i> BÁLDI 71
anomalidens	<i>Echinophoria intermedia andreasii</i> NOSZKY 83
anomaloides	<i>Lima cancellata angusta</i> NOSZKY 76
antalas	<i>Thyasira vara angusta</i> BÁLDI 81
antiquus	<i>Archaeoacanthocircus angustiannulatus</i> KOZUR, MOIX & OZSVÁRT 21
aperta	<i>Flabellipecten burdigalensis angustiformis</i> BÁLDI 74
applanata	<i>Rhinolophus macrorhinus anomalidens</i> TOPÁL 157
archiinexpectatus	<i>Nummulites anomalooides</i> KECSKEMÉTI 45
arcuatocaudata	<i>Terebratula (Waldheimia) antalas</i> COQUAND 37
arcuatodecrescens	<i>Glis antiquus</i> KORMOS 143
arcuatospirata	<i>Huglusphaera aperta</i> KOZUR, MOIX & OZSVÁRT 21
arcuatostriata	<i>Globigerina applanata</i> HANTKEN 42
arcuatus	<i>Balanus amphitrite archi-inexpectatus</i> KOLOSVÁRY 111
	<i>Ficula tenuis arcuato-caudata</i> NOSZKY 84
	<i>Otodus arcuato-decrescens</i> NEUGEBOREN 126
	<i>Surcula regularis arcuatospirata</i> NOSZKY 91
	<i>Cristellaria (Robulina) arcuatostrriata</i> HANTKEN 67
	<i>Megalodus arcuatus</i> VÉGH-NEUBRANDT 28
	<i>Polyonyx arcuatus</i> MÜLLER & COLLINS 65

arnaudi	<i>Cyclas arnaudi</i> COQUAND 32
arvicolinus	<i>Siphneus arvicolinus</i> NEHRING 162
asperum	<i>Nodophthalmidium asperum</i> GÖRÖG 97
aspinosa	<i>Huglusphaera aspinosa</i> KOZUR, MOIX & OZSVÁRT 21
aspirata	<i>Acera aspirata</i> SZÓTS 51
asteriscus	<i>Favosites asteriscus</i> FRECH 18
asymmetrica	<i>Mioranina asymmetrica</i> MÜLLER 115
atavus	<i>Hystrix vinogradovi atavus</i> JÁNOSSY 143
augustini	<i>Aporrhais augustini</i> COQUAND 38
auriculata	<i>Bythinella auriculata</i> SZÓTS 52
austriaca	<i>Mitra goniophora austriaca</i> CSEPREGHY-MEZNERICCS 107
austriacus	<i>Sorex savini austriacus</i> KORMOS 163
bachmayeri	"Xaiva" <i>bachmayeri</i> MÜLLER 120
bacilloides	<i>Nodosaria bacilloides</i> HANTKEN 69
baconica	<i>Harpactocarcinus telegi-rothi baconica</i> TOMOR-THIRRING 64
baconicum	<i>Thecocathus baconica</i> KOLOSVÁRY 32
bactchisaraiensis	<i>Dicerocardium baconicum</i> VÉGH-NEUBRANDT 28
badenis	<i>Parallelodon baconicum</i> VÉGH 29
badensis	<i>Nummulina bactchisaraiensis</i> ROZLOZSNIK 42
bajotensis	<i>Metopograpsus badenis</i> MÜLLER 115
bakonycsernyensis	<i>Clavatula (Surcula) consobrina badensis</i> CSEPREGHY-MEZNERICCS 104
bakonyensis	<i>Roualtia lapugensis badensis</i> CSEPREGHY-MEZNERICCS 109
balatonica	<i>Rhaetomegalodon bajotensis</i> VÉGH-NEUBRANDT 29
balaticum	<i>Astrorhiza bakonycsernyensis</i> OZSVÁRT 41
baloghi	<i>Ceratotrochus bakonyensis</i> KOLOSVÁRY 37
baranensis	<i>Triadogigantocypris balatonica</i> MONOSTORI 31
basilaevi	<i>Encoiloceras balaticum</i> VÖRÖS 30
beaumonti	<i>Plerophyllum (Ufimia) baloghi</i> KOLOSVÁRY 18
bellardii	<i>Myotis baranensis</i> KORMOS 150
beremendensis	<i>Delphinula scobia basilaevi</i> NOSZKY 83
berenderi	<i>Aptychus beaumontii</i> COQUAND 34
biaense	<i>Murex (Pteronotus) detritus bellardii</i> NOSZKY 87
bicarinata	<i>Cricetinus beremendensis</i> HÍR 139
bicarinatus	<i>Mustela beremendensis</i> PETÉNYI in KUBINYI 149
binominata	<i>Otis khosatzkii beremendensis</i> JÁNOSSY 134
bipartita	<i>Pliolagus beremendensis</i> KORMOS 154
bistriata	<i>Turritella (Mesalia) berenderi</i> SZÓTS 59
bisulcata	<i>Chlamys biaense</i> CSEPREGHY-MEZNERICCS 100
blainvillei	<i>Melanopsis doboi bicarinata</i> SCHRETER 121
bockhi	<i>Pliolagus beremendensis</i> KORMOS 154
bodvanus	<i>Rostellaria bicarinata</i> BÁLDI 89
bogschi	<i>Anisus bicarinatus</i> SZÓTS 51
bogschi	<i>Nodosaria binominata</i> FRANZENAU 97
bipartita	<i>Chlamys palmata bipartita</i> CSEPREGHY-MEZNERICCS 100
bistriata	<i>Volutilithes (Athleta) ficalina bistriata</i> NOSZKY 93
bisulcata	<i>Turritella (Haustator) strangulata bisulcata</i> NOSZKY 92
blainvillei	<i>Aptychus blainvillei</i> COQUAND 39
bockhi	<i>Caranx böckhi</i> GORJANOVIĆ-KRAMBERGER 123
bodvanus	<i>Nummulina böckhi</i> ROZLOZSNIK 42
bogschi	<i>Vexillum böckhi</i> SZÓTS 59
bogschi	<i>Leptodontomys bodvanus</i> JÁNOSSY 145
bogschi	<i>Cardium scobinula bogschi</i> NOSZKY 72
bogschi	<i>Mangelia (Enatoma) bogschi</i> BÁLDI 86
bogschi	<i>Natica bogschi</i> SZÓTS 56

- bolcskeyensis *Antedon bolcskeyensis* SZALAI 120
 boldii *Tonohamites boldii* SZIVES & MONKS 40
 bolkayi *Dolomys episcopalis bolkayi* KORMOS 141
 bombitus *Nummulites bombitus* HOTTINGER 45
 bor *Sorex bor* REUMER 162
 borealis *Turdoidea borealis* JÁNOSSY 135
 boreaui *Melania boreaui* COQUAND 33
Physa boreaui COQUAND 33
 borsodensis *Arca borsodensis* CSEPREGHY-MEZNERICS 99
Chama gryphoides borsodensis CSEPREGHY-MEZNERICS 100
Clavatula (Surcula) nodosa borsodensis CSEPREGHY-MEZNERICS 105
Episoriculus borsodensis JÁNOSSY 142
Miltha (Eomittha) borsodensis CSEPREGHY-MEZNERICS 102
Nassa (?Tritia) borsodensis CSEPREGHY-MEZNERICS 107
Sigaretus borsodensis CSEPREGHY-MEZNERICS 110
Bittium botense CSEPREGHY-MEZNERICS 104
Genota (Pseudotoma) bonelli botensis CSEPREGHY-MEZNERICS 106
Bayania boussaci SZÓTS 51
Nummulina boussaci ROZLOZNIK 42
Peccholia argentea brachiumbonata NOSZKY 78
Trochosmilia brachypoda REUSS 49
Xiphothecaella brevicaudata KOZUR, MOIX & OZSVÁRT 26
Callianassa brevimanus BEURLEN 95
Alces brevirostris KRETZOI in JÁNOSSY 135
Arca (Acar) dactylus brevis NOSZKY 71
Sulcogladius collegnii brevis NOSZKY 90
Zhamojdasphaera rigoi brevispinosa KOZUR, MOIX & MOSTLER 27
Conus (Leptoconus) dujardini brevispiratus NOSZKY 83
Rhabdophyllia budense KOLOSVÁRY 48
Cultellus budensis BALDI 73
Flabellina budensis HANTKEN 68
Marginulina budensis HANTKEN 69
Nodosaria budensis HANTKEN 69
Paraxanthosia budensis MÜLLER & COLLINS 65
Ranella (Apollon) gigantea budensis NOSZKY 89
Robulina budensis HANTKEN 70
Tauraxinus budensis NOSZKY 80
 bukkense *Astarte solidula bukkense* CSEPREGHY-MEZNERICS 99
Pyrene (Anachis) guembeli bukkense CSEPREGHY-MEZNERICS 109
Chama bukkensis CSEPREGHY-MEZNERICS 99
Lonsdaleoides bükkense KOLOSVÁRY 18
Surcula regularis bulbiformis NOSZKY 91
Surcula bulbosa NOSZKY 90
Allocricetus bursae SCHAUB 135
Arca caillatiformis SZÓTS 49
Rostellaria hungarica calcarata NOSZKY 90
Ampullina (Globularia) telegdirothi callosa NOSZKY 81
Globularia gibberosa callosa BÁLDI 85
Chenopus callosus TELELDI ROTH 82
Corbula carentonensis COQUAND 32
Mosasaurus carentonensis COQUAND 133
Leuciscus carpathicus BÖHM 126
Rakosia carupoides MÜLLER 118

cavidens	<i>Lamna cavidens</i> NEUGEBOREN 125
chamaeformis	<i>Venus chamaeformis</i> CSEPREGHY-MEZNERICS 103
chinensis	<i>Haplothecia? chinensis</i> FRECH 18
	<i>Lima chinensis</i> LÓCZY 28
cholnokyi	<i>Spirillina chinensis</i> LÖRENTHEY 17
cingulata	<i>Bythinia (?) cholnokyi</i> SCHLOSSER 121
cladangiae	<i>Pseudotoma orbignyi cingulata</i> NOSZKY 88
clarae	<i>Creusia spinulosa cladangiae</i> KOLOSVÁRY 112
clatratus	<i>Galeodea echinophora clarae</i> NOSZKY 85
clava	<i>Acamptochetus clatratus</i> BÁLDI 81
claviformis	<i>Lagena clava</i> OZSVÁRT 42
collaris	<i>Hypoxiphothecaella claviformis</i> KOZUR, MOIX & OZSVÁRT 22
collectiva	<i>Podobursa claviformis</i> KOZUR, MOIX & OZSVÁRT 24
collispannoni	<i>Dolichotoma collaris</i> NOSZKY 83
colorata	<i>Lyria collaris</i> NOSZKY 86
commenticius	<i>Drillia collectiva</i> NOSZKY 83
compressa	<i>Procapra (Protetraceros) collispannoni</i> KRETZOI 155
concavus	<i>Tricolia colorata</i> SZÓTS 59
concentrica	<i>Mediocris commenticius</i> KAZÁR 148
condamyi	<i>Siliqua harmati compressa</i> NOSZKY 80
confluentiseptatus	<i>Pagurus concavus</i> MÜLLER 115
conica	<i>Discocyclina concentrica</i> KECSKEMÉTI 41
conspicuus	<i>Corbula condamyi</i> COQUAND 32
constricta	<i>Phineus confluentiseptatus</i> KOLOSVÁRY 18
contorta	<i>Adeorbisina conica</i> SZABÓ 33
convexa	<i>Turbanilla conica</i> SZÓTS 59
coronatum	<i>Fusus conspicuus</i> COQUAND 38
coronensis	<i>Spinoprotunuma? constricta</i> KOZUR, MOIX & OZSVÁRT 25
costatus	<i>Nodosaria (Dentalina) contorta</i> HANTKEN 69
costulatum	<i>Malletia caterini convexa</i> NOSZKY 77
crassa	<i>Ruesticyrtium coronatum</i> KOZUR, MOIX & OZSVÁRT 24
crenellaeformis	<i>Microtus coronensis</i> KORMOS 148
cretaceum	<i>Epistomarooides costatus</i> OZSVÁRT 41
csakvarensse	<i>Amussium costulatum</i> NOSZKY 71
csakvarensis	<i>Fusus pergracilis crassa</i> NOSZKY 85
csepreghymeznericsae	<i>Spondylus crenellaeformis</i> NOSZKY 80
cserhatensis	<i>Buccinum cretaceum</i> COQUAND 38
csobankanus	<i>Vexillum csákvárense</i> SZÓTS 59
curtimanus	<i>Cyclostrema csákvárensis</i> SZÓTS 53
curvicostata	<i>Chlamys (Aequipecten) csepreghy-meznericsae</i> BÁLDI 73
cylindrica	<i>Beguina (Carditamera) striatellata</i> cserhátensis CSEPREGHY-MEZNERICS 99
dacicus	<i>Paguristes cserhátensis</i> MÜLLER 115
dacus	<i>Paphia waldmanni</i> cserhátensis Cs. MEZNERICS 103
dakotana	<i>Hypsolophiodon csobánkanus</i> KRETZOI 143
dalmatinus	<i>Dardanus curtimanus</i> MÜLLER & COLLINS 62
darchiaci	<i>Calamophyllia curvicostata</i> KOLOSVÁRY 48
	<i>Zebinella decussata curvicostata</i> NOSZKY 94
	<i>Clavulina cylindrica</i> HANTKEN 41
	<i>Muscardinus dacicus</i> KORMOS 149
	<i>Titanosaurus dacus</i> NOPCSA 134
	<i>Cadurcopsis dakotana</i> KRETZOI 139
	<i>Dolomys dalmatinus</i> KORMOS 141
	<i>Nummulina bronniarti d'archiaci</i> ROZLOZNIK ex HANTKEN & MADARÁSZ 43

- darnoensis
declinatus
dejtarensse
dendricola
densespiratum
densicosta
- densicostatus
densiradiata
densitexta
dentatus
denticulata
denticulatus
depressa
- depressosuturata
desgodinsi
dicampyla
digitata
diodesensis
divaricata
doboi
- dregeri
dubia
- dudarensis
ebeczkyi
edentata
edentula
eduardi
effusa
- egerensis
ehiki
- elegans
- ellipticus
elongata
- elongatula
- Chlamys darnóensis* CSEPREGHY-MEZNERICSS 100
Mesolambrus declinatus MÜLLER & COLLINS 64
Dentalium densitextum dejtarensse BÁLDI 81
Stephanosmilia dendricola KOLOSVÁRY 48
Bittium reticulatum densespiratum BÁLDI 81
Pleurotomaria sismondaei densicosta NOSZKY 88
Spondylus tenuispina densicosta NOSZKY 80
Fusus (Aptyxis) lóczyi densicostatus NOSZKY 84
Actinopteria? densiradiata LÓCZY 19
Dentalium haeringense densitexta NOSZKY 81
Theodoxus (Calvertia?) grateloupianus dentatus KÓKAY 110
Longoporellana denticulata MÜLLER & COLLINS 64
Budapanopeus denticulatus MÜLLER & COLLINS 61
Lamna depressa NEUGEBOREN 125
Lucina (Megaxinus) bellardiana depressa NOSZKY 76
Melanella naumanni depressosuturata BÁLDI 87
Productus (Marginifera) desgodinsi LÓCZY 20
Cristellaria dicampyla FRANZENAU 96
Chenopus speciosus digitata TELEGI ROTH 82
Crossotonotus diodesensis MÜLLER 114
Triloculina divaricata FRANZENAU 98
Chrysodomus dobói NOSZKY 82
Melanopsis doboi SCHRÉTER 121
Lucina (Myrthea) elongata dregeri NOSZKY 76
Chonetella dubia LÓCZY 20
Gibbula dubia BÁLDI 85
Nummulites dudarensis KECSKEMÉTI 46
Castor eberezkyi KRENNER 139
Hinia fortecostata edentata BÁLDI 86
Quinqueloculina peregrina edentula FRANZENAU 98
Karadagithyris eduardi VÖRÖS 36
Ficula tenuis effusa NOSZKY 84
Surcula effusa NOSZKY 90
Volutilithes (Neoathleta) affinis effusa NOSZKY 93
Tinostoma egerensis BÁLDI 91
Allocricetus éhiki SCHAUB 136
Chenopus uttingerianus éhiki NOSZKY 82
Carcharodon elegans NEUGEBOREN 123
Plecantium elegans HANTKEN 70
Xiphothecaella elegans KOZUR, MOIX & OZSVÁRT 26
Prochlorodius ellipticus MÜLLER & COLLINS 65
Avicula hirundo elongata NOSZKY 72
Cominella hungarica elongata NOSZKY 83
Drillia obtusa elongata NOSZKY 83
Hypoxiphothecaella elongata KOZUR, MOIX & OZSVÁRT 22
Lamna (Odontaspis) elongata NEUGEBOREN 125
Limopsis retifera elongata NOSZKY 76
Lucina (Myrthea) elongata NOSZKY 76
Nassa (Telasco) neuvillei elongata NOSZKY 87
Sulcoglaucus collegioides elongata NOSZKY 90
Syndesmia elongata NOSZKY 80
Textularia elongata HANTKEN 70
Oliva (Olivella) inflata elongatula NOSZKY 88

elongatus	<i>Ovocarcinus elongatus</i> MÜLLER & COLLINS 64
eotvoesi	<i>Dromilites eotvoesi</i> MÜLLER 114
epianthus	<i>Clypeaster epianthus</i> MEZNERICS 120
erecta	<i>Arca (Acar) dactylus erecta</i> NOSZKY 71
	<i>Lucina (Megaxinus) elliptica erecta</i> NOSZKY 76
	<i>Pleurotoma coronata erecta</i> NOSZKY 88
estramontis	<i>Rhinolophus estramontis</i> TOPÁL 156
estramosensis	<i>Myotis estramosensis</i> TOPÁL 150
estramosi	<i>Porzana estramosi</i> JÁNOSSY 135
europaea	<i>Blarinella europaea</i> REUMER 138
euthiraeformis	<i>Triton euthiraeformis</i> NOSZKY 91
evae	<i>Ethusa evae</i> MÜLLER & COLLINS 63
evanicsi	<i>Scaphites (Scaphites) evanicsi</i> SZIVES 40
exalata	<i>Gryphaea bronniarti exalata</i> NOSZKY 75
excellens	<i>Modiolus excellens</i> CSEPREGHY-MEZNERICS 103
exogyroides	<i>Ostrea lamellosa exogyroides</i> CSEPREGHY-MEZNERICS 103
extinctorius	<i>Helcion extinctorius</i> COQUAND 33
faludyi	<i>Kahlerosphaera faludyi</i> KOZUR, MOIX & OZSVÁRT 22
februarius	<i>Brachynotus februarius</i> MÜLLER 112
fejervaryi	<i>Mimomys fejérváryi</i> KORMOS 148
	<i>Testudo fejérváryi</i> SZALAI 133
felmenesensis	<i>Prionechinus felmenensis</i> LAMBERT & THIÉRY 120
fissidens	<i>Crossopus fissidens</i> PETÉNYI in KUBINYI 140
foezyi	<i>Hypoxiphothecaella foezyi</i> KOZUR, MOIX & OZSVÁRT 22
foldvari	<i>Lucina (Dentilucina) meneghinii földvárii</i> NOSZKY 77
formosa	<i>Pentaglis földvárii</i> KRETZOI 153
fortestriata	<i>Frondicularia formosa</i> FRANZENAU 96
fossata	<i>Euthria subnodosata fortestriata</i> CSEPREGHY-MEZNERICS 106
fossatus	<i>Dromilites fossata</i> MÜLLER & COLLINS 62
fossilis	<i>Gemmacarcinus fossatus</i> MÜLLER & COLLINS 63
	<i>Balanus improvisus fossilis</i> KOLOSVÁRY 111
	<i>Lamna (Odontaspis) ferox fossilis</i> NEUGEBOREN 125
	<i>Talpa vulgaris fossilis</i> PETÉNYI in KUBINYI 163
fotensis	<i>Pecten fótensis</i> CSEPREGHY-MEZNERICS 103
foveolata	<i>Vertebralina foveolata</i> FRANZENAU 98
fragilis	<i>Thalamita fragilis</i> MÜLLER 119
franciae	<i>Ovamene franciae</i> MÜLLER & COLLINS 64
franciscae	<i>Daphnella franciscae</i> NOSZKY 83
franzenau	<i>Hopkinsina franzenau</i> MAJZON 97
	<i>Nassa (Hima) franzenau</i> CSEPREGHY-MEZNERICS 108
	<i>Semipecten Mayeri</i> franzenau NOSZKY 79
frentanaeformis	<i>Turborotalia munda</i> franzenau SZTRÁKOS 70
frequens	<i>Nummulina granifera frentanaeformis</i> ROZLOZNIK 43
	<i>Marginella frequens</i> SZÖTS 54
	<i>Mathilda frequens</i> SZÖTS 56
furcatolateratum	<i>Dicerocardium furcatolateratum</i> VÉGH-NEUBRANDT 28
furcastostriata	<i>Spinoprotunuma? furcastostriata</i> KOZUR, MOIX & OZSVÁRT 25
fusiformis	<i>Podobursa fusiformis</i> KOZUR, MOIX & OZSVÁRT 24
gaali	<i>Agriarctos gaali</i> KRETZOI 135
	<i>Saxicava gaali</i> NOSZKY 79
gaboraroni	<i>Pisanianura gaboraroni</i> CSEPREGHY-MEZNERICS 109
galaczi	<i>Bakonydraco galaczi</i> ÖSI, WEISHAMPEL & JIANU 127
	<i>Dichotomosella galaczi</i> VÖRÖS 36

- galaczi *Limea (Pseudolimea) galaczi* SZENTE 33
 gallicus *Priabonocarcinus gallicus* MÜLLER & COLLINS 65
 gantensis *Cylichna gántensis* SZÖTS 53
Murex gántensis SZÖTS 56
Solariella gántensis SZÖTS 58
Trinacria gántensis SZÖTS 50
Voluta (Lyria) gárdonyii NOSZKY 93
Globuligerina geczyi GÖRÖG 31
Nerinea gemmifera COQUAND 39
Ruesticyrtium georgi georgi KOZUR, MOIX & OZSVÁRT 24
Natica gervaisi COQUAND 39
Dentalina gigantea HANTKEN 41
Muscardinus giganteus JÁNOSSY 149
Lingulina glabra HANTKEN 42
Syringocapsa glabra KOZUR, MOIX & OZSVÁRT 26
Trapezia glaessneri MÜLLER 119
Globigerina globosa HANTKEN 42
Ebalia globulosa MÜLLER 114
Chonetes flemmingi gobica LÓCZY 20
Helicigona goemoerensis SOÓS 121
Leo gombaszögensis KRETZOI 144
Ursus etruscus gombaszögensis KRETZOI 164
Spinomersinella goricanae KOZUR, MOIX & OZSVÁRT 25
Sinophyllum gracile KOLOSVÁRY 19
Leda (Lembulus) gracilis NOSZKY 75
Pleurotoma rotata gracilis NOSZKY 88
Rostellaria hungarica gracilis NOSZKY 90
Sorex gracilis PETÉNYI in KUBINYI 163
Surcula beyrichi gracilis NOSZKY 90
Bythinella gracillima SZÖTS 52
Volutilithes (Neothleteta) obliqua gracilis NOSZKY 94
Tectura tauroconica gradiformis NOSZKY 91
Assiminea gránásensis SZÖTS 51
Natica gránásensis SZÖTS 56
Mastodon (Bunolophodon) grandincisivum SCHLESINGER 148
Asthenotoma graniformis SZÖTS 51
Robulina granulata HANTKEN 47
Turbo (Senectus) ranellaformis granulata NOSZKY 92
Panopeus granulatus MÜLLER & COLLINS 65
Micromithrax? grippi MÜLLER 115
Macularia lartetii grundensis CSEPREGHY-MEZNERICS 107
Pyrene (Anachis) bellardii grussbachensis CSEPREGHY-MEZNERICS 109
Dentalina gümbeli HANTKEN 68
Rhynchonella guerrizila COQUAND 36
Trochus guilhoti COQUAND 33
Rhaetomegalodon incisus gutnici VÉGH-NEUBRANDT 29
Eulimella guttulina SZÖTS 54
Perumys gyulavarii KRETZOI & VÖRÖS 153
Lytoceras haasi GÉCZY 36
Vaginulinopsis hagni OZSVÁRT 47
Kahlerosphaera hamvasi KOZUR, MOIX & OZSVÁRT 23
Balanus hantkeni KOLOSVÁRY 61
Cyathophyllia hantkeni REUSS 48

hantkeni	<i>Cylichna hantkeni</i> SZŐTS 54 <i>Glandulina hantkeni</i> FRANZENAU 96 <i>Lima (Acesta) miocenica</i> hantkeni NOSZKY 76 <i>Marginella hantkeni</i> SZŐTS 55 <i>Nerita hantkeni</i> SZŐTS 56 <i>Tritaxilina hantkeni</i> CUSHMAN 70
harmati	<i>Aequipecten deletus</i> harmati NOSZKY 70 <i>Belone harmati</i> WEILER 123 <i>Malletia harmati</i> BÁLDI 77 <i>Sepia harmati</i> SZÖRÉNYI 94 <i>Siliqua harmati</i> NOSZKY 80 <i>Vexillum (Costellaria) harmati</i> CSEPREGHY-MEZNERICS 110
haueri	<i>Lamna haueri</i> NEUGEBOREN 125 <i>Oxyrhina haueri</i> NEUGEBOREN 126
haydnii	<i>Petrolisthes haydnii</i> MÜLLER 117
heberti	<i>Nerita heberti</i> SZŐTS 56
heckeliana	<i>Oxyrhina heckeliana</i> NEUGEBOREN 126
hegeduesi	<i>Calliostoma (Ampullotrochus) elegantulum</i> hegeduesi BÁLDI 81
helbingi	<i>Baranogale helbingi</i> KORMOS 138
helenae	<i>Balanus amphitrite helenae</i> KOLOSVÁRY 111 <i>Galeodes taurinensis helenae</i> NOSZKY 85
helleri	<i>Prosmilia helenae</i> KOLOSVÁRY 19
hevesensis	<i>Eliomys querinus helleri</i> JÁNOSSY 142 <i>Aequipecten opercularis hevesensis</i> SCHRÉTER 98
hidasesensis	<i>Phos hevesensis</i> BÁLDI 88
hidasensis	<i>Arca (Anadara) hidasesensis</i> STRAUSZ 99 <i>Myxas hidasesensis</i> CSEPREGHY-MEZNERICS 107 <i>Rissoa hidasesensis</i> CSEPREGHY-MEZNERICS 109
hintoni	<i>Valvata hidasesensis</i> KÓKAY 110
hintonianus	<i>Pitymys hintoni</i> KRETZOI 154
hodnaensis	<i>Clethrionomys hintonianus</i> KRETZOI 139
hodnensis	<i>Isastraea hodnaensis</i> COQUAND 31
hofmanni	<i>Lima hodnensis</i> COQUAND 33
holitschensis	<i>Arcopsis (Arcopsis) hofmanni</i> BÁLDI 71
holocristatus	<i>Lima (Acesta) miocenica hofmanni</i> NOSZKY 76
hontensis	<i>Phoca holitschensis</i> BRÜHL 154
honti	<i>Murex (Alipurpura) holocristatus</i> NOSZKY 87
horogensis	<i>Cythara (Mangelia) hontensis</i> CSEPREGHY-MEZNERICS 105
horvathi	<i>Lathyrus (Dolicolathyrus) hontensis</i> CSEPREGHY-MEZNERICS 106
hosiusi	<i>Nassa (Uzita) hontensis</i> CSEPREGHY-MEZNERICS 108
humilis	<i>Odontostomia hontensis</i> CSEPREGHY-MEZNERICS 108
humilispira	<i>Balanus tintinnabulum</i> honti KOLOSVÁRY 112
hungarica	<i>Thecocystrella horogensis</i> PÁLFY 31
	<i>Graysonites horvathi</i> SZIVES 39
	<i>Glandulina hosiusii</i> FRANZENAU 97
	<i>Carcharodon humilis</i> KOCH 123
	<i>Volutilithes (Neoathleta) consanguinea humilis</i> NOSZKY 93
	<i>Marginella crassula humilispira</i> SZŐTS 54
	<i>Archicoenopsammia hungarica</i> KOLOSVÁRY 48
	<i>Chlamys scabrella hungarica</i> CSEPREGHY-MEZNERICS 101
	<i>Clemmys hungarica</i> SZALAI 127
	<i>Discocyclina hungarica</i> KECSKEMÉTI 41
	<i>Ebalia hungarica</i> MÜLLER 114

- hungarica *Galeodes (Volema) cornuta hungarica* CSEPREGHY-MEZNERICS 106
Hastula hungarica CSEPREGHY-MEZNERICS 106
Lepidocyclina (Eulepidina) hungarica KECSKEMÉTI 68
Manis hungarica KORMOS 147
Microdrillia hungarica BÁLDI 87
Necroteuthis hungarica KRETZOI 94
Petényia hungarica KORMOS 154
Philbertia hungarica CSEPREGHY-MEZNERICS 108
Pholadomyia (Procardia) canavarii hungarica NOSZKY 79
Placosmilia? hungarica KOLOSVÁRY 38
Pseudaspidura hungarica KOLOSVÁRY 96
Rostellaria hungarica NOSZKY 89
Scissurella hungarica SZŐTS 58
Stylacropora hungarica KOLOSVÁRY 48
Testudo hungarica SZALAI 133
Zebina hungarica SZŐTS 60
hungaricum *Prodinotherium hungaricum* ÉHIK 155
Scalpellum hungaricum SZÖRÉNYI 95
hungaricus *Trochocerithium hungaricum* CSEPREGHY-MEZNERICS 110
Amynodon hungaricus KRETZOI 137
Balanus hungaricus KOLOSVÁRY 111
Clavus (Brachytoma) hungaricus BÁLDI 105
Dicerorhinus orientalis hungaricus GAÁL 141
Dolomys hungaricus KORMOS 141
Harpactocarcinus hungaricus TOMOR-THIRRING 63
Lepidopus hungaricus BÖHM 125
Lyreidus hungaricus BEURLEN 95
Mammonteus hungaricus KRETZOI 146
Mesocetus hungaricus KADIĆ 148
Pachygrapsus hungaricus MÜLLER 115
Palicus hungaricus MÜLLER 116
Sciurus whitei hungaricus JÁNOSSY 161
Sirenaeus hungaricus KRETZOI 162
Smerdis hungaricus WEILER 126
Undulocyathus hungaricus KOLOSVÁRY 32
Bison bonasus hungarorum KRETZOI 138
Turricula ilonae BÁLDI 92
Spongotorilispinus inaequispinosus KOZUR, MOIX & MOSTLER 25
Bellerophon (Bucania?) incerta LÓCZY 19
Melania (Balanocochlis) incerta CSEPREGHY-MEZNERICS 107
Reophax incerta FRANZENAU 98
Metula reticulata incisa NOSZKY 87
Turritella incisaformis CSEPREGHY-MEZNERICS 110
Pleurotoma duchatelli incostata NOSZKY 88
Marginulina indifferens HANTKEN 69
Clavatula (Surcula) inermiformis CSEPREGHY-MEZNERICS 104
Hypoxiphothecaella inflata KOZUR, MOIX & OZSVÁRT 22
Megalodus arcuatus inflatus VÉGH-NEUBRANDT 28
Volutilithes intercostaeciliata NOSZKY 93
Dentalina intermedia HANTKEN 41
Strix intermedia JÁNOSSY 135
Echinocardium intermedium LÓCZY 120
Dimydon similis interstricta NOSZKY 73

intersulcatum	<i>Cardium (?Trachycardium) neglectum intersulcatum</i> BÁLDI 72
irregularis	<i>Gaudryina irregularis</i> HANTKEN 68
jabdas	<i>Volutilithes apenninica irregularis</i> NOSZKY 93
jablonszkyi	<i>Terebratula (Waldheimia) jabdas</i> COQUAND 37
janossyi	<i>Fasciolaria jablonszkyi</i> NOSZKY 84
jarnacensis	<i>Cricetus janossyi</i> HÍR 139
jaworskii	<i>Deindsdorfia janossyi</i> REUMER 140
juglans	<i>Megaderma janossyi</i> TOPÁL 148
julieni	<i>Myotis janossyi</i> TOPÁL 150
julii	<i>Pannonicits? janossyi</i> RABEDER 153
kambueheli	<i>Anomia jarnacensis</i> COQUAND 32
kansuensis	<i>Corbuloides jarnacensis</i> COQUAND 32
karinthyi	<i>Pecten jarnacensis</i> COQUAND 33
kashaii	<i>Limopsis retifera jaworskii</i> NOSZKY 76
katalinae	<i>Chlorodiella juglans</i> MÜLLER 113
kayseri	<i>Fusus julieni</i> COQUAND 38
kemencensis	<i>Physocardia julii</i> VÉGH-NEUBRANDT 29
kerekhegyensis	<i>Basiliocostella kambueheli</i> DULAI, BITNER & MÜLLER 40
kerepesiensis	<i>Phillipsia kansuensis</i> LÓCZY 20
kerteszi	<i>Kahlerosphaera karinthyi</i> KOZUR, MOIX & OZSVÁRT 23
kiscellensis	<i>?Metascaphites kashaii</i> SZIVES 39
kiscelliensis	<i>Dufrenoyia katalinae</i> SZIVES 39
kismartonensis	<i>?Nautilus kayseri</i> LÓCZY 19
kissi	<i>Clypeaster kemencensis</i> MEZNERICS 120
kochi	<i>Discorbis kerékhegyensis</i> FRANZENAU 96
kocsisi	<i>Gaudryina kerékhegyensis</i> FRANZENAU 96
koeneni	<i>"Callianassa" kerepesiensis</i> MÜLLER 113
koestleri	<i>Kahlerosphaera kerteszi</i> KOZUR, MOIX & OZSVÁRT 23
kokayi	<i>Sepia kiscellensis</i> WAGNER 94
kondai	<i>Eudolium fasciatum</i> kiscelliensis NOSZKY 84
kopeki	<i>Leda (Lembulus) elata</i> kiscelliensis NOSZKY 75
kormosi	<i>Cibicides kismartonensis</i> FRANZENAU 96
korobkovi	<i>Dibunophyllum kissi</i> KOLOSVÁRY 18
kovacsii	<i>Euthria kochi</i> NOSZKY 84
kovacsiensis	<i>Panopaea (Glycymeris) kochi</i> NOSZKY 78
kreitneri	<i>Lytoceras amplum</i> kocsisi GÉCZY 36
krepuskai	<i>Cytherea (Meretrix) incrassata</i> koeneni NOSZKY 73
kretzoi	<i>Gyroidinoides koestleri</i> OZSVÁRT 42
grenneri	<i>Kahlerosphaera koestleri</i> KOZUR, MOIX & OZSVÁRT 23
Quinqueloculina krenneri	<i>Porcellana (Pisidia) kokayi</i> MÜLLER 118
Volutilithes multicostata	<i>Lokutella kondai</i> VÖRÖS 36
Myophoria kreitneri	<i>Nummulites kopeki</i> KECSKEMÉTI 46
Macrochilina kreitneri	<i>Desmana kormosi</i> SCHREUDER 141
Prospalax kretzoi	<i>Portlandia korobkovi</i> BÁLDI 79
Clavatula (Surcula) krenneri	<i>Modiola mytiloides</i> kovácsi NOSZKY 77
Quinqueloculina krenneri	<i>Nummulina kovácsiensis</i> ROZLOZNIK ex HANTKEN & MADARÁSZ 43
Volutilithes multicostata	<i>Macrochilina kreitneri</i> LÓCZY 19
Myotis kretzoi	<i>Myophoria kreitneri</i> LÓCZY 28
Prospalax kretzoi	<i>Clavatula (Surcula) krenneri</i> CSEPREGHY-MEZNERICS 104
Quinqueloculina krenneri	<i>Quinqueloculina krenneri</i> FRANZENAU 98
Volutilithes multicostata	<i>Volutilithes multicostata</i> krepuskai NOSZKY 93
Myotis kretzoi	<i>Myotis kretzoi</i> TOPÁL 150
Prospalax kretzoi	<i>Prospalax kretzoi</i> JÁNOSSY 156

- kubacskai *Nemopteryx kubacskai* WEILER 126
 kubinyii *Cristellaria (Robulina) kubinyi* HANTKEN 67
Soriculos kubinyii KORMOS 163
 kulcsari *Lucina globulosa* kulcsári NOSZKY 77
 kutassyi *Calyptraea kutassyi* CSEPREGHY-MEZNERICS 104
Hungariella kutassyi SZABÓ 30
Deitanites labatlanensis COMPANY, FÓZY, SANDOVAL & TAVERA 39
 labatlanensis *Titanaster labiostoma* SZÖRÉNYI 67
 labiostoma *Gyroidina laciñata* FRANZENAU 97
 laciniata *Fusus longiroster laevior* NOSZKY 85
 laevisor *Neaera (Cuspidaria) scalarina laevisor* NOSZKY 77
 laevis *Cytherea (Margovoluta) bellardii laevis* NOSZKY 83
Surcula moulensis laevis NOSZKY 90
Telphusograpsus laevis LÖRENTHEY 66
Turbo (Bolma) muricata laevis NOSZKY 91
Volutilithes elevata laevis NOSZKY 93
Allophaiomys laguroides KORMOS 136
Encoiloceras lajosi VÖRÖS 30
Testudo lambrechti SZALAI 133
Tellina (Capsa) lamellosa NOSZKY 80
Genota (Pseudotoma) bonelli lapugyensis CSEPREGHY-MEZNERICS 106
Nummulina lászlói ROZLOZNIK 44
Oxyrhina lata NEUGEBOREN 126
Psammobia (Psammocula) tauroplana lata NOSZKY 79
Siliqua harmati lata NOSZKY 80
Syrnola laterariae BÁLDI 91
Pugilina aequalis lathyroides NOSZKY 89
Chenopus callosus latialatus NOSZKY 82
Archaeoacanthocircus latiannulatus KOZUR, MOIX & OZSVÁRT 21
Pagurus latidactylus MÜLLER & COLLINS 65
Ruesticyrtium latidentatum KOZUR, MOIX & OZSVÁRT 25
Alatipicapora latoalata KOZUR, MOIX & OZSVÁRT 20
Lagotona lázári KRETZOI 144
Erinaceus lechei KORMOS 142
Nucula ledaeformis NOSZKY 78
Balanus legányii KOLOSVÁRY 111
Bittium legányii CSEPREGHY-MEZNERICS 104
Chrysodomus legányii NOSZKY 82
Patellina legányii KENAWY & NYIRÓ 70
Plesiothyreus legányii CSEPREGHY-MEZNERICS 109
"Pylopagurus" legányii MÜLLER 118
Turricula (Knefastia) legányii BÁLDI 92
Turritella leoperdites COQUAND 39
Clavatula letkésensis CSEPREGHY-MEZNERICS 105
Nodosaria letkésiensis FRANZENAU 97
Truncatulina letkésiensis FRANZENAU 98
Ammonites letourneuxi COQUAND 34
Calliphylloceras liasicum GÉCZY 34
Stephanoceras liechtensteinii PAPP 36
Balanus amphitrite litoralis KOLOSVÁRY 111
Ruesticyrtium lobatum KOZUR, MOIX & OZSVÁRT 25
Actaeites lobatus MÜLLER & COLLINS 61
Baranomys lóczyi KORMOS 138

loczyi	<i>Chlorodiella loczyi</i> MÜLLER 113 <i>Cyathophyllum lóczyi</i> FRECH 18 <i>Cybium lóczyi</i> WEILER 123 <i>Fusulinella lóczyi</i> LÖRENTHEY 17 <i>Fusus (Aptyxis) lóczyi</i> NOSZKY 84 <i>Lucina (Myrthea) spinifera lóczyi</i> NOSZKY 77 <i>Parthenope loczyi</i> MÜLLER 116 <i>Perisphinctes lóczyi</i> PAPP 36 <i>Prionechinus loczyi</i> LAMBERT & THIÉRY 120 <i>Scalpellum lóczyi</i> SZÖRÉNYI 95
loerentheyi	<i>Trapezia loerentheyi</i> MÜLLER 66
longicauda	<i>Ficula condita longicauda</i> NOSZKY 84 <i>Fusus retrorsicosta longicauda</i> NOSZKY 85
longicaudata	<i>Xiphothecaella longicaudata</i> KOZUR, MOIX & OZSVÁRT 27
longiceras	<i>Podobursa longiceras</i> longiceras KOZUR, MOIX & OZSVÁRT 24
longicontraseptatum	<i>Plerophyllum (Ufimia) longicontraseptatum</i> KOLOSVÁRY 18
longidentata	<i>Siphonaptera longidentata</i> GÖRÖG 98
longimanus	<i>Diogenes longimanus</i> MÜLLER & COLLINS 62
longinodosa	<i>Echinophoria rondeletii longinodososa</i> NOSZKY 84
longispina	<i>Murex guembeli longispina</i> NOSZKY 87 <i>Spondylus longispina</i> NOSZKY 80
longispinosum	<i>Pararuesticyrtium longispinosum</i> KOZUR, MOIX & OZSVÁRT 23
longus	<i>Trochocyathus longus</i> REUSS 49
lorentheyi	"Carcinus" lórentheyi MÜLLER 113 <i>Panopaea (Glycymeris) lórentheyi</i> NOSZKY 78 <i>Surcula lórentheyi</i> NOSZKY 90
luisae	<i>Galeodea sconsoides luisae</i> NOSZKY 85
lycaonoides	<i>Xenocyon lycaonoides</i> KRETZOI 169
macrognathus	<i>Sorex araneus macrognathus</i> JÁNOSSY 162
macrorhinus	<i>Rhinolophus macrorhinus</i> TOPÁL 157
mactraeformis	<i>Thecocystathus mactraeformis</i> KOLOSVÁRY 32
madachi	<i>Harpa bellardii madáchi</i> NOSZKY 86
madaraszi	<i>Nummulites madaraszi</i> HANTKEN 46
magnum	<i>Calliphylloceras altisulcatum magnum</i> GÉCZY 34
magnus	<i>Achaeus magnus</i> MÜLLER 112 <i>Petrolisthes magnus</i> MÜLLER 117
maior	<i>Nesodon imbricatus maior</i> KRETZOI 151 <i>Nummulites subtilis maior</i> KECSKEMÉTI 47
majeri	<i>Arcopagia majeri</i> SZÖTS 49
major	<i>Favosites goldfussi major</i> FRECH 18
majori	<i>Vespertilio majori</i> KORMOS 169
majzoni	<i>Eponides majzoni</i> NYÍRŐ 68 <i>Nodosaria (Dentalina) majzoni</i> JASKÓ 69 <i>Nummulites majzoni</i> KECSKEMÉTI 46 <i>Tellina (Peronidea) budensis majzoni</i> NOSZKY 80
makadii	<i>Iharkutosuchus makadii</i> ÓSI, CLARK & WEISHAMPEL 131
malyinkae	<i>Polycoelia malyinkae</i> KOLOSVÁRY 19
manyense	<i>Cardium manyense</i> KÓKAY 99
maresi	<i>Isocardia maresi</i> COQUAND 32
margarethae	<i>Galeodea echinophora margarethae</i> NOSZKY 85
margaritacea	<i>Turritella sandbergeri margarethae</i> GAÁL 92
margaritodon	<i>Heterostegina margaritacea</i> MAJZON 68 <i>Sorex margaritodon</i> KORMOS 163

- marginatostrata
 marginatus
 marginuliniformis
 margitae
 mariae
 martelina
 mathiasi

 matraverebelyensis
 matrensis
 maubeugei
 mazana
 medioumbonata
 menisa
 meridionalis
 merlai
 mersinensis

 merzbacheri
 michelottii
 microdon
 microstriatus
 mikszathi
 milleri
 minima

 minor
 minuta

 minutus
 miocaenica
 miocaenicus
 miocenicus
 mitraeformis
 moixi
 montisantonii
 morensis
 mostleri
 mucronata
 multifilosa
 multispinosa

 multisulcata
 munieri
 nabdalsae
 naefi
 nankingensis
 manus

Lima (Acesta) miocenica marginatostrata NOSZKY 76
Anapagurus marginatus MÜLLER 112
Amphicoryne marginuliniformis NYÍRÖ 67
Cryptoplax margitae DULAI 98
Solarium mariae BÁLDI 90
Mustela martelina PETÉNYI 150
Charybdis (Goniosupradens) mathiasi MÜLLER 113
Chenopus mathiasi NOSZKY 82
Actinometra matraverebelyensis SZALAI 120
Diogenes matrensis MÜLLER 114
Phylloceras? baconicum maubeugei GÉCZY 36
Rhynchonella mazana COQUAND 36
Saxicava rugosa medioumbonata NOSZKY 79
Pecten menisa COQUAND 33
Rhynchonella meridionalis COQUAND 37
Hammatoceras planinsigne merlai GÉCZY 35
Hypoxiphotheca mersinensis KOZUR, MOIX & OZSVÁRT 22
Pararuesticyrtium mersinensis KOZUR, MOIX & OZSVÁRT 23
Podobura mersinensis KOZUR, MOIX & OZSVÁRT 24
Zhamojdasphaera latispinosa mersinensis KOZUR, MOIX & MOSTLER 27
Pleuromya merzbacheri PAPP 33
Murex (Phyllonotus) rufus michelottii NOSZKY 87
Promimomys microdon JÁNOSSY 156
Pyrgo microstriatus OZSVÁRT 47
Umbrella mikszathi NOSZKY 92
Dolomys milleri NEHRING 142
Eoplax minima MÜLLER & COLLINS 63
Lamna minima NEUGEBOREN 125
Sphenia minima CSEPREGHY-MEZNERICS 103
Crassatella (Crassatina) bosqueti minor BÁLDI 73
Pecten (Amussiopecten) burdigalensis minor TELELDI ROTH 79
Cristellaria minuta HANTKEN 67
Lamina minuta NEUGEBOREN 125
Stylocyathus minutus KOLOSVÁRY 38
Tritodynamia miocaenica MÜLLER 119
Portunus (Monomia) miocaenicus MÜLLER 118
Anapagurus miocenicus MÜLLER 112
Lyria taurinia mitraeformis NOSZKY 86
Spongotorilispinus moixi KOZUR & MOSTLER 25
Cerithium calcaratum montis-antonii SZŐTS 53
Trinacria móreensis SZŐTS 50
Ruesticyrtium mostleri KOZUR, MOIX & OZSVÁRT 25
Textularia carinata mucronata JASKÓ 70
Melanopsis doboi multifilosa SCHRÉTER 121
Karnospongella multispinosa KOZUR, MOIX & MOSTLER 23
Spinomersinella multispinosa KOZUR, MOIX & OZSVÁRT 25
Latrunculus (Peridipsacus) caronis multisulcata NOSZKY 86
Rissoa munieri SZŐTS 58
Turbo nabdalsae COQUAND 34
Archaeosepia naefi SZÖRÉNYI 60
Lingulina nankingensis LÖRENTHEY 17
Cricetus cricetus nanus SCHAUB 140
Ungaromys nanus KORMOS 164

nassaeformis	<i>Egerea collectiva nassaeformis</i> NOSZKY 84
naticoides	<i>Deshayesia naticoides</i> SZÓTS 54
neglecta	<i>Petényia neglecta</i> KRETZOI 154
nehringi	<i>Dolomys nehringi</i> KRETZOI 142
neocomiensis	<i>Heliopora neocomiensis</i> KOLOSVÁRY 38
neogenica	<i>Dromia neogenica</i> MÜLLER 114
neogenicus	<i>Portunus neogenicus</i> MÜLLER 118
neogradiensis	<i>Antedon neogradiensis</i> SZALAI 120
neoscalarina	<i>Cuspidaria neoscalarina</i> BÁLDI 73
nicaisei	<i>Rhynchonella nicaisei</i> COQUAND 37
niobe	<i>Spiriferina nicaisei</i> COQUAND 37
nitens	<i>Turbo nicaisei</i> COQUAND 34
nodosa	<i>Coelosmilia niobe</i> KOLOSVÁRY 38
nogradensis	<i>Adeorbisina nitens</i> SZABÓ 33
	<i>Clavatula (Perrona) vindobonensis nodosa</i> CSEPREGHY-MEZNERICS 105
	<i>Teredo anguinea nodosa</i> NOSZKY 81
	<i>Astarte (Goodallia) triangularis nigradensis</i> CSEPREGHY-MEZNERICS 99
	<i>Cardita (Cardiocardita) nigradensis</i> Cs. MEZNERICS 99
	<i>Clavatula nigradensis</i> CSEPREGHY-MEZNERICS 105
	<i>Lymnaea pachygaster nigradensis</i> CSEPREGHY-MEZNERICS 106
	<i>Mactra nigradensis</i> CSEPREGHY-MEZNERICS 102
	<i>Rissoina (Zebinella) nigradensis</i> CSEPREGHY-MEZNERICS 109
nopcsai	<i>Trionyx nopcsai</i> SZALAI 134
noszkyi	<i>Andromacheia noszkyi</i> KOLOSVÁRY 111
novemdentatum	<i>Asthenotoma noszkyi</i> BÁLDI 81
nuda	<i>Cardiomya noszkyi</i> BÁLDI 72
nudata	<i>Ceratotrochus noszkyi</i> KOLOSVÁRY 37
nudus	<i>Drillia noszkyi</i> Cs. MEZNERICS 105
nysti	<i>Hinia (Telasco) schlotheimi noszkyi</i> BÁLDI 86
obesaformis	<i>Melongena noszkyi</i> GAÁL 87
obovatooides	? <i>Turbinoberis noszkyi</i> KOLOSVÁRY 49
obtusa	<i>Undulocyathus noszkyi</i> KOLOSVÁRY 32
obtusata	<i>Ruesticyrtium georgi novemdentatum</i> KOZUR, MOIX & OZSVÁRT 25
octospinosa	<i>Callianassa nuda</i> BEURLEN 95
oersi	<i>Dolichotoma collaris nudata</i> NOSZKY 83
oligocaenica	<i>Murex trigonalis nudus</i> NOSZKY 87
oligocenica	<i>Parvisipho nudus</i> SZÓTS 57
oligocenicum	<i>Nucula praeelongata nysti</i> NOSZKY 78
oligodon	<i>Montlivaltia obesaformis</i> KOLOSVÁRY 38
oligoobscura	<i>Glycymeris (Glycymeris) latiradiata obovatooides</i> BÁLDI 74
oligotuberculatus	<i>Crocidura obtusa</i> KRETZOI 140
oppenheimi	<i>Gibbula (Colliculus) biangula obtusata</i> CSEPREGHY-MEZNERICS 106
orbicularis	<i>Ethusa octospinosa</i> MÜLLER 114
	<i>Ebalia oersi</i> MÜLLER 114
	<i>Trinacria oligocaenica</i> BÁLDI 81
	<i>Cytherocardia oligocenica</i> NOSZKY 73
	<i>Drillia crispata oligocenica</i> NOSZKY 83
	<i>Cardium (Ringicardium) grateloupi oligocenicum</i> NOSZKY 72
	<i>Amblycoptus oligodon</i> KORMOS 137
	<i>Trochus (Solariella) oligoobscura</i> NOSZKY 91
	<i>Paguristes oligotuberculatus</i> MÜLLER & COLLINS 64
	<i>Nummulina oppenheimi</i> ROZLOZNIK 44
	<i>Lucina (Myrthea) rectangularis orbicularis</i> NOSZKY 77

- orchardi
orientalis
- orientoromana
orlovi
ornatissima
ornatissimus
osborni
ostramosensis
ostramosi
ouarsenissensis
ovalis
ovata
ovatus
ozsvarti
palatinus
palermineus
palfyi
pamuki
pancici
pannoniae
pannonica
- pannonicus
- pappi
- paradaubentoni
paratethyensis
parva
parvispira
parvula
parvus
patella
pectinata
pentacantha
pentagonalis
penzesgyoerensis
peracuta
peraffinis
- Xiphophecaella orchardi* KOZUR, MOIX & OZSVÁRT 27
Chonetes orientalis LÓCZY 20
Moniliopsis (Bathytona) cataphracta orientalis CSEPREGHY-MEZNERICS 107
Pitaria lamarcki orientalis CSEPREGHY-MEZNERICS 103
Clavatula orientoromana BÁLDI 105
Eomellivora orlovi KRETZOI 142
Dorippe ornatissima MÜLLER 114
Vermetus ornatissimus SZŐTS 59
Hemilophodon osborni KRETZOI 143
Mimomys ostramosensis JÁNOSSY & VAN DER MEULEN 148
Erinaceus ostramosi JÁNOSSY 143
Trochus ouarsenissensis COQUAND 34
Archaeocanthoncircus ovalis KOZUR, MOIX & OZSVÁRT 21
Globularia ovata BÁLDI 85
Erycites ovatus GÉCZY 34
Spongotorilispinus ozsvartii KOZUR, MOIX & MOSTLER 26
Pusionella pseudofusus palatinus STRAUSZ 109
Foetorius palermineus PETÉNYI 143
Veloruesticyrtium palfyi KOZUR, MOIX & OZSVÁRT 26
Kahlerosphaera pamuki KOZUR, MOIX & OZSVÁRT 23
Gervillia pancici RADOVANOVIĆ 32
Dystychoceras pannoniae KRETZOI 142
Abra pannonica SZŐTS 49
Eomaldivia pannonica MÜLLER & COLLINS 63
Leda (Yoldia) perovalis pannonica NOSZKY 75
Marginella pannonica SZŐTS 55
Microsolena rotula pannonica KOLOSVÁRY 31
Nummulina striata pannonica ROZLOZNIK 44
Odostomia pannonica SZŐTS 56
Pachyura pannonica KORMOS 152
Praepusa pannonica KRETZOI 155
Sabatia (Mnestia) pannonica CSEPREGHY-MEZNERICS 109
Sphyraena pannonica WEILER 127
Balanus pannonicus KOLOSVÁRY 111
Gadus (Merlangus) pannonicus KOCH 124
Hubacyon (Kanicyon) pannonicus KRETZOI 143
Lagurus pannonicus KORMOS 144
Limnocardium pappi STRAUSZ 102
Neritopsis pappi KUTASSY 30
Pholas (Martesia) pappi SZŐTS 50
Myotis paradaubentoni TOPÁL 151
Pilumnopeus paratethyensis MÜLLER 118
Galathea (Acanthogalathea) parva MÜLLER & COLLINS 63
Lithoconus ineditus parvispira NOSZKY 86
Bulimina parvula FRANZENAU 67
Palaeograpsus parvus MÜLLER & COLLINS 65
Spirillina plana patella LÖRENTHEY 17
Vulvulina pectinata HANTKEN 48
Protomunida pentacantha MÜLLER & COLLINS 66
Kromtis pentagonalis MÜLLER & COLLINS 64
Nummulites penzesgyoerensis KECSKEMÉTI 46
Lucina spissistriata peracuta NOSZKY 77
Ammonites peraffinis COQUAND 34

peralta	<i>Lucina (Myrthea) spinifera peralta</i> NOSZKY 77
	<i>Nucula sulcifera peralta</i> NOSZKY 78
	<i>Turbo (Senectus) ranellaformis peralta</i> NOSZKY 92
percarinata	<i>Turritella beyrichi percarinata</i> TELELDI ROTH 92
percassus	<i>Echinolampas percrassus</i> MEZNERICS 120
percrecca	<i>Anas crecca percrecca</i> JÁNOSSY 134
perforans	<i>Nerinea perforans</i> COQUAND 33
pertenuis	<i>Nodosaria pertenuis</i> FRANZENAU 97
perturbata	<i>Plectina perturbata</i> FRANZENAU 97
petenyii	<i>Mimomys petényii</i> MÉHELY 148
	<i>Prodeinotherium petenyii</i> VÖRÖS 155
	<i>Vormela petényii</i> KRETZOI 169
peyssoneli	<i>Terebratula peyssoneli</i> COQUAND 37
pilgrimi	<i>Pannonicits piligrimi</i> KORMOS 153
pilisensis	<i>Theodoxus (Vittoclithon) pilisensis</i> BÁLDI 91
pisum	<i>Ampullaria pisum</i> COQUAND 33
pitymyoides	<i>Mimomys pitymyoides</i> JÁNOSSY & VAN DER MEULEN 149
plana	<i>Acirsia plana</i> CSEPREGHY-MEZNERICS 104
	<i>Pseudopatellina plana</i> KENAWY & NYÍRÓ 70
	<i>Terebratulina plana</i> MEZNERICS 111
	<i>Vaginulina plana</i> NYÍRÓ 70
planidentatus	<i>Pilumnomimus planidentatus</i> MÜLLER & COLLINS 65
planus	<i>Corallicarcinus planus</i> MÜLLER & COLLINS 62
platyodon	<i>Amphicynops platyodon</i> KRETZOI 137
platysoma	<i>Conocyclus platysoma</i> SZÖRÉNYI 66
plesiobathystoma	<i>Echinolampas plesiobathystoma</i> SZÖRÉNYI 66
plioacaenius	<i>Allophaiomys plioacaenius</i> KORMOS 136
	<i>Plecotus (Plecotus) plioacaenius</i> TOPÁL 154
plioacaenus	<i>Lates plioacaenus</i> KOCH 125
polonica	<i>Cythara (Mangelia) vulpecula polonica</i> CSEPREGHY-MEZNERICS 105
polydectes	<i>Stephanosmilia polydectes</i> KOLOSVÁRY 38
pongraczi	<i>Axinus (Cryptodon) subangulatus pongráczi</i> NOSZKY 72
porvaensis	<i>Triloculina porvaensis</i> HANTKEN 47
postdelphinensis	<i>Rhinolophus postdelphinensis</i> TOPÁL 157
posterior	<i>Pliomys posterior</i> JÁNOSSY 154
posthumus	<i>Rabdocidaris posthumus</i> PÁVAY 67
praebonasia	<i>Tetrastes praebonasia</i> JÁNOSSY 135
praedesertum	<i>Limnocardium praedesertum</i> STRAUSZ 102
praeglacialis	<i>Alopex praeaglacialis</i> KORMOS 136
	<i>Cricetus cricetus praeaglacialis</i> SCHAUB 140
	<i>Rhinolophus euryale praeaglacialis</i> KORMOS 156
praeloriger	<i>Sicista praeloriger</i> KORMOS 161
praenivalis	<i>Mustela praenivalis</i> KORMOS 150
praenoriticus	<i>Neomegalodon (Gemmelarodus) paronai praenoriticus</i> VÉGH-NEUBRANDT 28
praepapillosum	<i>Cardium (Parvicardium) praepapillosum</i> BÁLDI 72
praespinulosa	<i>Creusia spinulosa praespinulosa</i> KOLOSVÁRY 112
praestrigosus	<i>Jujubinus (Strigosella) multicingulatus praestrigosus</i> BÁLDI 86
praetimidus	<i>Lepus praetimidus</i> KRETZOI in JÁNOSSY 145
praetypica	<i>Mastodon (Mammuth) americanus praetypica</i> SCHLESINGER 147
	<i>Trilophodon angustidens praetypica</i> TASNÁDI KUBACSKA 164
primigenius	<i>Citellus primigenius</i> KORMOS 139
primitivus	<i>Cymonomus primitivus</i> MÜLLER & COLLINS 62
prisca	<i>Leda (Costatoleda) psammobiaeformis prisca</i> NOSZKY 75

- procera *Xiphothecaella procera* KOZUR, MOIX & OZSVÁRT 27
 procerus *Clavulinoides procerus* OZSVÁRT 41
 profunda *Lucina spinifera profunda* NOSZKY 77
 profundiformis *Polycoelia profundiformis* KOLOSVÁRY 19
 progressa *Pachycrocuta robusta progressa* KRETZOI 151
 progressus *Baranomys (Warthamys) kowalskii progressus* JÁNOSSY 138
"Pliomys" progressus KRETZOI 154
Pecten promontorensis CSEPREGHY-MEZNERICS 103
 propinqua *Cristellaria propinqua* HANTKEN 67
psammobiaeformis *Leda (Costatoleda) psammobiaeformis* TELELDI ROTH 75
psammophila *Mioxaiava psammophila* MÜLLER 115
pseudoacuticosta *Lagenia pseudoacuticosta* FRANZENAU & MAJZON 97
pseudoallixi *Persicula pseudoallixi* SZŐTS 57
pseudobarbata *Arca (Barbatia) pseudobarbata* SZALAI 99
pseudobliquistoma *Pithocerithium pseudobliquistoma* SZALAI 109
pseudocarinata *Siphonina pseudocarinata* NYÍRÓ 70
pseudodeformis *Calyptarea pseudodeformis* BÁLDI 82
pseudogamlitzensis *Pirenella picta pseudogamlitzensis* STRAUSZ 108
pseudohungaricus *Cadulus pseudohungaricus* SZŐTS 50
pseudomumiola *Cingula pseudomumiola* SZŐTS 53
pseudonana *Marginella pseudonana* SZŐTS 55
pseudonassoides *Raphitoma pseudonassoides* BÁLDI 89
pseudoruellensis *Odostomia pseudoruellensis* SZŐTS 57
pseudoscharbergana *Nodosaria pseudo-scharbergana* FRANZENAU 97
pseudospinulosa *Cristellaria pseudo-spinulosa* FRANZENAU 96
pseudosubangulatus *Anisus pseudosubangulatus* SZŐTS 51
pteriiforme *Dicerocardium pteriiforme* VÉGH-NEUBRANDT 28
pulcherrima *Bythinella pulcherrima* SZŐTS 52
punctata *Dymia fragilis punctata* NOSZKY 73
pupoides *Erato laevis punctata* CSEPREGHY-MEZNERICS 106
purbeckensis *Genota ramosa pupoides* CSEPREGHY-MEZNERICS 106
pygmaea *Corbula purbeckensis* COQUAND 32
pyruloides *Cyrena sirena pygmaea* NOSZKY 73
quadrangleculata *Egeraea collectiva pyruloides* NOSZKY 84
quadratum *Assiminea quadrangulata* SZŐTS 51
quadricingulata *Calliphylloceras altisulcatum quadratum* GÉCZY 34
quadricostata *Trochosmilia 4-cingulata* KOLOSVÁRY 49
quinquepetallus *Seila quadricostata* SZŐTS 58
racmecskeensis *Antedon quinquepetallus* SZALAI 120
radiata *Testudo rácmecskeensis* SZALAI 133
radula *Myophoria radiata* LÓCZY 28
rakosense *Trachypirimela radula* MÜLLER 119
rakosensis *Chlamys rákosense* CSEPREGHY-MEZNERICS 100
rakosiensis *Asthenognathus rakosensis* MÜLLER 112
rakuszi *Pagurus rakosensis* MÜLLER 116
ranellaeformis *Quinqueloculina rákosiensis* FRANZENAU 98
rara *Plerophyllum (Ufimia) rakuszi* KOLOSVÁRY 19
raricosta *Turbo (Senectus) ranellaeformis* NOSZKY 92
raricostata *Cerithiella rara* SZŐTS 53
raricostatum *Emarginula elongata raricosta* NOSZKY 84
rarilamellosa *Styliola maxima raricostata* NOSZKY 90
Latirus cognatus rarilamellosa NOSZKY 86

rarinodosa	<i>Ranella (Apollon) gigantea rarinodosa</i> NOSZKY 89
raristriata	<i>Surcula moulinsi rarisstriata</i> NOSZKY 90
rarivaricosa	<i>Murex (Haustellum) sismondae rarivaricosa</i> NOSZKY 87
ratoti	<i>Triadomegalodon rátóti</i> VÉGH-NEUBRANDT 30
recta	<i>Tellina (Peronea) nitida recta</i> NOSZKY 81
rectangularis	<i>Archaeoacanthocircus rectangularis</i> KOZUR, MOIX & OZSVÁRT 21
reitzi	<i>Nodosaria (Dentalina) reitzi</i> HANTKEN 69
rejtoei	<i>Kahlerosphaera rejtoei</i> KOZUR, MOIX & OZSVÁRT 23
reticulata	<i>Dolichotoma subdenticulata reticulata</i> NOSZKY 83
retifera	<i>Chenopus mathiasi retifera</i> NOSZKY 82
rex	<i>Echinophoria intermedia retifera</i> NOSZKY 84
rigida	<i>Nucula sulcifera retifera</i> NOSZKY 78
rigoi	<i>Pleurotomaria retifera</i> NOSZKY 88
robusta	<i>Turbo (Senectus) ranellaformis retifera</i> NOSZKY 92
robustus	<i>Mimomys rex</i> KORMOS 149
rogeri	<i>Sculptoplax rigida</i> MÜLLER & COLLINS 66
romerii	<i>Zhamojdasphaera rigoi rigoi</i> KOZUR, MOIX & MOSTLER 27
romhanyensis	<i>Surcula regularis robusta</i> NOSZKY 91
rostrata	<i>Myotis bechsteini robustus</i> TOPÁL 150
rotaridesi	<i>Erycites ovatus? rogeri</i> GÉCZY 35
rotundidorsatum	<i>Spondylus rogeri</i> CSEPREGHY-MEZNERICS 80
rotundum	<i>Turritella romerii</i> SZŐTS 59
rozlozsni	<i>Pulvinulina romhanyensis</i> FRANZENAU 70
rugosa	<i>Aloidis carinata rostrata</i> CSEPREGHY-MEZNERICS 99
rugosum	<i>Barbastella rostrata</i> TOPÁL 138
rupelica	<i>Volutilithes rotaridesi</i> NOSZKY 94
rupeliensis	<i>Haplophragmium rotundidorsatum</i> HANTKEN 68
saccoi	<i>Flabellum rotundum</i> KOLOSVÁRY 48
sanctipetri	<i>Pholadomya (Procardia) canavarii rozlozsni</i> NOSZKY 79
sandbergeri	<i>Tympanotonus rozlozsni</i> SZŐTS 59
sasselloniensis	<i>Melanopsis doboi rugosa</i> SCHRÉTER 121
scabra	<i>Nodophthalmidium rugosum</i> GÖRÖG 97
scalena	<i>Heterostegina rupelica</i> MAJZON 68
sceptrumides	<i>Thaumastocheles rupeliensis</i> BEURLEN 95
schafarziki	<i>Galeodocassis fuchsii saccoi</i> NOSZKY 85
schaubi	<i>Placenta saccoi</i> NOSZKY 79
schlichti	<i>Umbrella sancti-petri</i> NOSZKY 93
scholzi	<i>Lucina (Cardiolucina) striatula sandbergeri</i> NOSZKY 77
schröteri	<i>Aequipecten deletus sasselloniensis</i> NOSZKY 71
schwageri	<i>Upogebia scabra</i> MÜLLER 119
scrobiculatus	<i>Bulimina scalena</i> FRANZENAU 96
	<i>Podobursa sceptrumides</i> KOZUR, MOIX & OZSVÁRT 24
	<i>Perna maxillata schafarziki</i> NOSZKY 79
	<i>Pholadomya schafarziki</i> PAPP 33
	<i>Myotis schaubi</i> KORMOS 151
	<i>Nummulites schaubi</i> KECSKEMÉTI 47
	<i>Glandulina schlichti</i> FRANZENAU 97
	<i>?Metascaphites scholzi</i> SZIVES 39
	<i>Cladocora schréteri</i> KOLOSVÁRY 37
	<i>Leda (Yoldia) schréteri</i> NOSZKY 75
	<i>Pecten (Aequipecten) schréteri</i> NOSZKY 79
	<i>Cristellaria schwageri</i> HANTKEN 41
	<i>Pycnodus scrobiculatus</i> REUSS 126

- semicostata
semilaevis
seminuda
semistriata
semseyi
semseyiana
serrata
serratulum
setosa
silasensis
simplex
sinuosa
skofleki
solitaria
solymarensis
soosi
soproniensis
speciosa
spelaeoides
spinicosta
spiniferus
spinosa
spiralostriata
splendens
splendidum
ssemenowi
stehlini
steiningeri
stellata
stephani
strandii
strangulatus
stredae
stredai
striata
striatissimus
strigosa
stromeri
subcolumbella
subcrebricosta
subcylindrica
- Cassidaria nodosa semicostata* NOSZKY 82
Dentalina semilaevis HANTKEN 68
Nassa (Hima) turbinella semilaevis CSEPREGHY-MEZNERICS 108
Lingulina costata seminuda HANTKEN 69
Odostomia semistriata SZŐTS 57
Neohyaenodon semseyi KRETZOI 151
Pyrula (Melongena) semseyiana ERDŐS 89
Lamna (Odontaspis) serrata NEUGEBOREN 125
Cardium (Ringicardium) bükkianum serratulum NOSZKY 72
Dentalina setosa HANTKEN 68
Mimomys silasensis JÁNOSSY 149
Cominella hungarica simplex NOSZKY 83
Estramomys simplex JÁNOSSY 143
Nanomaja simplex MÜLLER & COLLINS 64
Nodosaria (Dentalina) simplex HANTKEN 69
Nodosinella simplex LÖRENTHEY 17
Ranella (Apollon) semilaevis simplex NOSZKY 89
Surcula beyrichii simplex NOSZKY 90
Terebra simplex TELELDI ROTH 91
Turbo (Senectus) ranellaeformis simplex NOSZKY 92
Lucina (Myrthea) spinifera sinuosa NOSZKY 77
Karstocricetus skofleki KORDOS 144
Mitrella solitaria BÁLDI 87
Cricetus runtonensis solymarensis HÍR 140
Nuculana solymarensis BÁLDI 78
Cytherea (Callista) taurorugosa soósi NOSZKY 73
Congeria soproniensis VITÁLIS 102
Lamna speciosa NEUGEBOREN 125
Canis spelaeoides KRETZOI 139
Semipecten spinicosta NOSZKY 80
Trophon varicosissimus spiniferus CSEPREGHY-MEZNERICS 110
Alatipicrapora spinosa KOZUR, MOIX & OZSVÁRT 20
Nassa (?Leiodomus) sturi spiralostriata CSEPREGHY-MEZNERICS 108
Marginulina splendens HANTKEN 69
Vexillum splendidum SZŐTS 60
Rhacophyllites ssemenowi PAPP 36
Ciconia stehlini JÁNOSSY 134
Haydnella steiningeri MÜLLER 114
Myotis steiningeri KORMOS 151
Isocrinus stellata SZALAI 120
Murex (Harmatia) stephani NOSZKY 87
Emys strandii SZALAI 127
Fusus strangulatus COQUAND 38
Hungariella stredae KUTASSY 30
Hammatoceras planiforme stredai GÉCZY 35
Patella protea stredai NOSZKY 88
Flabellina striata HANTKEN 41
Petrolisthes? striatissimus MÜLLER & COLLINS 65
Xiphothecaella strigosa KOZUR, MOIX & OZSVÁRT 27
Putorius stromeri KORMOS 156
Megalodon subcolumbella KOKEN 28
Mitra subcrebricosta SZŐTS 56
Marginella subcylindrica SZŐTS 55

- subfallax *Angulus (Peronidia) nysti subfallax* BÁLDI 71
 subfichteli *Glycymeris (Glycymeris) latiradiata subfichteli* BÁLDI 75
 subflabelliformis *Textilaria subflabelliformis* HANTKEN 47
 subfragilis *Spiriferina subfragilis* LÓCZY 31
 subglobosa *Dromilites subglobosa* MÜLLER & COLLINS 62
 sublaevigata *Clavatula sublaevigata* BÁLDI 105
 submelba *Apus submelba* JÁNOSSY 134
 submisera *Odostomia submisera* SZÖTS 57
 subparvocostata *Cardita (Cyclocardia) orbicularis subparvocostata* BÁLDI 72
 subplanulata *Nummulina subplanulata* ROZLOZNIK ex HANTKEN & MADARÁSZ 44
 subquadratus *Erycites subquadratus* GÉCZY 35
 subrotundatum *Circulum subrotundatum* SZÖTS 50
 subtapiroidea *Mastodon (Bunolophodon) angustidens subtapiroidea* SCHLESINGER 147
 subtrifaria *Seila subtrifaria* SZÖTS 58
 suemegensis *Menabites (Delawarella) suemegensis* FŐZY 39
 sulcata *Nummulites suemegensis* KECSEKÉTI 47
 Branchioplax *sulcata* MÜLLER & COLLINS 61
 Neaera (*Cuspidaria*) *sulcata* NOSZKY 78
 sulcatus *Plagiolophus sulcatus* BEURLEN 95
 sulcosa *Neaera (Tergulina) sulcosa* NOSZKY 78
 sulcosus *Spondylus sulcosus* NOSZKY 80
 Phyllosmilia *sümegensis* KOLOSVÁRY 38
 sumeghyi *Congeria sumeghyi* STRAUSZ 102
 superba *Frondicularia superba* HANTKEN 68
 supraoligocaenicus *Theodoxus (Vittocliton) supraoligocaenicus* BÁLDI 91
 supravariabilis *Odostomia supravariabilis* SZÖTS 57
 supravarians *Bayania supravarians* SZÖTS 52
 suttoensis *Testudo süttöensis* SZALAI 134
 szaboi *Clavulina szaboi* HANTKEN 41
 Parthenope *szaboi* MÜLLER 116
 szalaii *Sphaenia benghami szalaii* NOSZKY 80
 szechenyii *Lingulina széchenyi* LÖRENTHEY 17
 szekessyi *Loxonema széchenyi* LÓCZY 19
 Myophoria *széchenyi* LÓCZY 28
 sizerbi *Volutilithes (Athleta) consanguinea székessyi* NOSZKY 93
 szobensis *Kahlerosphaera sizerbi* KOZUR, MOIX & OZSVÁRT 23
 ?*Callianassa szobensis* MÜLLER 113
 szoerenyiae *Drillia granaria szobensis* CSEPREGHY-MEZNERICS 105
 szontaghi *Fissurina szobensis* FRANZENAU 96
 szorennyiae *Odontostomia szobensis* CSEPREGHY-MEZNERICS 108
 szorennyiae *Taurasia szobensis* CSEPREGHY-MEZNERICS 110
 szorennyiae *Hammatoceras tenerum* SZOERENYIAE GÉCZY 35
 szorennyiae *Thia szorennyiae* MÜLLER 119
 szorennyiae *Cardium (Parvicardium) transversale* SZONTAGHI NOSZKY 72
 szorennyiae *Euthriofusus szontaghi* NOSZKY 84
 szorennyiae *Lyria collaris szontaghi* NOSZKY 86
 szorennyiae *Aequipecten oligosquamosus szörényii* NOSZKY 71
 szotsi *Flabellum szótsi* KOLOSVÁRY 48
 szutschuanensis *Camarophoria szü-tschan-ensis* LÓCZY 20
 taegeri *Beguina taegeri* SZÖTS 50
 tardosi *Anomalospalax tardosi* KORDOS 137
 tarkoensis *Rhinolophus ferrumequinum tarkoensis* TOPÁL 156
 tasnadii *Aequipecten raulini tasnádii* NOSZKY 71

- tasnadii
 telegddii
 telegdirothi
 telegyi
 tellinoides
 tenerinodosa
 tergolaevius
 tetenyensis
 tetrapedis
 tetraspinosa
 tevesthenensis
 textilearoides
 thraciaeformis
 toarcense
 tokodensis
 topali
 tormai
 tornensis
 tortonica
 tothi
 transdanubica
 transiens
 transitus
 transsilvanica
 transylvanica
 transylvanicus
 triassica
 tricosta
 tricostata
 tricuspidata
 tridentata
 trigonalis
 trigonostomea
 triquetra
 trispinosa
 trispinosus
 trivaricosa
 trochiformis
 tuberosus
 tudiclaeformis
 tunicata
 turcensis
Sorex tasnádii KRETZOI 163
Pilumnus telegddii MÜLLER 118
Flabellipecten telegdi-rothi CSEPREGHY-MEZNERICCS 74
Harpactocarcinus telegdi-rothi TOMOR-THIRRING 63
Mitromorpha (Antimitra) telegdirothi BÁLDI 87
Surcula telegdirothi NOSZKY 91
Thracia telegdirothi NOSZKY 81
Volutilithes telegyi GAÁL 94
Gastrana tellinoides CSEPREGHY-MEZNERICCS 102
Volutilithes (Athleta) pygmaea tenerinodosa NOSZKY 94
Pterocera (Alaria) rochatiana tergolaevius NOSZKY 89
Chlorodiella mediterranea tetenyensis MÜLLER 113
Parthenope tetenyensis MÜLLER 116
"Pilumnopeus" tetenyensis MÜLLER 118
Alatipacipora tetrapedis KOZUR, MOIX & OZSVÁRT 21
Podobursa longiceras tetraspinosa KOZUR, MOIX & OZSVÁRT 24
Fusus tevesthenensis COQUAND 39
Gaudryina textilaroides HANTKEN 42
Lyonsia thraciaeformis NOSZKY 77
Alocolytoceras ophioneum toarcense GÉCZY 34
Argyrotheca tokodensis BITNER & DULAI 60
Amblycoptus topali JÁNOSSY 137
Hungarosaurus tormai ÓSI 127
Episoriculus tornensis JÁNOSSY 142
Mimomys tornensis JÁNOSSY & VAN DER MEULEN 149
Cymbaloporella tortonica FRANZENAU & MAJZON 96
Pliolagus tóthi KRETZOI 154
Pinna transdanubica VÉGH 29
Charonia (Sassia) tarbelliana transiens BÁLDI 82
Lagurus transiens JÁNOSSY 144
Archaeoanthocircus transitus KOZUR, MOIX & OZSVÁRT 21
Ditrupa transsilvanica MEZNERICCS 110
Goniocypoda transsilvanica BITTNER 63
Mitra goniophora transsylvania CSEPREGHY-MEZNERICCS 107
Balanus transylvanicus KOLOSVÁRY 112
Magyarosaurus transylvanicus HUENE 133
Serranus transylvanicus BÖHM 126
Spinoprotunuma triassica KOZUR, MOIX & OZSVÁRT 25
Amussium semiradiatum tricosta NOSZKY 71
Triphora tricostata SZÖTS 59
Vaginella tricuspidata ZORN & JANSEN 93
Calappa tridentata BEURLEN 95
Leda modesta trigonalis NOSZKY 75
Quinqueloculina ermani trigonostomea FRANZENAU 98
Bulimina triquetra FRANZENAU 67
Eomaldivia trispinosa MÜLLER & COLLINS 63
Daragrapsus trispinosus MÜLLER & COLLINS 62
Ocinebrina crassilabiata trivaricosa BÁLDI 88
Alvania montagui trochiformis CSEPREGHY-MEZNERICCS 104
Textularia tuberosus OZSVÁRT 47
Triton tudiclaeformis NOSZKY 91
Cristellaria (Marginulina) tunicata HANTKEN 67
Cornucardia turcensis VÉGH-NEUBRANDT 27

turcocampestris	<i>Actaea turcocampestris</i> MÜLLER 112
turcus	? <i>Pagurus turcus</i> MÜLLER 116
turkensis	<i>Spongotorilispinus turkensis</i> KOZUR, MOIX & MOSTLER 26
tuvalica	<i>Podobursa tuvalica</i> KOZUR, MOIX & OZSVÁRT 24
	<i>Stampfiella tuvalica</i> KOZUR, MOIX & OZSVÁRT 26
tuvalicus	<i>Spongotorilispinus tuvalicus</i> MOIX & MOSTLER 26
umbilicosiformis	<i>Latrunculus (Peridipsacus) eburnoides umbilicosiformis</i> TELELDI ROTH 86
umbonata	<i>Chama gryphina umbonata</i> CSEPREGHY-MEZNERICS 100
undulatocollaris	<i>Lacuna umbonata</i> SZÓTS 54
upponyensis	<i>Voluta suturalis undulatocollaris</i> NOSZKY 93
uromensis	<i>Phyllosmilia upponyensis</i> KOLOSVÁRY 38
vaceki	<i>Archidiskodon meridionalis türömensis</i> VÖRÖS 138
vadaszi	<i>Ludwigia vaceki</i> GÉCZY 35
	<i>Lytoceras vaceki</i> GÉCZY 36
valdecarinata	<i>Balanus tintinnabulum vadászi</i> KOLOSVÁRY 112
varhegyensis	<i>Marginella (Volvarina) vadászi</i> BÁLDI 86
variabilis	<i>Panopaea (Pleuromya) vadászi</i> NOSZKY 78
variocostata	<i>Pleramplexus vadászi</i> KOLOSVÁRY 18
varpalotensis	<i>Stephanosmilia vadászi</i> KOLOSVÁRY 48
vatonnei	? <i>Turbinoberis vadászi</i> KOLOSVÁRY 49
vendli	<i>Raphitoma valdecarinata</i> BÁLDI 89
ventricosa	<i>Spatagoides (Toxopatagus) várhegyensis</i> SZÖRÉNYI 67
venusta	<i>Rhinolophus variabilis</i> TOPÁL 157
venzoi	<i>Bayania variocostata</i> SZÓTS 52
vertesensis	<i>Adeorbis várpalotensis</i> SZALAI 104
viai	<i>Cypricardia vatonnei</i> COQUAND 32
viallii	<i>Perna lamarcki vendli</i> NOSZKY 79
vighi	<i>Pleurotoma flexicostata ventricosa</i> NOSZKY 88
villei	<i>Uvigerina venusta</i> FRANZENAU 98
vindobonensis	<i>Murex (Pteronotus) detritus venzoi</i> NOSZKY 87
visegradensis	<i>Arca vértesensis</i> SZÓTS 49
vitalisi	<i>Cantharus vértesensis</i> SZÓTS 52
vogli	<i>Collonia vértesensis</i> SZÓTS 53
vonneguti	<i>Cylichna vértesensis</i> SZÓTS 54
waageni	<i>Cythara vértesensis</i> SZÓTS 54
	<i>Marginella vértesensis</i> SZÓTS 55
	<i>Seila vértesensis</i> SZÓTS 58
	<i>Terebellum vértesensis</i> SZÓTS 58
	<i>Kerepesia viai</i> MÜLLER 115
	<i>Costileioceras sinon viallii</i> GÉCZY 34
	<i>Clavella vighi</i> NOSZKY 82
	<i>Lima (Plagiostoma) villei</i> COQUAND 33
	<i>Lathyrus crassus vindobonensis</i> CSEPREGHY-MEZNERICS 106
	<i>Murex (Muricantha) sedgwicki vindobonensis</i> CSEPREGHY-MEZNERICS 107
	<i>Olivella (Lamprodoma) clavula vindobonensis</i> CSEPREGHY-MEZNERICS 108
	<i>Solenocurtus antiquatus vindobonensis</i> CSEPREGHY-MEZNERICS 103
	<i>Schizophrrys visegradensis</i> MÜLLER 119
	<i>Actaeon vitálisi</i> SZÓTS 51
	<i>Melanopsis vitálisi</i> STRAUSZ 107
	<i>Avicula hirundo vogli</i> NOSZKY 72
	<i>Lucina vogli</i> SZÓTS 50
	<i>Kahlerosphaera vonneguti</i> KOZUR, MOIX & OZSVÁRT 23
	<i>Nautilus (Temnocheilus) waageni</i> LÓCZY 19

- waageni *Reticularia waageni* LÓCZY 20
wagneri *Volutilithes (Neoathleta) affinis wagneri* NOSZKY 93
weemsi *Huglusphaera weemsi* KOZUR, MOIX & OZSVÁRT 21
wenzi *Fusus wenzi* NOSZKY 85
wronai *Panopeus wronai* MÜLLER 116
wuesti *Myotis wüsti* KORMOS 151
xyphodon *Lamna xyphodon* NEUGEBOREN 125
yini *Huglusphaera yini* KOZUR, MOIX & OZSVÁRT 22
yunnanensis *Productus yünنانensis* LÓCZY 20
zaboul *Rhynchonella zaboul* COQUAND 37
zakharovi *Huglusphaera zakharovi* KOZUR, MOIX & OZSVÁRT 22
zalaensis *Micromelania zalaensis* STRAUSZ 107
zamma *Lucina zamma* COQUAND 38
zelenkai *Limaria (Limatulella) langhiana zelenkai* BÁLDI 76
zimanyii *Amussium bronni zimányii* NOSZKY 71
zimmermanni *Capra (Capra) zimmermanni* KRETZOI 139
zircensis *Nummulites zircensis* KECSKEMÉTI 47
zitteli *Zebina zitteli* SZŐTS 60
zsivnyi *Aequipecten scabrellus zsivnyi* NOSZKY 71
Gymnosarda zsivnyi WEILER 124

References

- ALMÉRAS, Y. 1996. Les brachiopodes toarciens et aaléniens inférieurs du Bassin du Rhône. Paléontologie et biostratigraphie. Révision de la collection Dumortier et compléments. *Documents des Laboratoires de Géologie Lyon*, 138: 3-123. [Alm96]
- BÁLDI, T. 1960. Tortonische Molluskenfauna von „Badener Tegelfazies“ aus Szokolya, Nordungarn. *Annales Historico-Naturales Musei Nationalis Hungarici*, 52: 51-99. [Bál60]
- BÁLDI, T. 1962. *Glycymeris s. str.* des europäischen Oligozäns und Miozäns. *Annales Historico-Naturales Musei Nationalis Hungarici*, 54: 85-153. [Bál62]
- BÁLDI, T. 1963. Die oberoligozäne Molluskenfauna von Törökbálint. *Annales Historico-Naturales Musei Nationalis Hungarici*, 55: 71-107. [Bál63]
- BÁLDI, T. 1966. Die oberoligozäne Molluskenfauna von Eger und die Neuuntersuchung der Schichtfolge. *Annales Historico-Naturales Musei Nationalis Hungarici*, 58: 69-101. [Bál66]
- BÁLDI, T. 1973. *Mollusc Fauna of the Hungarian Upper Oligocene (Egerian). Studies in Stratigraphy, Palaeoecology, Palaeogeography and Systematics*. Akadémiai Kiadó, Budapest, 511 p. [Bál73]
- BÁLDI, T. 1986. *Mid-Tertiary Stratigraphy and Paleogeographic Evolution of Hungary*. Akadémiai Kiadó, Budapest, 201 p. [Bál86]
- BÁLDI, T., CSEPREGHY-MEZNERICS, I. & NYÍRÖ, M. R. 1964. La biostratigraphie des gisements oligocènes et miocènes de l'Est de la Montagne Börzsöny. *Annales Historico-Naturales Musei Nationalis Hungarici*, 56: 153-183. [Bál64]
- BÁLDI, T., KECSKEMÉTI, T., NYÍRÖ, M. R. & DROGER, C. W. 1961. Neue Angaben zur Grenzziehung zwischen Chatt und Aquitan in der Umgebung von Eger (Nordungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 53: 67-132. [Bál61]
- BÁLDI, T., LESS, G. & MANDIC, O. 1999. Some new aspects of the lower boundary of the Egerian stage (Oligocene, chronostratigraphic scale of the Paratethys area). *Abhandlungen der Geologischen Bundesanstalt*, 56(2): 653-668. [Bál99]
- BEURLEN, K. 1939. Neue Decapoden-Krebse aus dem ungarischen Tertiär. *Palaeontologische Zeitschrift*, 21: 135-160. [Beu39]
- BITNER, M. A. & DULAI, A. 2004. Revision of Miocene brachiopods of the Hungarian Natural History Museum, with special regard to the Meznerics collection. *Fragmента Palaeontologica Hungarica*, 22: 69-82. [Bit04]
- BITNER, M. A. & DULAI, A. 2008. Eocene micromorphic brachiopods from north-western Hungary. *Geologica Carpathica*, 59(1): 31-43. [Bit08]
- BITTNER, A. 1893. Decapoden des pannonicischen Tertiärs. *Sitzungsberichte der Mathematisch-Naturwissenschaftliche Classe der Kaiserlichen Akademie der Wissenschaften*, 102(1): 10-37. [Btt93]
- BODA, J. 1964. Magyarországi ósmaradványtípusok jegyzéke. *Ősállatok (Catalogus originalium fossilium Hungariae. Pars zoologica)*. Magyar Állami Földtani Intézet (Hungarian Geological Institute), Budapest, 229 p.
- BÖHM, B. 1941. Fosszilis halmaradványok az erdélyi Kovászna és Kommandó környékéről (Die Fossilien Fische von Kovászna und Kommandó in Siebenbürgen). *Magyar Királyi Földtani Intézet Évkönyve (Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt)*, 35(5): 179-203. [Böh41]
- BÖHM, B. 1942. Adatok a magyarországi harmadkori halfaunához (Beiträge zur tertiären Fischfauna Ungarns). *Geologica Hungarica, Series Palaeontologica*, 19: 1-42. [Böh42]
- BRAGA, G. & GRÜNING, A. 1975. Foraminiferi bentonici dell'Eocene superiore. *Schweizerische Paläontologische Abhandlungen*, 97: 98-111. [Bra75]
- BRÜHL, C. B. 1860. *Phoca holtschensis*, der fossile Phocafuss des Pester Universitäts-Museums, ein Unicum. *Mittheilungen aus dem Kaiserlich-Königlichen Zoologischen Institute der Universität Pest*, 2: 1-17. [Brü60]
- CICHA, I., RÖGL, F., RUPP, C. & ČTYROKA, J. 1998. Oligocene-Miocene Foraminifera of the Central Paratethys. *Abhandlungen der Seckenbergischen Naturforschenden Gesellschaft*, 549: 1-325. [Cic98]

- COMPANY, M., FŐZY, I., SANDOVAL, J. & TAVERA, J. M. 2006. *Deitanites* n. g. and other related ammonites. Their significance within the family Holcodiscidae (Lower Cretaceous, Mediterranean region). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1: 1-14. [Com06]
- COQUAND, H. 1841. Mémoire sur les *Aptychus*. *Bulletin de la Société Géologique de France*, 12: 376-391. [Coq41]
- COQUAND, H. 1860. *Synopsis des animaux et des végétaux fossiles observés dans les formations secondaires de la Charente, de la Charente-Inférieure et de la Dordogne*. Barlatier-Feissat et Demonchy, Marseille, 146 p. [Coq60]
- COQUAND, H. 1862. *Géologie et paléontologie de la région sud de la province de Constantine*. Société d'Émulation, Marseille, 341 p. [Coq62]
- COQUAND, H. 1880. Études supplémentaires sur la paléontologie algérienne faisant suite à la description géologique et paléontologique de la région sud de la province de Constantine. *Bulletin de l'Académie d'Hippone*, 15: 1-449. [Coq80]
- CUSHMAN, J. A. 1936. New genera and species of the families Verneuilinidae and Valvulinidae of the subfamily Virgulininae. *Cushman Laboratory for Foraminiferal Research, Special Publication*, 6: 1-71. [Cus36]
- CUVILLIER, J. & SZAKALL, V. 1949. Foraminifères d'Aquitaine. Première partie (Reophacidae à Nonionidae). *Société Nationale des Pétroles d'Aquitaine, Toulouse*: 1-112. [Cuv49]
- Cs. MEZNERICS, I. 1950. Néhány eddig ismeretlen és új forma a K-Cserhát tortonai rétegeiből [Some previously unknown and new forms from the Tortonian strata of eastern Cserhát]. *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 80(10-12): 395-404. [CsM50]
- CsÁSZÁR, G. (ed.) 1997. *Basic Litostratigraphic Units of Hungary. Charts and short descriptions (Magyarország litosztratigráfiai alapegységei. Táblázatok és rövid leírások)*. Geological Institute of Hungary, Budapest, 114 p.
- CSEPREGHY-MEZNERICS, I. 1950. A hidasi (Baranya m.) tortonai fauna (Die Tortonische Fauna von Hidas (Kom. Baranya, Ungarn)). *Magyar Állami Földtani Intézet Évkönyve (Annals of the Hungarian Geological Institute)*, 39(2): 1-115. [C-M50]
- CSEPREGHY-MEZNERICS, I. 1951. A salgótarjánvidéki slir és pectenes homokkő faunája [The fauna of schlier and *Pecten*-bearing sandstone around Salgótarján]. *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 81(7-9): 303-319. [C-M51]
- CSEPREGHY-MEZNERICS, I. 1952. Paläontologische Seltenheiten in der Fauna von Szob (Őslénytani ritkaságok a szobi faunából). *Annales Historico-Naturales Musei Nationalis Hungarici (Series Nova)*, 2: 225-231. [C-M52]
- CSEPREGHY-MEZNERICS, I. 1953. Mittelmiozäne Pleurotomen aus Ungarn (Magyarországi középmiocén Pleurotomák). *Annales Historico-Naturales Musei Nationalis Hungarici (Series Nova)*, 4: 5-22. [C-M53]
- CSEPREGHY-MEZNERICS, I. 1954. A keletcserháti helvétai és tortónai fauna (Helvetische und Tortonische Fauna aus dem östlichen Cserhátgebirge). *Magyar Állami Földtani Intézet Évkönyve (Annals of the Hungarian Geological Institute)*, 41(4): 1-185. [C-M54]
- CSEPREGHY-MEZNERICS, I. 1956. A szobi és letkési puhatestű fauna (Die Molluskenfauna von Szob und Letkés). *Magyar Állami Földtani Intézet Évkönyve (Annals of the Hungarian Geological Institute)*, 45(2): 363-477. [C-M56]
- CSEPREGHY-MEZNERICS, I. 1960. Pectinidés du Néogène de la Hongrie et leur importance stratigraphique. *Mémoires de la Société Géologique de France (Nouvelle Série)*, 92: 1-56. [C-M60]
- CSEPREGHY-MEZNERICS, I. 1961. Quelques Lamellibranches rares du Miocène de la Hongrie. *Annales Historico-Naturales Musei Nationalis Hungarici*, 53: 133-137. [C-M61]
- CSEPREGHY-MEZNERICS, I. 1969. Nouvelles Gastropodes et Lamellibranches pour la faune hongroise des gisements tortoniens-inférieurs de la Montagne de Bükk. *Annales Historico-Naturales Musei Nationalis Hungarici*, 61: 63-127. [C-M69]
- DARAKCHIEVA, S. 1999. *Small Tertiary Foraminifers from Bulgaria. Atlas*. Sofia, 92 p. [Dar99]
- DULAI, A. 2001. Middle Miocene (Badenian) Polyplacophora (Mollusca) remains from borehole Szokolya-2 (Börzsöny Mts, Hungary, Central Paratethys). *Fragmenta Palaeontologica Hungarica*, 19: 39-49. [Dul01]

- DULAI, A., BITNER, M. A. & MÜLLER, P. 2008. A monospecific assemblage of a new rhynchonellide brachiopod from the Paleocene of Austria. *Fossils and Strata*, 54: 193–201. [Dul08]
- ÉHIK, J. 1930. *Prodinotherium hungaricum* nov. gen., nov. spec. *Geologica Hungarica, Series Palaeontologica*, 6: 1–18. [Éhi30]
- ERDŐS, L. 1900. Új *Pyrula* faj Pomáz fiatalabb harmadkori üledékeiből (Eine neue *Pyrula*-Species aus den jüngeren Tertiärschichten von Pomáz). *Földtani Közlöny (Geologische Mittheilungen)*, 30(10–12): 262–266, 296–301. [Erd00]
- FILIPESCU, S. 2001. Cenozoic lithostratigraphic units in Transylvania. In: BUCUR, I. I., FILIPESCU, S. & SÁSÁRAN, E. (eds.), *Algae and carbonate platforms in western part of Romania. 4th Regional Meeting of IFAA Cluj-Napoca 2001 – Field Trip Guidebook*. Cluj University Press, Cluj-Napoca, pp. 75–91.
- FOSTOWICZ-FRELIK, Ł. 2007. Revision of *Hypolagus* (Mammalia: Lagomorpha) from the Plio-Pleistocene of Poland: qualitative and quantitative study. *Annales Zoologici*, 57(3): 541–590. [F-F07]
- FOSTOWICZ-FRELIK, Ł. & GASPARIK, M. 2006. The taxonomic status of leporid remains from Ördög-lyuk Cave, Solymár (Hungary). *Acta Zoologica Cracoviensia*, 49A(1–2): 151–161. [F-F06]
- FŐZY, I. 2001. Campanian (Late Cretaceous) cephalopods from Sümeg (Transdanubian Central Range, Hungary). *Fragmenta Palaeontologica Hungarica*, 19: 25–37. [Főz01]
- FRANZENAU, Á. 1881. Adatok a rákosi (Budapest) felső mediterrán emelet foraminifera faunájához (Beitrag zur Foraminiferen Fauna der Rákoser (Budapest) Ober-Mediterran Stufe). *Földtani Közlöny (Geologische Mittheilungen)*, 11(1–3): 31–55, 83–107. [Fra81]
- FRANZENAU, Á. 1892. A romhányi tályag (Der Tegel von Romhány). *Természettrajzi Füzetek*, 15(3): 107–113, 138–143. [Fra92]
- FRANZENAU, Á. 1894a. Adatok Letkés faunájához [Data to the fauna of Letkés]. *Mathematikai és Természettudományi Közlemények (Mathematische und Naturwissenschaftliche Berichte aus Ungarn)*, 26(1): 1–36. [Fra94a]
- FRANZENAU, Á. 1894b. Fossile Foraminiferen von Markuševci, in Kroatien. *Glasnik Hrvatskoga Naravoslovnoga Društva*, 6: 249–291. [Fra94b]
- FRANZENAU, Á. & MAJZON, L. 1956. New and interesting Foraminifera species. *Annales Historico-Naturales Musei Nationalis Hungarici (Series Nova)*, 7: 211–220. [Fra56]
- FRECH, F. 1897. Palaeozoos korállok. In: Gróf Széchenyi Béla keletázsiai utjának tudományos eredménye, vol. 3, pt. 1D. Kilián, Budapest, pp. 193–200. [Fre97]
- FRECH, F. 1899. Palaeozoische Korallen. In: Wissenschaftliche Ergebnisse der Reise des Grafen Béla Széchenyi in Ostasien, vol. 3, pt. 1D. Hölzel, Wien, pp. 229–236. [Fre99]
- GAÁL, I. 1938a. Az egrikkel azonos harmadkori puhatestűek Balassa-Gyarmaton és az oligocén-kérdés (Über die mit der Egerer gleichalterige tertiäre Molluskenfauna von Balassa-Gyarmat und das Oligozän-Problem). *Annales Historico-Naturales Musei Nationalis Hungarici*, 31: 1–87. [Gaá38a]
- GAÁL, I. 1938b. A keleti orrszarvú – *Dicerorhinus orientalis* Schloss. – új alakjának csontmaradványai Magyarországon [Bone remains of a new form of the oriental rhinoceros (*Dicerorhinus orientalis* Schloss.) from Hungary]. *Természettudományi Közlöny Pótfüze*, 1938(10–12): 130–138. [Gaá38b]
- GAÁL, I. 1943. Alsó-pliocén emlősmaradványok Hatvanból (Unterpliozäne Säugetierreste aus Hatvan in Ungarn). *Geologica Hungarica, Series Palaeontologica*, 20: 1–120. [Gaá43]
- GASPARIK, M. 1993. Deinotheres (Proboscidea, Mammalia) of Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici*, 85: 3–17. [Gas93]
- GASPARIK, M. 2001. Neogene proboscidean remains from Hungary; an overview. *Fragmenta Palaeontologica Hungarica*, 19: 61–77. [Gas01]
- GASPARIK, M. 2007. “Elephants” in the cellar. A revision of the Neogene proboscidean remains, damaged in the fire of the Hungarian Natural History Museum in 1956. *Fragmenta Palaeontologica Hungarica*, 24–25: 83–91. [Gas07]
- GÉCZY, B. 1966. Ammonoides jurassiques de Csernye, Montagne Bakony, Hongrie. Part I. (Hammatoceratidae). *Geologica Hungarica, Series Palaeontologica*, 34: 1–276. [Géc66]
- GÉCZY, B. 1967. Ammonoides jurassiques de Csernye, Montagne Bakony, Hongrie. Part II. (excl. Hammatoceratidae). *Geologica Hungarica, Series Palaeontologica*, 35: 1–413. [Géc67]
- GORJANOVIĆ-KRAMBERGER, K. 1902a. Palaeo-ichthyologai adalékok. *Magyar Királyi Földtani Intézet Évkönyve*, 14(1): 1–20. [Gor02a]

- GORJANOVIĆ-KRAMBERGER, K. 1902b. Palaeoichthyologische Beiträge. *Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt*, 14(1): 1–21. [Gor02b]
- GÖRÖC, Á. 1992. Sarmatian foraminifera of the Zsámbék Basin, Hungary. *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Geologica*, 29: 31–153. [Gör92]
- GÖRÖC, Á. 1994. Early Jurassic planktonic foraminifera from Hungary. *Micropaleontology*, 40(3): 255–260. [Gör94]
- GRADSTEIN, F., OGG, J. & SMITH, A. (eds.) 2004. *A Geologic Time Scale 2004*. Cambridge University Press, Cambridge, 589 p.
- HABLY, L., ERDEI, B. & KVAČEK, Z. 2001. 19th century's palaeobotanical types and originals of the Hungarian Natural History Museum. *Studia Naturalia*, 13: 1–235.
- HABLY, L. & SZAKÁLY, M. 1989. *The Catalogue of Leaf-Fossil Types Preserved in Hungary*. Akadémiai Kiadó, Budapest, 253 p.
- HAGN, H. 1956. Geologische und paläontologische Untersuchungen im Tertiär des Monte Brione und seiner Umgebung (Gardasee, Ober-Italien). *Palaeontographica, Abteilung A*, 107: 67–210. [Hag56]
- HAGN, H. 1960. Die stratigraphischen, paläogeographischen und tektonischen Beziehungen zwischen Molasse und Helvetikum im östlichen Oberbayern. *Geologica Bavarica*, 44: 1–208. [Hag60]
- HANTKEN, M. 1868. A kis-czelli tállyag foraminiferái [Foraminifers of the Kiscell Clay]. *Magyar honi Földtani Társulat Munkálatai*, 4: 75–96. [Han68]
- HANTKEN, M. 1875a. Die Fauna der *Clavulina Szabói* Schichten. I. Theil. Foraminiferen. *Mittheilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt*, 4(1): 1–93. [Han75a]
- HANTKEN, M. 1875b. A *Clavulina Szabói* rétegek faunája. I. rész. Foraminiferák. *Magyar Királyi Földtani Intézet Évkönyve*, 4(1): 1–82. [Han75b]
- HANTKEN, M. 1883a. Die *Clavulina Szabói*-Schichten im Gebiete der Euganeen und der Meeralpen und die cretacische Scaglia in den Euganeen. *Mathematische und Naturwissenschaftliche Berichte aus Ungarn*, 2: 121–169. [Han83a]
- HANTKEN, M. 1883b. A *Clavulina Szabói*-rétegek, az Euganeák és a tengeri Alpok területén és a kréta-korú „Scaglia” az Euganeákban. *Értekezések a Természettudományok Köréből*, 13: 1–47. [Han83b]
- HARRIS, J. M. 1973. *Prodeinotherium* from Gebel Zelten, Libya. *Bulletin of the British Museum (Natural History), Geology*, 23(5): 285–348. [Har73]
- HELLER, F. 1930. Eine Forest-Bed-Fauna aus der Sackdillinger Höhle (Oberpfalz). *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Abhandlungen, Abteilung B*, 63: 247–298. [Hel30]
- HEMMER, H. 2001. Die Feliden aus dem Epivilafranchium von Untermaßfeld. In: KAHLKE, R.-D. (ed.), *Das Pleistozän von Untermaßfeld bei Meiningen (Thüringen). Teil 3. Monographien des Römisch-Germanischen Zentralmuseums Mainz*, 40(3): 699–782. [Hem01]
- HÍR, J. 1994. *Cricetinus beremendensis* sp. n. (Rodentia, Mammalia) from the Pliocene fauna of Berekend 15. (S Hungary). *Fragmenta Mineralogica et Palaeontologica*, 17: 71–89. [Hir94]
- HÍR, J. 1996. *Cricetinus janossyi* sp. n. (Rodentia, Mammalia) from the Pliocene fauna of Osztramos 7. (N Hungary). *Fragmenta Mineralogica et Palaeontologica*, 18: 79–90. [Hir96]
- HÍR, J. 1997. *Cricetus runtonensis solymarensis* ssp. n. (Mammalia, Rodentia) from the Late Middle Pleistocene fauna of Solymár. *Annales Historico-Naturales Musei Nationalis Hungarici*, 89: 23–42. [Hir97]
- HORVÁTH, M. 2002. Data to revision and distribution of small foraminifera species described by Hantken (1868, 1875). Part I. Textulariidae and Miliolidae. *Fragmenta Palaeontologica Hungarica*, 20: 25–42. [Hor02]
- HORVÁTH, M. 2003. Data to revision and distribution of small foraminifera species described by Hantken (1868, 1875). Part II. Nodosariidae and Vaginulinidae. *Fragmenta Palaeontologica Hungarica*, 21: 5–32. [Hor03]
- HOTTINGER, L. 1977. Foraminifères operculiniformes. *Mémoires du Muséum National d'Histoire Naturelle, Série C, Sciences de la Terre*, 40: 1–159. [Hot77]
- HUENE, F. 1932. Die fossile Reptil-Ordnung Saurischia, ihre Entwicklung und Geschichte. *Monographien zur Geologie und Palaeontologie, Serie I*, 4: 1–361. [Hue32]
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE 1999. *International Code of Zoological Nomenclature*. The International Trust for Zoological Nomenclature, London, 306 p.

- JÁNOSSY, D. 1962. Vorläufige Mitteilung über die Mittelpaläozäne Vertebratenfauna der Tarkö-Felsnische (NO-Ungarn, Bükk-Gebirge). *Annales Historico-Naturales Musei Nationalis Hungarici*, 54: 155–176. [Ján62]
- JÁNOSSY, D. 1965. Nachweis einer jungmittelpleistozänen Kleinvertebratenfauna aus der Felsnische Uppony I. (Nordungarn). *Karszt- és Barlangkutatás*, 4: 55–68. [Ján65]
- JÁNOSSY, D. 1969a. Uj Eomyida (Rodentia, Mammalia) a bódvaszilasi osztramosi köfejtő 3. lelet-helyének alsópleisztocén faunájából (A new eomyid (Rodentia, Mammalia) from the lowest Pleistocene of Hungary). *Óslénytani Viták (Discussiones Palaeontologicae)*, 13: 5–40. [Ján69a]
- JÁNOSSY, D. 1969b. Stratigraphische Auswertung der europäischen mittelpaläozänen Wirbeltierfauna. Teil II. *Berichte der Deutschen Gesellschaft für Geologische Wissenschaften. Reihe A, Geologie und Paläontologie*, 14(5): 573–643. [Ján69b]
- JÁNOSSY, D. 1970. Ein neuer Eomyide (Rodentia, Mammalia) aus dem Ältestenpaläozän („Oberes Villafrankium“, Villányium) des Osztramos (Nordostungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 62: 99–113. [Ján70]
- JÁNOSSY, D. 1972a. Middle Pliocene microvertebrate fauna from the Osztramos Loc. 1. (Northern Hungary). *Annales Historico-Naturales Musei Nationalis Hungarici*, 64: 27–52. [Ján72a]
- JÁNOSSY, D. 1972b. Ein kleiner *Hystrix* aus Altpaläozän der Fundstelle Osztramos 8. (Nordungarn). *Vertebrata Hungarica*, 13: 163–182. [Ján72b]
- JÁNOSSY, D. 1972c. Die Mittelpaläozäne Vogelfauna der Stránská Skála. *Anthropos*, 20: 35–64. [Ján72c]
- JÁNOSSY, D. 1973. New species of *Episoriculus* from the Middle Pliocene of Osztramos (North Hungary). *Annales Historico-Naturales Musei Nationalis Hungarici*, 65: 49–55. [Ján73]
- JÁNOSSY, D. 1974a. New “Middle Pliocene” microvertebrate fauna from Northern Hungary (Osztramos Loc. 9). *Fragmenta Mineralogica et Palaeontologica*, 5: 17–27. [Ján74a]
- JÁNOSSY, D. 1974b. Die mittelpaläozäne Vogelfauna von Hundsheim (Niederösterreich). *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abteilung I*, 182(6–8): 211–257. [Ján74b]
- JÁNOSSY, D. 1979. Plio-Pleistocene bird remains from the Carpathian Basin. IV. Anseriformes, Gruiformes, Charadriiformes, Passeriformes. *Aquila*, 85: 11–39. [Ján79]
- JÁNOSSY, D. 1986. *Pleistocene Vertebrate Faunas of Hungary*. Akadémiai Kiadó, Budapest, 208 p.
- JÁNOSSY, D. 1991. Late Miocene bird remains from Polgárdi (W-Hungary). *Aquila*, 98: 13–35. [Ján91]
- JÁNOSSY, D. 1992. Lower Pleistocene bird remains from Beremend (S-Hungary, Loc. 15. and 16.). *Aquila*, 99: 9–25. [Ján92]
- JÁNOSSY, D. & VAN DER MEULEN, A. J. 1975. On *Mimomys* (Rodentia) from Osztramos-3, North Hungary. *Koninklijke Nederlandse Akademie van Wetenschappen, Series B*, 78(5): 381–391. [Ján75]
- JASKÓ, S. 1940. A Rima és Tarna közének oligocén rétegei és kövületei (Die Versteinerungen der Oligozienschichten zwischen den Flüssen Rima und Tarna (Nordungarn)). *Földtani Közlöny (Geologische Mitteilungen)*, 70(10–12): 294–317, 369–373. [Jas40]
- KADIĆ, O. 1907a. *Mesocetus hungaricus* Kadić. Egy új balaenopteridafaj a borbolyai miocén rétegekből. *Magyar Királyi Földtani Intézet Évkönyve*, 16(2): 21–86. [Kad07a]
- KADIĆ, O. 1907b. *Mesocetus hungaricus* Kadić, eine neue Balaenopteridenart aus dem Miozän von Borbolya in Ungarn. *Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt*, 16(2): 23–91. [Kad07b]
- KAZÁR, E. 2005. A new kentriodontid (Cetacea: Delphinoidea) from the Middle Miocene of Hungary. *Mitteilungen aus dem Museum für Naturkunde in Berlin, Geowissenschaftliche Reihe*, 8: 53–73. [Kaz05]
- KAZÁR, E. 2006. *Sophianaecetus*, a replacement name for *Mediocris* (Cetacea: Delphinoidea: Kentriodontidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin, Fossil Record*, 9(2): 260. [Kaz06]
- KECSKEMÉTI, T. 1959. Die Discocycliniden des südlichen Bakonygebirges. *Annales Historico-Naturales Musei Nationalis Hungarici*, 51: 31–84. [Kec59]
- KECSKEMÉTI, T. 1970. Neue Nummuliten-Arten aus dem Bakonygebirge (Transdanubien, Ungarn). I. Teil. *Annales Historico-Naturales Musei Nationalis Hungarici*, 62: 53–65. [Kec70]
- KECSKEMÉTI, T. 1974. Neue Nummuliten-Arten aus dem Bakonygebirge (Transdanubien, Ungarn). II. Teil. *Annales Historico-Naturales Musei Nationalis Hungarici*, 66: 33–46. [Kec74]

- KECSKEMÉTI, T. & NAGY, I. Z. 1987. Palaeontological collections of the Hungarian Natural History Museum (Budapest). *Annals of the History of Hungarian Geology, Special Issue*, 1: 151–155.
- KECSKEMÉTI, T. & VAŇOVÁ, M. 1972. Nummulites of the Dorog–Štúrovo basin. *Zborník Geologických Vied, Západné Karpaty*, 17: 105–145. [Kec72]
- KENAWY, A. I. & NYÍRÓ, R. M. 1967. Zwei neue Foraminiferen aus dem Oberoligozän in Eger (Nordungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 59: 103–107. [Ken67]
- KOCH, A. 1904a. Kövült czápafogak és emlősmaradványok Felsőesztergályról, Nógrád vármegyében (Fossile Haifischzähne und Säugetierreste von Felsőesztergály, im Komitate Nógrád). *Földtani Közlöny (Geologische Mitteilungen)*, 34(5–7): 190–202, 260–273. [Koc04a]
- KOCH, A. 1904b. A beocsini cementmárga kövült halai (Die fossilen Fische des Beocsiner Cementmergels). *Annales Historico-Naturales Musei Nationalis Hungarici*, 2: 1–72. [Koc04b]
- KÓKAY, J. 1967. A Bakony-hegység felsőtortonai képződményei (Obertortonische Ablagerungen des Bakonygebirges). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 97(1): 74–90. [Kök67]
- KOKEN, E. 1913. Beiträge zur Kenntnis der Schichten von Heiligenkreuz (Abteital, Südtirol). *Abhandlungen der Kaiserlich-Königlichen Geologischen Reichsanstalt*, 16(4): 1–43. [Kok13]
- KOLOSVÁRY, G. 1941a. *Balanus hungaricus* n. sp. *Földtani Közlöny (Geologische Mitteilungen)*, 71(7–12): 282–284. [Kol41a]
- KOLOSVÁRY, G. 1941b. Ein neuer Ophiurites von Kiscell (Ungarn). *Palaeontologische Zeitschrift*, 22(3–4): 307–309. [Kol41b]
- KOLOSVÁRY, G. 1942. Über tertiäre Balaniden Ungarns. II. *Palaeontologische Zeitschrift*, 23(1–2): 203–205. [Kol42]
- KOLOSVÁRY, G. 1947. Eine neue Balanide aus dem ungarischen Eozän. *Annales Historico-Naturales Musei Nationalis Hungarici*, 40(8): 305–307. [Kol47]
- KOLOSVÁRY, G. 1948. Helvéti emeletbeli új balanidák Várpalotáról (New balanids from the Middle Miocen of Várpalota in Hungary). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 78: 102–112. [Kol48]
- KOLOSVÁRY, G. 1949a. Új balanidák a hazai harmadkorból (New balanids from the Hungarian Tertiary age). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 79(1–4): 111–118. [Kol49a]
- KOLOSVÁRY, G. 1949b. Dunántúli eocén-korallok (The Eocene corals of the Hungarian Transdanubian province). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 79(5–8): 141–242. [Kol49b]
- KOLOSVÁRY, G. 1950. Négy új Balanida a magyar harmadkorból (Descriptions of 3 new fossil Tertiary barnacles from Hungary). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 80(7–9): 271–276. [Kol50]
- KOLOSVÁRY, G. 1951a. Magyarország permo-karbon koralljai (The Permo-Carboniferous corals of Hungary). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 81(1–3, 4–6): 4–56, 171–185. [Kol51a]
- KOLOSVÁRY, G. 1951b. Szabadbattyáni alsó-karbon korallok (The Lower-Carboniferous corals from Hungary). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 81(4–6): 275–283. [Kol51b]
- KOLOSVÁRY, G. 1952. A stratigraphical study on some Tertiary balanids from Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici (Series Nova)*, 2: 233–236. [Kol52]
- KOLOSVÁRY, G. 1954a. Magyarország kréta-időszaki koralljai (Les Coralliaires du Crétacé de la Hongrie). *Magyar Állami Földtani Intézet Évkönyve (Annals of the Hungarian Geological Institute)*, 42(2): 67–163. [Kol54a]
- KOLOSVÁRY, G. 1954b. Adatok a magyarországi júra-időszaki korallok ismeretéhez (Beiträge zur Kenntnis der fossilen Korallen der Jurazeit in Ungarn). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 84(3): 235–243. [Kol54b]
- KORDOS, L. 1981. Some complements to the knowledge of a Middle Eocene Sirenia, *Sirenavus hungaricus* Kretzoi, 1941. *Fragmenta Mineralogica et Palaeontologica*, 10: 75–78. [Krd81]
- KORDOS, L. 1985. Lower Turolian (Neogene) *Anomalospalax* gen. n. from Hungary and its phylogenetic position. *Fragmenta Mineralogica et Palaeontologica*, 12: 27–42. [Krd85]
- KORDOS, L. 1987. *Karstocricetus skofleki* gen. n., sp. n. and the evolution of the Late Neogene Cricetidae in the Carpathian Basin. *Fragmenta Mineralogica et Palaeontologica*, 13: 65–88. [Krd87]

- KORETSKY, I. A. 2001. Morphology and systematics of Miocene Phocinae (Mammalia: Carnivora) from Paratethys and the North Atlantic region. *Geologica Hungarica, Series Palaeontologica*, 54: 1-109. [Kry01]
- KORMOS, T. 1926. *Amblycoptus oligodon* n. g. & n. sp. Új cickány-féle a magyarországi pliocénból (*Amblycoptus oligodon* n. g. & n. sp. Eine neue Spitzmaus aus dem ungarischen Pliozän). *Annales Historico-Naturales Musei Nationalis Hungarici*, 24: 352-391. [Krm26]
- KORMOS, T. 1930. Diagnosen neuer Säugetiere aus der oberpliozänen Fauna des Somlyóberges bei Püspökfürdő. *Annales Historico-Naturales Musei Nationalis Hungarici*, 27: 237-246. [Krm30]
- KORMOS, T. 1931. Über eine neuentdeckte Forestbed-Fauna in Dalmatien. *Palaeobiologica*, 4: 113-136. [Krm31]
- KORMOS, T. 1932a. Neue Wühlmäuse aus dem Oberpliocän von Püspökfürdő. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Abhandlungen, Abteilung B*, 69: 323-346. [Krm32a]
- KORMOS, T. 1932b. Die Füchse des ungarischen Oberpliozäns. *Folia Zoologica et Hydrobiologica*, 4(2): 167-188. [Krm32b]
- KORMOS, T. 1933a. Revision der präglazialen Wühlmäuse vom Gesprengberg bei Brassó in Siebenbürgen. *Paleontologische Zeitschrift*, 15(1): 1-21. [Krm33a]
- KORMOS, T. 1933b. *Baranomys lóczyi* n. g. n. sp., új rágcsáló a magyarországi felső pliocénből (*Baranomys lóczyi* n. g. n. sp. ein neues Nagetier aus dem Oberpliocän Ungarns). *Állattani Közlemények*, 30(1-2): 45-54. [Krm33b]
- KORMOS, T. 1934a. Felsőpliocénkori új rovarevők, denevérek és rágcsálók Villány környékéről (Neue Insektenfresser, Fledermäuse und Nager aus dem Oberpliozän der Villányer Gegend). *Földtani Közlöny (Geologische Mitteilungen)*, 64(10-12): 296-321. [Krm34a]
- KORMOS, T. 1934b. Neue und wenig bekannte Musteliden aus dem ungarischen Oberpliozän. *Folia Zoologica et Hydrobiologica*, 5(2): 129-158. [Krm34b]
- KORMOS, T. 1934c. Az euráziai nyulak származástani problémája (Zur Frage der Abstammung eurasiatischer Hasen). *Állattani Közlemények*, 31(1-2): 65-78. [Krm34c]
- KORMOS, T. 1934d. *Manis hungarica* n. sp., das erste Schuppentier aus dem europäischen Oberpliozän. *Folia Zoologica et Hydrobiologica*, 6(1): 87-94. [Krm34d]
- KORMOS, T. 1937. A hundsheimi fossilis kisemlősök revíziója (Revision der Kleinsäuger von Hundsheim in Niederösterreich). *Földtani Közlöny (Geologische Mitteilungen)*, 67(1-3, 4-6): 23-37, 157-171. [Krm37]
- KOZUR, H. W., MOIX, P. & OZSVÁRT, P. 2007a. Characteristic Nassellaria of the lower Tuvalian (Upper Triassic) *Spongotorilispinus moixi* Zone of the Hügülü Unit in the Mersin Mélange, south-eastern Turkey. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 90(3): 151-173. [Koz07a]
- KOZUR, H. W., MOIX, P. & OZSVÁRT, P. 2007b. Stratigraphically important Spumellaria and Entactinaria from the lower Tuvalian (Upper Triassic) of the Hügülü Unit in the Mersin Mélange, south-eastern Turkey. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 90(3): 175-195. [Koz07b]
- KOZUR, H. W., MOIX, P. & OZSVÁRT, P. 2007c. Further new Nassellaria of the lower Tuvalian (Upper Triassic) *Spongotorilispinus moixi* Zone of the Hügülü Unit in the Mersin Mélange. *Bulletin de la Société Vaudoise des Sciences Naturelles*, 90(4): 197-215. [Koz07c]
- KRENNER, J. S. 1867. Ajnácskő ősemlösei [Fossil mammals from Ajnácskő]. *Magyarhoni Földtani Társulat Munkálatai*, 3: 114-132. [Krn67]
- KRETZOI, M. 1938. Die Raubtiere von Gombaszög nebst einer Übersicht der Gesamtfauna (Ein Beitrag zur Stratigraphie des Altquartaers). *Annales Historico-Naturales Musei Nationalis Hungarici*, 31: 88-157. [Kre38]
- KRETZOI, M. 1940. Alttertiäre Perissodactylen aus Ungarn (Óharmadkori magyar Perissodactylák). *Annales Historico-Naturales Musei Nationalis Hungarici*, 33: 87-98. [Kre40]
- KRETZOI, M. 1941a. Weitere Beiträge zur Kenntnis der Fauna von Gombaszög (Újabb megfigyelések a gombaszögi faunáról). *Annales Historico-Naturales Musei Nationalis Hungarici*, 34: 105-139. [Kre41a]
- KRETZOI, M. 1941b. *Sirenavus hungaricus* n. g., n. sp., ein neuer Prorastomide aus dem Mitteleozän (Lutetium) von Felsőgalla in Ungarn (Új Prorastomida (*Sirenavus hungaricus* n. g., n. sp.) Felsőgalla középső eocénjéből). *Annales Historico-Naturales Musei Nationalis Hungarici*, 34: 146-156. [Kre41b]

- KRETZOI, M. 1941c. Ausländische Säugetierfossilien der ungarischen Museen (1-4.). *Földtani Közlöny (Geologische Mitteilungen)*, 71(1-6): 170-176. [Kre41c]
- KRETZOI, M. 1941d. Ösemlősmaradványok Bettfiáról (Die unterpleistozäne Säugetierfauna von Bettia bei Nagyvárad). *Földtani Közlöny (Geologische Mitteilungen)*, 71(7-12): 235-261, 308-335. [Kre41d]
- KRETZOI, M. 1941e. Szarmatakori antilop Sopronból (Neue Antilopen-Form aus dem soproner Sarmat). *Földtani Közlöny (Geologische Mitteilungen)*, 71(7-12): 261-268, 336-343. [Kre41e]
- KRETZOI, M. 1941f. A magyar mammut (*Mammonteus hungaricus* n. sp.) (*Mammonteus hungaricus*, ein neues Waldmammut aus Ungarn). *Földtani Közlöny (Geologische Mitteilungen)*, 71(7-12): 268-270, 343-345. [Kre41f]
- KRETZOI, M. 1941g. Fóka-maradványok az érdi szarmatából (Seehund-Reste aus dem Sarmat von Érd bei Budapest). *Földtani Közlöny (Geologische Mitteilungen)*, 71(7-12): 274-279, 350-356. [Kre41g]
- KRETZOI, M. 1942a. *Necroteuthis* n. g. (Ceph. Dibr., Necroteuthidae n. f.) aus dem Oligozän von Budapest und das System der Dibranchiata (*Necroteuthis* n. g. a kiscelli oligocénból). *Földtani Közlöny (Geologische Mitteilungen)*, 72(1-3): 99-100, 124-138. [Kre42a]
- KRETZOI, M. 1942b. Ausländische Säugetierfossilien der ungarischen Museen (5-6.). *Földtani Közlöny (Geologische Mitteilungen)*, 72(1-3): 139-148. [Kre42b]
- KRETZOI, M. 1942c. A tigrisgörény, görény és nyérc a magyar pleisztocénben (Tigeriltis, Iltis und Nerz im ungarischen Pleistozän). *Földtani Közlöny (Geologische Mitteilungen)*, 72(4-12): 237-255, 323-344. [Kre42c]
- KRETZOI, M. 1942d. Két új Agriotheriida a magyar pannonból (Zwei neue Agriotheriiden aus dem ungarischen Pannon). *Földtani Közlöny (Geologische Mitteilungen)*, 72(4-12): 257-259, 350-353. [Kre42d]
- KRETZOI, M. 1942e. Kecskek a magyar diluviumban (*Capra* im ungarischen Diluvium). *Földtani Közlöny (Geologische Mitteilungen)*, 72(4-12): 259-262, 353-356. [Kre42e]
- KRETZOI, M. 1942f. Präökupierte und durch ältere zu ersetzenende Säugetiernamen. *Földtani Közlöny (Geologische Mitteilungen)*, 72(4-12): 345-349. [Kre42f]
- KRETZOI, M. 1943a. Ein neuer Muscardinide aus dem ungarischen Miozän (Új pele a magyar miocénból). *Földtani Közlöny (Geologische Mitteilungen)*, 73(1-3): 182, 271-273. [Kre43a]
- KRETZOI, M. 1943b. Bemerkungen über *Petényia*. *Földtani Közlöny (Geologische Mitteilungen)*, 73(4-9): 607-608. [Kre43b]
- KRETZOI, M. 1946. On *Bison bonusus hungarorum* n. ssp. *Annales Historico-Naturales Musei Nationalis Hungarici*, 39(5-6): 105-107. [Kre46]
- KRETZOI, M. 1955. *Dolomys* und *Ondatra*. *Acta Geologica*, 3(4): 347-355. [Kre55]
- KRETZOI, M. 1956. A Villányi hegység alsó-pleisztocén gerinces-faunái (Die altpleistozänen Wirbeltierfaunen des Villányer Gebirges). *Geologica Hungarica, Series Palaeontologica*, 27: 1-264. [Kre56]
- KRETZOI, M. 1958. New names for arvicolid homonyms. *Annales Historico-Naturales Musei Nationalis Hungarici*, 50: 55-58. [Kre58]
- KRETZOI, M. 1959. Insectivoren, Nagetiere und Lagomorphen der jüngstpliozänen Fauna von Csarnóta im Villányer Gebirge (Südungarn). *Vertebrata Hungarica*, 1(2): 237-246. [Kre59]
- KRETZOI, M. 1961. Zwei Myospalaciden aus Nordchina. *Vertebrata Hungarica*, 3(1-2): 123-136. [Kre61]
- KRETZOI, M. 1962. A csarnótai fauna és faunaszint (Fauna und Faunenhorizont von Csarnóta). *Magyar Állami Földtani Intézet Évi Jelentése (Annual Report of the Hungarian Geological Institute)*, 1959: 297-395. [Kre62]
- KRETZOI, M. 1965. Die Hipparion-Fauna von Győrszentmárton in NW-Ungarn. *Annales Historico-Naturales Musei Nationalis Hungarici*, 57: 127-143. [Kre65]
- KRETZOI, M. 1969. A magyarországi quarter és pliocén szárazföldi biosztratigráfiájának vázlata (Sketch of the Late Cenozoic (Pliocene and Quaternary) terrestrial stratigraphy of Hungary). *Földrajzi Közlemények*, 17(3): 179-204.
- KRETZOI, M. 1985. Neuer Amphicyonide aus dem Altpannon von Pécs (Südungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 77: 65-68. [Kre85]
- KRETZOI, M. & VÖRÖS, I. 1989. On a new caviomorph rodent from Peru. *Fragmenta Mineralogica et Palaeontologica*, 14: 111-116. [Kre89]

- KUBINYI, F. 1856. Emlősök és hüllők maradványai a beremendi csonttorlatban [Mammal and reptile remains from the bone breccia of Beremend]. *Magyar Akadémiai Értesítő*, 16(2): 69–77. [Kub56]
- KUTASSY, A. 1927. Beiträge zur Stratigraphie und Paläontologie der alpinen Triasschichten in der Umgebung von Budapest. *Magyar Királyi Földtani Intézet Évkönyve (Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt)*, 27(2): 107–175. [Kut27]
- KUTASSY, E. 1933. Újabb adatok a Budapest-környéki dachsteini mészkő faunájának ismeretéhez (Weitere Beiträge zur Kenntnis der Fauna des Dachstein-Kalkes in der Umgebung von Budapest). *Matematikai és Természettudományi Értesítő (Mathematischer und Naturwissenschaftlicher Anzeiger der Ungarischen Akademie der Wissenschaften)*, 49: 222–250. [Kut33]
- LAMBERT, J. & THIÉRY, P. 1911. *Essai de nomenclature raisonnée des échinides*, fasc. 3. Ferrière, Chaumont, 161–240 p. [Lam11]
- LE LOEFF, J. 1993. European titanosaurids. *Revue de Paléobiologie, Volume Spécial*, 7: 105–117. [LeL93]
- LESS, G. 1991. A Bükk felső-oligocén nagy foraminiferái (Upper Oligocene larger foraminifers of the Bükk Mountains). *Magyar Állami Földtani Intézet Évi Jelentése (Annual Report of the Hungarian Geological Institute)*, 1989: 411–465. [Les91]
- LÓCZY, L. 1877. Néhány echinoidea a Fehér-Körös völgy neogen-rétegeiből (Einige Echinoiden aus den Neogen-Ablagerungen des Weissen Körösthales). *Természetrájzi Füzetek (Naturhistorische Hefte)*, 1(1): 39–44, 61–67. [Lóc77]
- LÓCZY, L. 1897. A fosszilis emlős- és puhatestű állatmaradványok leírása és a palaeontológiai-stratigraphiai eredmények. In: *Gróf Széchenyi Béla keletázsai utjának tudományos eredménye*, vol. 3, pt. 1A–C. Kilián, Budapest, pp. 7–192. [Lóc97]
- LÓCZY, L. 1899. Beschreibung der fossilen Reste von Wirbelthieren und von Mollusken und die palaeontologisch-stratigraphischen Ergebnisse. In: *Wissenschaftliche Ergebnisse der Reise des Grafen Béla Széchenyi in Ostasien*, vol. 3, pt. 1A–C. Hölzel, Wien, pp. 11–228. [Lóc99]
- LÖRENTHEY, E. 1899. Mikroskopische Untersuchungen der palaeozoischen Gesteine. In: *Wissenschaftliche Ergebnisse der Reise des Grafen Béla Széchenyi in Ostasien*, vol. 3, pt. 1E. Hölzel, Wien, pp. 237–304. [Lör99]
- LÖRENTHEY, E. 1903. Neuere Beiträge zur tertiären Decapodenfauna Ungarns. *Mathematische und Naturwissenschaftliche Berichte aus Ungarn*, 18: 98–120. [Lör03]
- LÖRENTHEY, I. 1897. Khinai palaeozoos kőzetek mikroszkópikus vizsgálata. In: *Gróf Széchenyi Béla keletázsai utjának tudományos eredménye*, vol. 3, pt. 1E. Kilián, Budapest, pp. 201–256. [Lör97]
- LÖRENTHEY, I. 1901. Paleontológiai tanulmányok a harmadkorú rákok köréből. III. Újabb adatok Magyarország harmadkorú rákfaunájához [Paleontological studies on Tertiary decapods. III. New data on the Tertiary decapod fauna of Hungary]. *Mathematikai és Természettudományi Közlemények*, 27(5): 803–840. [Lör01]
- MAJZON, L. 1960. Magyarországi paleogén Foraminifera-szintek (Paleogene Foraminifera horizons of Hungary). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 90(3): 355–362. [Maj60]
- MAJZON, L. 1962. Hantken Miksa „Clavulina Szabói rétegek faunája” című művének nevezéktani módosítása (Nomenclatural modification of the paper “The fauna of the beds with *Clavulina Szabói*” by Miksa Hantken). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 92(7–10): 268–273. [Maj62]
- MÉHELY, L. 1914a. Fibrinae Hungariae. Die ternären und quartären wurzelzähnigen Wühlmäuse Ungarns. *Annales Historico-Naturales Musei Nationalis Hungarici*, 12: 155–243. [Méh14a]
- MÉHELY, L. 1914b. Fibrinae Hungariae. Magyarország harmad- és negyedkorú gyökeresfogú poczkai. Magyar Tudományos Akadémia Mathematikai és Természettudományi Bizottsága, Budapest, 102 p. [Méh14b]
- MÉSZÁROS, L. 1997. *Kordosia*, a new genus for some Late Miocene Amblycoptini shrews (Mammalia, Insectivora). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 2: 65–78. [Més97]
- MÉSZÁROS, L. 1999. An exceptionally rich Soricidae (Mammalia) fauna from the upper Miocene localities of Polgárdi (Hungary). *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Geologica*, 32: 5–34. [Més99]
- MEZNERICS, I. 1941. Neue Stachelhäuter (Echinodermen) aus dem Miocän Ungarns (Új magyarországi miocén tuskésbőrűek). *Annales Historico-Naturales Musei Nationalis Hungarici*, 34: 83–96. [Mez41]

- MEZNERICS, I. 1944a. Die Brachiopoden des ungarischen Tertiärs (Magyarországi harmadkori Brachiopodák). *Annales Historico-Naturales Musei Nationalis Hungarici*, 36: 10–60. [Mez44a]
- MEZNERICS, I. 1944b. Ditrupa-reste aus Ungarn. *Annales Historico-Naturales Musei Nationalis Hungarici*, 37: 40–47. [Mez44b]
- MŁYNARSKI, M. 1966. Die fossilen Schildkröten in den ungarischen Sammlungen. *Acta Zoologica Cracoviensis*, 11(8): 223–288. [Mly66]
- MOIX, P., KOZUR, H. W., STAMPFLI, G. M. & MOSTLER, H. 2007. New paleontological, biostratigraphic and paleogeographic results from the Triassic of the Mersin Mélange, SE Turkey. In: LUCAS, S. G. & SPIELMANN, J. A. (eds.), *The Global Triassic. New Mexico Museum of Natural History and Science Bulletin*, 41: 282–311. [Moi07]
- MONOSTORI, M. 1991. *Triadogigantocypris balatonica* n. g. n. sp.: a giant ostracode from the Hungarian Triassic. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1991(2): 91–96. [Mon91]
- MÜLLER, P. 1974a. Decapoda (Crustacea) fauna a budapesti miocénból (1) (Les faunes de Crustacés Décapodes des calcaires miocènes de Budapest (1)). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 104(1): 119–132. [Mül74a]
- MÜLLER, P. 1974b. Decapoda (Crustacea) fauna a budapesti miocénból (2) (Faune de Décapodes (Crustacés) du Miocène de Budapest (2)). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 104(3): 275–287. [Mül74b]
- MÜLLER, P. 1975a. Decapoda (Crustacea) fauna a budapesti miocénból (3) (Faune de Décapodes (Crustacés) du Miocène de Budapest (3)). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 105(4): 506–515. [Mül75a]
- MÜLLER, P. 1975b. *Trapezia* (Crustacea, Decapoda) a magyar eocénból és miocénból (*Trapezia* (Crustacea, Decapoda) dans l'Eocene et le Miocene de Hongrie). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 105(4): 516–523. [Mül75b]
- MÜLLER, P. 1976. Decapoda (Crustacea) fauna a budapesti miocénból (4) (Faune de Décapodes (Crustacés) dans le Miocène de Budapest (4)). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 106(2): 149–160. [Mül76]
- MÜLLER, P. 1978. Decapoda (Crustacea) fauna a budapesti miocénból (5) (Faune de Décapodes (Crustacea) dans le Miocène de Budapest (5)). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 108(3): 272–312. [Mül78]
- MÜLLER, P. 1984. A bádeni emelet tízalábú rákjai (Decapod Crustacea of the Badenian). *Geologica Hungarica, Series Palaeontologica*, 42: 1–317. [Mül84]
- MÜLLER, P. M. 2006. New decapods from the Miocene of Hungary – with remarks about their environment. *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 136(1): 37–49. [Mül06]
- MÜLLER, P. & COLLINS, J. S. H. 1991a. Late Eocene coral-associated decapods (Crustacea) from Hungary. *Contributions to Tertiary and Quaternary Geology*, 28(2–3): 47–92. [Mül91a]
- MÜLLER, P. & COLLINS, J. S. H. 1991b. *Palaeograpsus parvus* (Crustacea, Decapoda), a replacement name for *Palaeograpsus bittneri* Müller & Collins, 1991, non *Palaeograpsus bittneri* Morris & Collins, 1991. *Contributions to Tertiary and Quaternary Geology*, 28(4): 140. [Mül91b]
- NEHRING, A. 1883. Eine fossile *Siphneus*-Art (*Siphneus arvicolinus* n. sp.) aus lacustrinen Ablagerungen am oberen Hoangho. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin*, 1883: 19–24. [Neh83]
- NEHRING, A. 1898. Über *Dolomys* nov. gen. foss. *Zoologischer Anzeiger*, 21(549): 13–16. [Neh98]
- NEUGEBOREN, J. L. 1850. Die vorweltlichen Squaliden-Zähne aus dem Grobkalke bei Portsesd am Altfuß unweit Talmats. *Archiv des Vereins für Siebenbürgische Landeskunde*, 4(2): 1–44. [Neu50]
- NEUGEBOREN, J. L. 1851. Die vorweltlichen Squaliden-Zähne aus dem Grobkalke bei Portsesd am Altfuß unweit Talmats. *Archiv des Vereins für Siebenbürgische Landeskunde*, 4(3): 151–213. [Neu51]
- NOPCSA, F. 1915a. Erdély dinosaurusai. *Magyar Királyi Földtani Intézet Évkönyve*, 23(1): 1–23. [Nop15a]
- NOPCSA, F. 1915b. Die Dinosaurier siebenbürgischen Landesteile Ungarns. *Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Reichsanstalt*, 23(1): 1–24. [Nop15b]
- NOSZKY, J. 1936. Az egri felső Cattien molluszkafaunája (Die Molluskenfauna des Oberen Cattiens von Eger, in Ungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 30: 53–115. [Nos36]

- NOSZKY, J. 1939. A kiscelli agyag molluszka-faunája. I. rész. Lamellibranchiata (Die Molluskenfauna des Kisceller Tones (Rupelien) aus der Umgebung von Budapest. I. Teil. Lamellibranchiata). *Annales Historico-Naturales Musei Nationalis Hungarici*, 32: 19–146. [Nos39]
- NOSZKY, J. 1940. A kiscelli agyag molluszka-faunája. II. rész. Loricata, Gastropoda, Scaphopoda (Die Molluskenfauna des Kisceller Tones (Rupelien) aus der Umgebung von Budapest. II. Teil. Loricata, Gastropoda und Scaphopoda). *Annales Historico-Naturales Musei Nationalis Hungarici*, 33: 1–80. [Nos40]
- NYÍRÓ, M. R. 1960. Auswertung der Foraminiferen aus den transdanubischen tortonischen Beckenablagerungen. *Annales Historico-Naturales Musei Nationalis Hungarici*, 52: 33–50. [Nyí60]
- NYÍRÓ, M. R. 1961. A new foraminifer species from the Oligocene layers of Törökbálint. *Annales Historico-Naturales Musei Nationalis Hungarici*, 53: 49–50. [Nyí61]
- OZSVÁRT, P. 2003. Five new species of benthic Foraminifera from the Hungarian Paleogene Basin. *Fragmenta Palaeontologica Hungarica*, 21: 1–4. [Ozs03]
- OZSVÁRT, P. 2007. Middle and Late Eocene benthic foraminiferal fauna from the Hungarian Paleogene Basin: systematics and paleoecology. *Geologica Pannonica Special Publication*, vol. 2. Hantken Press, Budapest, 123 p. [Ozs07]
- ŐSI, A. 2005. *Hungarosaurus tormai*, a new ankylosaur (Dinosauria) from the Upper Cretaceous of Hungary. *Journal of Vertebrate Paleontology*, 25(2): 370–383. [Ősi05a]
- ŐSI, A., CLARK, J. M. & WEISHAMPEL, D. B. 2007. First report on a new basal eusuchian crocodyliform with multicusped teeth from the Upper Cretaceous (Santonian) of Hungary. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 243(2): 169–177. [Ősi07]
- ŐSI, A., WEISHAMPEL, D. B. & JIANU, C. M. 2005. First evidence of azhdarchid pterosaurs from the Late Cretaceous of Hungary. *Acta Palaeontologica Polonica*, 50(4): 777–787. [Ősi05b]
- PÁLFY, J. 2003. The Pelsonian brachiopod fauna of the Balaton Highland. In: VÖRÖS, A. (ed.), *The Pelsonian Substage on the Balaton Highland (Middle Triassic, Hungary)*. *Geologica Hungarica, Series Palaeontologica*, 55: 139–158. [Pál03]
- PAPP, K. 1907. Beschreibung der während der Forschungsreisen M. v. Déchys im Kaukasus gesammelten Versteinerungen. In: DÉCHY, M. (ed.), *Kaukasus. Reisen und Forschungen im kaukasischen Hochgebirge. Band III*. Dietrich Riemer Verlag, Berlin, pp. 141–173. [Pap07]
- PÁVAY, E. 1874a. Die fossilen Seeigel des Ofner Mergels. *Mittheilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt*, 3(2): 9–179. [Páv74a]
- PÁVAY, E. 1874b. A budai márga ásatag tüskönczei. *Magyar Királyi Földtani Intézet Évkönyve*, 3(2): 165–335. [Páv74b]
- PERVINQUIÈRE, L. 1907. *Études de paléontologie tunisienne. I. Céphalopodes des terrains secondaires*. De Rudeval, Paris, 438 p. [Per07]
- PETÉNYI, S. J. 1864. A beremendi mészkarbány természetrajz- és öslénytanilag Petényi Salamon által leírva [The natural history and paleontology of the limestone quarry at Beremend, described by Salamon Petényi]. In: KUBINYI, F. (ed.), *Petényi S. János hátrahagyott munkái [Posthumous works of János S. Petényi]*. Magyar Tudományos Akadémia, Pest, pp. 37–81. [Pet64]
- PILLER, W. E., HARZHAUSER, M. & MANDIC, O. 2007. Miocene Central Paratethys stratigraphy – current status and future directions. *Stratigraphy*, 4(2–3): 151–168.
- RABEDER, G. 1972. Die Insectivoren und Chiropteren (Mammalia) aus dem Altpaleozän von Hundsheim (Niederösterreich). *Annalen des Naturhistorischen Museums in Wien*, 76: 375–474. [Rab72]
- RABEDER, G. 1976. Die Carnivoren (Mammalia) aus dem Altpaleozän von Deutsch-Altenburg 2. Mit Beiträgen zur Systematik einiger Musteliden und Caniden. *Beiträge zur Paläontologie von Österreich*, 1: 5–119. [Rab76]
- RABEDER, G. 1981. Die Arvicoliden (Rodentia, Mammalia) aus dem Pliozän und dem älteren Pleistozän von Niederösterreich. *Beiträge zur Paläontologie von Österreich*, 8: 1–373. [Rab81]
- RABEDER, G. 1983. *Mimomys malezi* n. sp., ein neuer Arvicolide (Rodentia) aus dem Altpaleozän von Dalmatien. *Beiträge zur Paläontologie von Österreich*, 10: 1–13. [Rab83]
- RADOVANOVIC, S. 1900. Über die unterliassische Fauna von Vrška Čuka in Ostserbien. *Annales Géologiques de la Péninsule Balkanique*, 5(2): 60–70. [Rdv00]

- RADULoviĆ, V., BošKOviĆ, D. & RABRENOViĆ, D. 1997. On the Dogger (Aalenian) age of the Vrška Čuka coal overbed, eastern Serbia. *Annales Géologiques de la Péninsule Balkanique*, 61(2): 139–151. [Rdl97]
- REUMER, W. F. 1985. Ruscinian and early Pleistocene Soricidae (Insectivora, Mammalia) from Tegelen (The Netherlands) and Hungary. *Scripta Geologica*, 73: 1–173. [Rem85]
- REUSS, A. E. 1844. *Geognostische Skizzen aus Böhmen. Zweiter Band: Das Kreide-Gebirge des westlichen Böhmens*. Medau, 304 p. [Res44]
- REUSS, A. E. 1846. *Die Versteinerungen der Böhmischen Kreideformation. Zweite Abtheilung*. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 148 p. [Res46]
- REUSS, A. E. 1870. Oberoligocäne Korallen aus Ungarn. *Sitzungsberichte der Mathematisch-Naturwissenschaftliche Classe der Kaiserlichen Akademie der Wissenschaften*, 61(1): 37–56. [Res70]
- ROZLOZSNIK, P. 1924a. Bevezetés a nummulinák és assilinák tanulmányozásába. *Magyar Királyi Földtani Intézet Évkönyve*, 26(1): 1–136. [Roz24a]
- ROZLOZSNIK, P. 1924b. Nummulinák Magyarország óharmadkori rétegeiből. Néhai Hantken Miksa és Madarász Zsigmond Ede hátrahagyott tábláinak magyarázata gyanánt [Nummulites from lower Tertiary strata of Hungary. Explanation to the plates in legacy of the late Miksa Hantken and Zsigmond Ede Madarász]. *Földtani Szemle*, 1(4): 159–190. [Roz24b]
- ROZLOZSNIK, P. 1927. Einleitung in das Studium der Nummulinen und Assilinen. *Mitteilungen aus dem Jahrbuche der Königlichen Ungarischen Geologischen Anstalt*, 26(1): 1–154. [Roz27]
- ROZLOZSNIK, P. 1929. Studien über Nummulinen. *Geologica Hungarica, Series Palaeontologica*, 2: 89–248. [Roz29]
- SCHAUB, H. 1981. Nummulites et Assilines de la Téthys paléogène. Taxinomie, phylogénèse et biostratigraphie. *Schweizerische Paläontologische Abhandlungen*, 104–106: 1–238. [Sch81]
- SCHAUB, S. 1930. Quartäre und jungtertiäre Hamster. *Abhandlungen der Schweizerischen Paläontologischen Gesellschaft*, 49: 1–49. [ScS30]
- SCHLESINGER, G. 1917. Die Mastodonten des K. K. Naturhistorischen Hofmuseums. Morphologisch-phylogenetische Untersuchungen. *Denkschriften des K. K. Naturhistorischen Hofmuseums*, 1: 1–230. [ScG17]
- SCHLESINGER, G. 1922. Die Mastodonten der Budapester Sammlungen. (Untersuchungen über Morphologie, Phylogenie, Ethologie und Stratigraphie europäischer Mastodonten.) *Geologica Hungarica*, 2(1): 1–284. [ScG22]
- SCHLOSSER, M. 1906. Über fossile Land- und Süßwassergastropoden aus Centralasien und China. *Annales Historico-Naturales Musei Nationalis Hungarici*, 4: 372–405. [ScM06]
- SCHRÉTER, Z. 1929. A borsod-hevesi szén és lignitterületek bányaföldtani leírása [Geology of the Borsod-Heves coal and lignite mining district]. *Magyar Királyi Földtani Intézet Kiadványai*, 1929: 5–390. [ScZ29]
- SCHRÉTER, Z. 1975. Tanulmány az alsópleisztocén korú Melanopsidák köréből [Studies on Lower Pleistocene Melanopsidae]. *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 105(1): 1–22. [ScZ75]
- SCHREUDER, A. 1940. A revision of the fossil water-moles (Desmaninae). *Archives Néerlandaises de Zoologie*, 4(2–3): 201–333. [ScA40]
- SPASSOV, N. & GERAADS, D. 2004. *Tragopontax Pilgrim*, 1937 and *Miotragocerus Stromer*, 1928 (Mammalia, Bovidae) from the Turolian of Hadjidimovo, Bulgaria, and a revision of the Late Miocene Mediterranean Boselaphini. *Geodiversitas*, 26(2): 339–370. [Spa04]
- STRAUSZ, L. 1940. Die pannonische Molluskenfauna der Tiefbohrung von Magyarszentmiklós (A magyarszentmiklói mélyfúrás pannonkorai molluszkái). *Annales Historico-Naturales Musei Nationalis Hungarici*, 33: 81–86. [Str40]
- STRAUSZ, L. 1942. Das Pannon des mittleren Westungarns (A Dunántúl középső részének pannón-kori rétegei). *Annales Historico-Naturales Musei Nationalis Hungarici*, 35: 1–102. [Str42]
- STRAUSZ, L. 1950. Óslénytani adatok Baranyából (Deux faunes miocènes de la Montagne Mecsek, Hongrie). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 80(7–9): 238–246. [Str50]
- STRAUSZ, L. 1954. Várpalotai felső-mediterrán csigák (Les Gastropodes du méditerranéen supérieur (Tortonien) de Várpalota). *Geologica Hungarica, Series Palaeontologica*, 25: 1–150. [Str54]

- STRAUSZ, L. 1955. Cerithium-félék a Dunántúl középső-miocén rétegeiből (Mittelmiozäne Cerithien Transdanubiens). *Magyar Állami Földtani Intézet Évkönyve (Annals of the Hungarian Geological Institute)*, 43(1): 3–271. [Str55]
- SZABÓ, J. 1996. Some new species of *Adeorbisina* (Adeorbisininae, Colloniidae, Trochoidea) in the Bajocian gastropod fauna of Somhegy (Bakony Mts., Hungary). *Fragmenta Mineralogica et Palaeontologica*, 18: 63–70. [Szb96]
- SZABÓ, J. 2007. Initial notes to a revision of Late Triassic gastropods from Budapest (Hungary): *Hungariella* Kutassy, 1933 (Neritopsidae). *Fragmenta Palaeontologica Hungarica*, 24–25: 69–75. [Szb07]
- SZALAI, T. 1926a. A várpalotai középmiocén faunája (Die Mittelmiozäne Fauna von Várpalota). *Annales Historico-Naturales Musei Nationalis Hungarici*, 24: 331–347. [Sz126a]
- SZALAI, T. 1926b. Adatok a harmadkori Crinoideák kérdéséhez (Daten zur Frage der Tertiärcrinoiden). *Földtani Közlöny (Geologische Mitteilungen)*, 55: 169–174, 339–341. [Sz126b]
- SZALAI, T. 1930. Biomorphische und methodologisch-systematische Untersuchungen an rezenten und fossilen Testudinaten. *Palaeobiologica*, 3: 347–364. [Sz130]
- SZALAI, T. 1934. Die fossilen Schildkröten Ungarns. *Folia Zoologica et Hydrobiologica*, 6(2): 97–142. [Sz134]
- SZENTE, I. 1995. Bivalvia from the Bathonian (Middle Jurassic) of the Mecsek Mts, Hungary. *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Geologica*, 30: 93–109, 221–224. [Sze95]
- SZIVES, O. 2007a. Aptian stage. In: SZIVES, O. (ed.), *Aptian–Campanian ammonites of Hungary. Geologica Hungarica, Series Palaeontologica*, 57: 31–74. [Szi07a]
- SZIVES, O. 2007b. Albian stage. In: SZIVES, O. (ed.), *Aptian–Campanian ammonites of Hungary. Geologica Hungarica, Series Palaeontologica*, 57: 75–122. [Szi07b]
- SZIVES, O. & MONKS, N. 2002. Heteromorph ammonites from the Tata Limestone Formation (Aptian-Lower Albian), Hungary. *Palaeontology*, 45(6): 1137–1149. [Szi02]
- SZÖRÉNYI, E. 1929. A budai márga és faunája [The Buda Marl and its fauna]. *Földtani Szemle Melléklete*, 1929: 1–44. [Szö29]
- SZÖRÉNYI, E. 1933. Adatok a harmadkori *Sepia*-félék ismeretéhez, néhány új magyarországi faj alapján (Neue tertiäre Sepiinae aus Ungarn nebst Bemerkungen zum zeitlichen Auftreten und zur Entwicklung der Gattung *Sepia*). *Földtani Közlöny (Geologische Mitteilungen)*, 63(7–12): 183–189. [Szö33]
- SZÖRÉNYI, E. 1934. Oligocén *Scalpellum* maradványok Magyarországról (Scalpellumreste aus dem ungarischen Oligozän). *Földtani Közlöny (Geologische Mitteilungen)*, 64(10–12): 273–277. [Szö34]
- SZÖTS, E. 1938. A móri Antalhegy óharmadkori képződményei (Die paläogenen Bildungen des Antalberges). *Földtani Szemle Melléklete*, 1938: 1–42. [Szö38]
- SZÖTS, E. 1953. Magyarország eocén puhatestű I. Gántkörnyéki eocén puhatestűek (Mollusques éocènes de la Hongrie I. Les mollusques éocènes des environs de Gánt). *Geologica Hungarica, Series Palaeontologica*, 22: 1–270. [Szö53]
- SZTRÁKOS, K. 1973. Révision des espèces "*Globigerina*" *applanata* et "*Globigerina*" *globosa* décrites par M. Hantken d'Euganea (Italie). *Revue de Micropaléontologie*, 16(4): 224–228. [Szt73]
- SZTRÁKOS, K. 1974. Paleogene planktonic foraminiferal zones in northeastern Hungary. *Fragmenta Mineralogica et Palaeontologica*, 5: 29–80. [Szt74]
- SZTRÁKOS, K. 1979. La stratigraphie, paléoécologie, paléogéographie et les foraminifères de l'Oligocène du nord-est de la Hongrie. *Cahiers de Micropaléontologie*, 3: 1–95. [Szt79]
- TASNÁDI KUBACSKA, A. 1939. *Trilophodon angustidens* Cuv. forma *praetypica* koponyamaradványa Zagyvapálfalváról (Ein primitiver Vertreter des Formenkreises *Trilophodon angustidens* Cuv. aus Ungarn). *Annales Historico-Naturales Musei Nationalis Hungarici*, 32: 154–164. [Tas39]
- TASNÁDI-KUBACSKA, A. & SOÓS, L. 1935. Die Mollusken- und Wirbeltierfauna des Plistozän und Ober-Pliozän von Gombaszög. *Annales Historico-Naturales Musei Nationalis Hungarici*, 29: 9–20. [Tas35]
- TELEGDI ROTH, K. 1914a. Eine oberoligozäne Fauna aus Ungarn. *Geologica Hungarica*, 1(1): 1–77. [Tel14a]
- TELEGDI ROTH, K. 1914b. Felső-oligocén fauna Magyarországból. *Geologica Hungarica*, 1(1): 1–66. [Tel14b]

- TOMOR-THIRRING, J. 1936. Őslénytani újdonságok a Bakonyhegységből (Paläontologische Neuigkeiten aus dem Bakony-Gebirge). *Földtani Közlöny (Geologische Mitteilungen)*, 66(1-3): 51–68. [Tom36]
- TOPÁL, G. 1963a. The bats of a Lower Pleistocene site from Mt. Kövesvárad near Répáshuta, Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici*, 55: 143–154. [Top63a]
- TOPÁL, G. 1963b. Description of a new bat, *Rhinolophus macrorhinus* sp. n. from the Lower Pleistocene of Hungary. *Vertebrata Hungarica*, 5(1-2): 219–228. [Top63b]
- TOPÁL, G. 1970. *Barbastella rostrata* n. sp. a Tarkói kőfülke középső pleisztocénjéből (*Barbastella rostrata* n. sp. from the Middle Pleistocene of the Tarkő niche, North-East Hungary). *Őslénytani Viták (Discussiones Palaeontologicae)*, 15: 5–18. [Top70]
- TOPÁL, G. 1974. The first record of *Megaderma* in Hungary (Pliocene sediments of Osztramos, Locality 10). *Vertebrata Hungarica*, 15: 95–104. [Top74]
- TOPÁL, G. 1975. A new fossil horseshoe bat (*Rhinolophus variabilis* n. sp.) from the Pliocene sediments of the Osztramos Hill, NE Hungary (Mammalia: Chiroptera). *Fragmenta Mineralogica et Palaeontologica*, 6: 5–29. [Top75]
- TOPÁL, G. 1979. Fossil bats of the *Rhinolophus ferrumequinum* group in Hungary (Mammalia: Chiroptera). *Fragmenta Mineralogica et Palaeontologica*, 9: 61–101. [Top79]
- TOPÁL, G. 1981. New fossil mouse-eared bat, *Myotis kretzoi* sp. n., from the Middle Pleistocene of Hungary (Mammalia: Chiroptera). *Fragmenta Mineralogica et Palaeontologica*, 10: 59–64. [Top81]
- TOPÁL, G. 1983. New and rare fossil mouse-eared bats from the Middle Pliocene of Hungary (Mammalia, Chiroptera). *Fragmenta Mineralogica et Palaeontologica*, 11: 43–54. [Top83]
- TOPÁL, G. 1989. New Tertiary plecotines from Hungary (Mammalia, Chiroptera). In: HANÁK, V., HORÁČEK, I. & GAISLER, J. (eds.), *European Bat Research 1987*. Charles University Press, Praha, pp. 77–86. [Top89]
- VADÁSZ, M. E. 1914. Magyarország mediterrán tuskésbőrűi. *Geologica Hungarica*, 1(2): 69–227. [Vad14]
- VADÁSZ, M. E. 1915. Die mediterranen Echinodermen Ungarns. *Geologica Hungarica*, 1(2): 79–253. [Vad15]
- VÉGH, S. 1964. A bakonyi fődolomit rétegtani kérdései (Stratigraphische Fragen des Hauptdolomits im Bakonygebirge). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 94(3): 327–339. [Vég64]
- VÉGH-NEUBRANDT, E. 1963. Nóri dachsteini mészkő az Északi Bakonyban (Norischer Dachsteinkalk im Nord-Bakony). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 93(3): 332–340. [V-N63]
- VÉGH-NEUBRANDT, E. 1969. Bemerkungen zur Gattung *Paramegalodus* und Aufstellung des neuen Genus *Rhaetomegalodon*. *Anzeiger der Mathematisch-Naturwissenschaftlichen Klasse der Österreichischen Akademie der Wissenschaften*, 1969(6): 121–125. [V-N69]
- VÉGH-NEUBRANDT, E. 1974. Új Megalodontacea fajok a magyarországi felsőtriaszból (Neue Megalodontacea-Arten aus der Obertrias von Ungarn). *Földtani Közlöny (Bulletin of the Hungarian Geological Society)*, 104(4): 10–39. [V-N74]
- VÉGH-NEUBRANDT, E. 1982. *Triassische Megalodontaceae. Entwicklung, Stratigraphie und Paläontologie*. Akadémiai Kiadó, Budapest, 526 p. [V-N82]
- VÉGH-NEUBRANDT, E., DUMONT, J.-F., GUTNIC, M., MARCOUX, J., MONOD, O. & POISSON, A. 1976. Megalodontidae du Trias supérieur dans la chaîne taurique (Turquie méridionale). *Géobios*, 9(2): 199–222. [V-N76]
- VIGH, G. 1914. Adatok az esztergomvidéki triász ismeretéhez (Beiträge zur Kenntnis der Trias in Komitate Esztergom). *Földtani Közlöny (Geologische Mitteilungen)*, 44(10-12): 572–577, 599–604. [Vig14]
- VITÁLIS, I. 1934. A *Congeria soproniensis* n. sp. (*Congeria soproniensis* n. sp.). *Matematikai és Természettudományi Értesítő (Mathematischer und Naturwissenschaftlicher Anzeiger der Ungarischen Akademie der Wissenschaften)*, 50: 509–519. [Vit34]
- VÖRÖS, A. 1983. Some new genera of Brachiopoda from the Mediterranean Jurassic. *Annales Historico-Naturales Musei Nationalis Hungarici*, 75: 5–25. [VöA83]

- VÖRÖS, A. 1995. Bathonian brachiopods of the Mecsek Mts (Hungary). *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio Geologica*, 30: 181–208, 237–238. [VöA95]
- VÖRÖS, A. 2001. Middle Triassic (Anisian) nautilid cephalopods from Aszófő (Balaton Highland, Hungary). *Fragmenta Palaeontologica Hungarica*, 19: 1–14. [VöA01]
- VÖRÖS, I. 1979. *Archidiskodon meridionalis* ürömensis n. ssp. from the Lower Pleistocene of the Carpathian Basin. *Fragmenta Mineralogica et Palaeontologica*, 9: 5–8. [VöI79]
- VÖRÖS, I. 1981. The skulls of mammoth in Hungary. *Fragmenta Mineralogica et Palaeontologica*, 10: 97–106. [VöI81]
- VÖRÖS, I. 1985. *Alces brevirostris* Kretzoi from the Ördöglyuk Cave at Solymár (Hungary). *Fragmenta Mineralogica et Palaeontologica*, 12: 59–66. [VöI85]
- VÖRÖS, I. 1989. *Prodeinotherium petenyii* sp. n. from the Lower Miocene at Putnok (North Hungary). *Fragmenta Mineralogica et Palaeontologica*, 14: 101–110. [VöI89]
- WAGNER, J. 1938. A kiscelli közép-oligocén (Rupélien) rétegek kétkopolyús Cephalopodái és új *Sepia*-félék a magyar eocénból (Die dibranchiaten Cephalopoden der mitteloligozänen (Rupélien) Tonschichten von Kiscell und eine Sepiinae aus dem ungarischen Eozän). *Annales Historico-Naturales Musei Nationalis Hungarici*, 31: 179–199. [Wag38]
- WEILER, W. 1933. Két magyarországi oligocénkorú halfauna (Zwei oligozäne Fischfaunen aus dem Königreich Ungarn). *Geologica Hungarica, Series Palaeontologica*, 11: 1–54. [Wei33]
- WEILER, W. 1935. *Nemopteryx kubacskei* n. sp. aus dem Kleinzeller Tegel bei Budapest, zugleich ein Beitrag zur Geschichte der Gattungen *Nemopteryx* Ag. und *Merluccius* L. *Palaeontologische Zeitschrift*, 17(1–2): 27–44. [Wei35]
- WEILER, W. 1938. Neue Untersuchungen an mitteloligozänen Fischen Ungarns. *Geologica Hungarica, Series Palaeontologica*, 15: 1–31. [Wei38]
- ZORN, I. & BOHN-HAVAS, M. 1997. Revision of Hungarian Tertiary holoplanktonic gastropods housed in the collections of the Hungarian Museum of Natural History in Budapest. In: DUDICH, E. & LOBITZER, H. (eds.), *Advances in Austrian-Hungarian Joint Geological Research*. Geological Institute of Hungary, Budapest, pp. 83–95. [Zor97]
- ZORN, I. & JANSEN, A. W. 1993. *Vaginella tricuspidata* sp. nov., a new holoplanktonic mollusc from the Late Oligocene of SW France and Hungary. *Contributions to Tertiary and Quaternary Geology*, 29(3–4): 61–71. [Zor93]

Type specimens are the scientifically most valuable assets in the collections of the Department of Geology and Paleontology of the Hungarian Natural History Museum. To make them more accessible to the wider paleontological community for research, a comprehensive type catalogue has been compiled for the first time. A listing of more than 2000 inventory lots with preserved and available type specimens comprises the bulk of the volume. In addition, lost or missing types are also included, many of them were destroyed in a devastating fire in 1956. Moving the collections to a new location in 2004–2006 and the development of digital collection databases created both a necessity and an opportunity of cataloguing the type specimens in our holding. Currently the collections contain the name-bearing types of 917 fossil invertebrate and vertebrate species and subspecies. Complemented by those with secondary or lost types, a total of 1181 species-group taxa are recorded in the catalogue which was compiled by a team of curators and research scientists.

